

ABBREVIATED NOTICE OF RESOURCE AREA DELINEATION

Filing Under the Massachusetts Wetlands Protection Act M.G.L. Chapter 131, Section 40 and the Town of Shutesbury Wetland Bylaw

Pratt South Project Pratt Corner Road Shutesbury, Massachusetts

Submitted to:

Shutesbury Conservation Commission

Shutesbury Town Hall 1 Cooleyville Road Shutesbury, Massachusetts 01072

Filed by:

W.D. Cowls, Inc.

134 Montague Road, P.O. Box 9677 North Amherst, Massachusetts 01059

Prepared by:

TRC Companies

650 Suffolk Street Lowell, Massachusetts 01854

October 2020



October 26, 2020

Town of Shutesbury Conservation Commission Shutesbury Town Hall 1 Cooleyville Road Shutesbury, MA 01072

RE: Pratt South Project
Pratt Corner Road
Abbreviated Notice of Resource Area Delineation (ANRAD)

Dear Commissioners:

TRC Companies (TRC) is writing on behalf of W.D. Cowls, Inc. to file an ANRAD for a parcel off Pratt Corner Road, Shutesbury, MA (Site) (Figure 1 in Attachment B). The Site consists of approximately 92.6 acres of a 140.18-acre parcel (listed by the Shutesbury tax assessor as Parcel ID ZU-2).

TRC conducted a wetland and waterbody delineation survey on July 29 and 30 and August 3, 2020. This survey resulted in an overall delineation of five wetlands and two streams. The total linear feet of wetland edge and other resource areas delineated during the wetland and waterbody survey effort for the Site, the focus of this ANRAD filing, are summarized in the following table:

Resource Area	Delineated Length (linear feet)
Bordering Vegetated Wetland	8,663
Bank	2,736

Please refer to Attachment B for survey methodology, delineated wetland descriptions, US Army Corps of Engineers Wetland Determination forms, site photographs, and figures showing the resource areas.

To assist your review, we have provided the following attachments:

- 1. Attachment A Abbreviated Notice of Resource Area Delineation Form & Wetland Fee Transmittal Form
- 2. Attachment B Wetland and Waterbody Delineation Report
- 3. Attachment C Abutter Information (Certified Abutter List, Abutter Notification & Affidavit of Service)
- 4. Attachment D Figure 1: Delineated Resources Map (September 2020)

Attachment B also includes the following figures:

Figure 1 – Project Location (September 2020)

Figure 2 – Wetland Delineation (September 2020)

We very much appreciate your review of this information. If you should have any questions, please do not hesitate to contact me at 978-656-3662 or via email at JBrandt@TRCcompanies.com.

Sincerely,

TRC Companies

Jeff Brandt

Senior Project Manager

Brandt



ATTACHMENT A Abbreviated Notice of Resource Area Delineation Form & Wetland Fee Transmittal Form





Important: When filling out forms on the computer, use only the tab key to move your cursor - do not use the return

key.

Note: Before completing this form consult your

local Conservation Commission regarding any municipal bylaw or ordinance.

Massachusetts Department of Environmental ProtectionBureau of Resource Protection - Wetlands

WPA Form 4A – Abbreviated Notice of Resource Area Delineation

Massachusetts Wetlands Protection Act M.G.L. c. 131, §40

⊃rov	ided by MassDEP:
	MassDEP File Number
	Document Transaction Number
	Shutesbury City/Town

A. General Information

Pratt Corner Road	Shutesbury	01072
a. Street Address	b. City/Town	c. Zip Code
Latituda and Langituda.	42.41192	-72.46679
Latitude and Longitude:	d. Latitude	e. Longitude
Map ZU	Lot 2	
f. Assessors Map/Plat Number	g. Parcel /Lot Number	
Applicant:		
a. First Name	b. Last Name	
W.D. Cowls, Inc.		
c. Organization		
P.O. Box 9677		
d. Mailing Address		
North Amherst	MA	01059
e. City/Town	f. State	g. Zip Code
h. Phone Number i. Fax Number	eturner@ariespowersy	ystems.com
h. Phone Number i. Fax Number	j. Email Address	
Property owner (if different from applicant):		han one owner (attach additic nd contact information)
a. First Name	b. Last Name	
c. Organization		
d. Mailing Address		
e. City/Town	f. State	g. Zip Code
h. Phone Number i. Fax Number	j. Email Address	
Representative (if any):		
Jeff	Brandt	
	h Camtaat Damaan Laat Nam	20
a. Contact Person First Name	 b. Contact Person Last Nan 	IE
	b. Contact Person Last Nan	ie
a. Contact Person First Name	D. Contact Person Last Nan	ie
a. Contact Person First Name TRC	D. Contact Person Last Nan	ie
a. Contact Person First Name TRC c. Organization	b. Contact Person Last Nan	ie
a. Contact Person First Name TRC c. Organization 650 Suffolk Street d. Mailing Address Lowell	MA	01854
a. Contact Person First Name TRC c. Organization 650 Suffolk Street d. Mailing Address		
a. Contact Person First Name TRC c. Organization 650 Suffolk Street d. Mailing Address Lowell	MA	01854 g. Zip Code

Fees will be calculated for online users.

a. Total Fee Paid

wpaform4a.doc • rev. 12/11 Page 1 of 4

b. State Fee Paid

c. City/Town Fee Paid



Massachusetts Department of Environmental ProtectionBureau of Resource Protection - Wetlands

WPA Form 4A – Abbreviated Notice of Resource Area Delineation

Massachusetts Wetlands Protection Act M.G.L. c. 131, §40

Provid	ded by MassDEP:
N	MassDEP File Number
Ī	Document Transaction Number
5	Shutesbury
(Citv/Town

B. Area(s) Delineated

	` '						
1.	Bordering \	Vegetated Wetland (BVW)	8,663 Linear Feet of Boundary Delineated				
2.	Check all n	nethods used to delineate the Border	ing Vegetated Wetland (B	√W) boundary:			
	a. MassDEP BVW Field Data Form (attached)						
	b. 🛛 Otl	her Methods for Determining the BVV	V boundary (attach docum	entation):			
	1. 🖂	50% or more wetland indicator plan	ts				
	2.	Saturated/inundated conditions exis	st				
	3.	Groundwater indicators					
	4. Direct observation						
	5. 🔀	Hydric soil indicators					
	6. Credible evidence of conditions prior to disturbance						
3.	Indicate an	y other resource area boundaries tha	at are delineated:				
Ва	nk			2,736			
	Resource Area			b. Linear Feet Delineated			
c. F	Resource Area			d. Linear Feet Delineated			

C. Additional Information

Applicants must include the following plans with this Abbreviated Notice of Resource Area Delineation. See instructions for details. **Online Users:** Attach the Document Transaction Number (provided on your receipt page) for any of the following information you submit to the Department.

- 1. ANRAD (Delineation Plans only)
- 2. SGS or other map of the area (along with a narrative description, if necessary) containing sufficient information for the Conservation Commission and the Department to locate the site. (Electronic filers may omit this item.)
- 3. Plans identifying the boundaries of the Bordering Vegetated Wetlands (BVW) (and/or other resource areas, if applicable).
- 4. \boxtimes List the titles and final revision dates for all plans and other materials submitted with this Abbreviated Notice of Resource Area Delineation.

wpaform4a.doc • rev. 12/11 Page 2 of 4



Massachusetts Department of Environmental ProtectionBureau of Resource Protection - Wetlands

WPA Form 4A – Abbreviated Notice of Resource Area Delineation

Massachusetts Wetlands Protection Act M.G.L. c. 131, §40

Prov	ided by MassDEP:
	MassDEP File Number
	Document Transaction Number
	Shutesbury
	Citv/Town

D. Fees

calculated and submitted to the 0 Wetland Fee Transmittal Form).	Conservation Commission and the Department (see Instructions and
	hall be assessed for projects of any city, town, county, or district of cognized Indian tribe housing authority, municipal housing authority, portation Authority.
Applicants must submit the follov Form) to confirm fee payment:	ving information (in addition to the attached Wetland Fee Transmittal
4005000	Contourle on 44, 2000

The fees for work proposed under each Abbreviated Notice of Resource Area Delineation must be

1205026	September 14, 2020		
2. Municipal Check Number	3. Check date		
1205034	September 14, 2020		
4. State Check Number	5. Check date		
TRC			
6. Payor name on check: First Name	7. Payor name on check: Last Name		

wpaform4a.doc • rev. 12/11 Page 3 of 4



Massachusetts Department of Environmental Protection Bureau of Resource Protection - Wetlands

WPA Form 4A – Abbreviated Notice of Resource Area Delineation

Massachusetts Wetlands Protection Act M.G.L. c. 131, §40

Provided by MassDEP:

MassDEP File Number

Document Transaction Number

Shutesbury City/Town

E. Signatures

I certify under the penalties of perjury that the foregoing Abbreviated Notice of Resource Area Delineation and accompanying plans, documents, and supporting data are true and complete to the best of my knowledge. I understand that the Conservation Commission will place notification of this Notice in a local newspaper at the expense of the applicant in accordance with the wetlands regulations, 310 CMR 10.05(5)(a).

I further certify under penalties of perjury that all abutters were notified of this application, pursuant to the requirements of M.G.L. c. 131, § 40. Notice must be made in writing by hand delivery or certified mail (return receipt requested) to all abutters within 100 feet of the property line of the project location.

I hereby grant permission, to the Agent or member of the Conservation Commission and the Department of Environmental Protection, to enter and inspect the area subject to this Notice at reasonable hours to evaluate the wetland resource boundaries subject to this Notice, and to require the submittal of any data deemed necessary by the Conservation Commission or Department for that evaluation.

I acknowledge that failure to comply with these certification requirements is grounds for the Conservation Commission or the Department to take enforcement action.

1. Signature of Applicant

3. Signature of Property Owner (if different)

5. Signature of Representative (if any)

2. Date

4. Date

6. Date

For Conservation Commission:

Two copies of the completed Abbreviated Notice of Resource Area Delineation (Form 4A), including supporting plans and documents; two copies of the ANRAD Wetland Fee Transmittal Form; and the city/town fee payment must be sent to the Conservation Commission by certified mail or hand delivery.

For MassDEP:

One copy of the completed Abbreviated Notice of Resource Area Delineation (Form 4A), including supporting plans and documents; one copy of the ANRAD Wetland Fee Transmittal Form; and a copy of the state fee payment must be sent to the MassDEP Regional Office (see Instructions) by certified mail or hand delivery. (E-filers may submit these electronically.)

The original and copies must be sent simultaneously. Failure by the applicant to send copies in a timely manner may result in dismissal of the Notice of Intent.

Important: When filling out forms on the computer, use only the tab key to move your cursor do not use the



return key.



☐ Online users: check box if fee exempt.

Massachusetts Department of Environmental ProtectionBureau of Resource Protection - Wetlands

ANRAD Wetland Fee Transmittal Form

Massachusetts Wetlands Protection Act M.G.L. c. 131, §40

Α.	App	licant Inform	nation						
1.	Location of Project:								
	Dratt (Corner Road (Parc	al ID: 7H-2\	Shutesbury					
		t Address	er ib. 20-2)	b. City/Town					
	\$987.5			1205034					
	c. Fee a			d. Check number	er				
	5 55 5				•				
2.	Applic	ant:							
					W.D. Cowls, Inc.				
	a. First	Name	b. Last Nam	e	c. Company				
	P.O. E	Box 9677			. ,				
		ng Address							
	North	Amherst			MA	01059			
	e. City/7				f. State	g. Zip Code			
		14-1702							
	h. Phon	e Number							
3.	Prone	rty Owner (if differ	ent).						
Ο.	Порс	ity Owner (ii diner	one).						
	a. First	Name	b. Last Nam	e	c. Company				
	d. Mailir	ng Address							
	e. City/1	Town			f. State	g. Zip Code			
	h. Phon	e Number							
R	Fees	2							
υ.	1 663	•							
app Are	olicable	project type). The	maximum fee for ea	ce Area Delineation i ach ANRAD, regardle vith a single-family h	ess of the num	ber of Resource			
	Borde	ring Vegetated We	tland Delineation Fe	ee:					
	1. 🗌	single family	a. feet of BVW	x \$2.00 =	b Foot	for DVAA			
	- 🖂	house project				for BVW			
	2. 🛛	all other	8,663	\$17,326		0 (maximum fee) for BVW			
		projects	a. feet of BVW	x \$2.00 =	D. Fee	IOL BAAA			
	Other	Resource Area (e.	g., bank, riverfront a	rea, etc.):					
	3. 🗌	single family							
		house project	a. linear feet	x \$2.00 =	b. Fee				
	4. 🛛	all other	2,736	\$5,472	\$0 (m	aximum fee)			
		projects	a. linear feet	x \$2.00 =	b. Fee				
			Total Fe	e for all Resource Ar	eas: $\frac{$2,000}{\text{Fee}}$	0			
				State share of filing	fee: $\frac{$987.5}{5.1/2}$	50 If total fee less \$12.50			
					\$1.01				
			City	/Town share of filing	fee: $\frac{\Psi^{1,01}}{6.4/2}$	f total foo plus \$12.50			

6. 1/2 of total fee **plus** \$12.50



Massachusetts Department of Environmental Protection

Bureau of Resource Protection - Wetlands

ANRAD Wetland Fee Transmittal Form

Massachusetts Wetlands Protection Act M.G.L. c. 131, §40

C. Submittal Requirements

a.) Send a copy of this form, with a check or money order for the state share of the fee, payable to the Commonwealth of Massachusetts, to:

Department of Environmental Protection Box 4062 Boston, MA 02211

- b.) **To the Conservation Commission:** Send the Abbreviated Notice of Resource Area Delineation; a **copy** of this form; and the city/town fee payment.
- c.) **To DEP Regional Office**: Send one copy of the Abbreviated Notice of Resource Area Delineation (and any additional documentation required as part of a Simplified Review Buffer Zone Project); a **copy** of this form; and a **copy** of the state fee payment. (E-filers of Notices of Intent may submit these electronically.)



Citizens Bank CONNECTICUT 51-7011/2111

CHECK DATE

September 14, 2020

Security Check Features Included Details on Back

PAY Nine Hundred Eighty Seven and 50/100 Dollars

AMOUNT

PAY TO THE ORDER OF

Commonwealth Of Massachusetts

\$ 987.50

Department of Environmental Protection

P.O. Box 4062 Boston, MA 02211

VOID AFTER 90 DAYS

AUTHORIZED SIGNATURE

#1205034# #211170114# 2232037104#



21 Griffin Road North Windsor, CT 06095 860.289.9692

EMILY BUSINESS FORMS 800.392.6018 DELTEK VISION

1205034

Check Date: 9/14/2020

Invoice Number	Date	Voucher	Amount	Discounts	Previous Pay	Net Amount
WPA 4A-SHUTESBURY 1	9/11/2020	007756821557	987.50	annimate-manda-mand-libration in the a		987.50
Commonwealth Of Massachusetts TOTAL		987.50			987.50	
Citizen Bank - Disbursement	1	030812				



Citizens Bank CONNECTICUT 51-7011/2111

CHECK DATE

September 14, 2020

PAY One Thousand Twelve and 50/100 Dollars

AMOUNT

PAY TO THE ORDER OF Town of Shutesbury

\$ 1,012.50

1 Cooleyville Road PO BOX 276

Shutesbury, MA 01072

VOID AFTER 90 DAYS

AUTHORIZED SIGNATURE

"1205026" "211170114" 2232037104"



21 Griffin Road North Windsor, CT 06095 860.289.9692

1205026

Invoice Number	Date	Voucher	Amount	Discounts	Previous Pay	Net Amount
WPA 4A FILING-PRATT	9/11/2020	007756821568	1,012.50			1,012.50
Town of Shutesbury		TOTAL	1,012.50			1,012.50
Citizen Bank - Disbursement	9	123516				

Check Date: 9/14/2020

ATTACHMENT B Wetland and Waterbody Delineation Report





Pratt South Project

Pratt Corner Road Shutesbury, Massachusetts

Prepared By:

TRC Wannalancit Mills 650 Suffolk Street Lowell, Massachusetts 01854

Wetland and Waterbody Delineation Report

August 2020



TABLE OF CONTENTS

1.0	INTR	RODUCTION					
2.0	REGI	JLATOR	Y AUTHORITY	1			
	2.1	United	States Army Corps of Engineers	1			
	2.2	Massa	achusetts Department of Environmental Protection	2			
	2.3	Town	of Shutesbury Conservation Commission	3			
3.0	PRO	PROJECT SITE CHARACTERISTICS					
	3.1	Hydrology					
		3.1.1	Floodplains	4			
	3.2	Federa	al and State Mapped Wetlands and Streams	4			
	3.3	Марре	ed Soils	4			
		3.3.1	Hydric Rating	5			
		3.3.2	Natural Drainage Class	6			
		3.3.3	Prime Farmland	6			
		3.3.4	Hydrologic Soil Groups	6			
4.0	WETI	LAND AN	ND STREAM DELINEATION METHODOLOGY	7			
	4.1	Non-wetland Aquatic Resource Methodology					
	4.2	Wetland Delineation Methodologies					
		4.2.1	Hydrophytic Vegetation Methodologies	8			
		4.2.2	Hydric Soil Methodologies	9			
		4.2.3	Wetland Hydrology Methodologies	9			
5.0	RESU	JLTS		9			
	5.1	Upland Areas					
	5.2	Delineated Wetlands and Waterbodies					
		5.2.1	Delineated Wetlands	10			
		5.2.2	Delineated Waterbodies	11			
6.0	CON	CLUSION	NS	12			
7.0	REFE	RENCES	S	13			



TABLES

Table 1: Mapped Soils					
APPENDIC	ES				
Appendix A	Figures				
Figure	e 1. Site Location Map				
Figure	e 2. Wetland Delineation				
Appendix B	Photographs				
Appendix C	Wetland Determination Data Forms				
Appendix D	NRCS Soil Report				
Appendix E	USGS StreamStats Report				



1.0 Introduction

This report presents the results of a wetland and waterbody delineation conducted on July 29, 30, 2020 and August 3, 2020 by TRC Companies, Inc. (TRC) south of Pratt Corner Road in the Town of Shutesbury, Franklin County, Massachusetts (Site). The survey included approximately 92.6 acres of the 140.18-acre parcel listed by the Shutesbury Tax Assessor as Parcel ID ZU-2.

The survey for wetlands and streams focused on the entire Site as well as adjacent parcels, when accessible, within 200 feet.

This report documents wetlands, streams, and other aquatic resources (ponds, lakes, impoundments, etc.) at the Site regardless of assumed jurisdictional status and addresses the implementation of local and state regulated buffer areas. To the extent practicable, the delineated resources were investigated to determine drainage patterns and a physical nexus to Waters of the United States (WOUS).

Appendix A provides a Site location map (Figure 1) and a map of the resources delineated by TRC (Figure 2). Appendix B includes representative photographs of the Site, Appendix C includes wetland determination data forms, and Appendix D contains the Natural Resources Conservation Service (NRCS) Soil Report. Appendix E contains the U.S. Geological Survey (USGS) StreamStats Report.

2.0 Regulatory Authority

2.1 United States Army Corps of Engineers

In accordance with Section 404 of the Clean Water Act (CWA), the United States Army Corps of Engineers (USACE) asserts jurisdiction over WOUS, defined as wetlands, streams, and other aquatic resources under the regulatory authority per Title 33 Code of Federal Regulations (CFR) Part 328, and the United States Environmental Protection Agency (EPA) per Title 40 CFR Part 230.3(s). Wetlands are defined as "those areas that are inundated or saturated by surface or ground water at a frequency and duration sufficient to support, and that under normal circumstances do support, a prevalence of vegetation typically adapted for life in saturated soil conditions" (EPA, 2019).

The USACE will assert jurisdiction over the following waters:

- Traditional navigable waters;
- Wetlands adjacent to traditional navigable waters;
- Non-navigable tributaries of traditional navigable waters that are relatively permanent where the
 tributaries typically flow year-round or have continuous flow at least seasonally (e.g., typically three
 months); and
- Wetlands that directly abut such tributaries.

The USACE will decide jurisdiction over the following waters based on analysis to determine whether they have significant nexus with a traditional navigable water:

- Non-navigable tributaries that are not relatively permanent;
- Wetlands adjacent to non-navigable tributaries that are not relatively permanent; and
- Wetlands adjacent to, but that do not directly abut, a relatively permanent non-navigable tributary.

The USACE generally will not assert jurisdiction over the following features:



- Swales or erosional features (e.g., gullies, small washes characterized by low volume, infrequent, or short duration flow); and
- Ditches (including roadside ditches) excavated wholly in and draining only uplands, and that do not carry a relatively permanent flow of water.

The USACE will apply the significant nexus standard as follows:

- A significant nexus analysis will assess the flow characteristics and functions of the tributary itself
 and the functions performed by all wetlands adjacent to the tributary to determine if they
 significantly affect the chemical, physical, and biological integrity of downstream traditional
 navigable waters; and
- Significant nexus includes consideration of hydrologic and ecologic factors.

The USACE also regulates navigable waters under Section 10 of the Rivers and Harbor Act (33 U.S.C. 401 et seq.), which requires that a permit must be issued by the USACE to construct any structure in or over any navigable WOUS, as well as any proposed action (such as excavation/dredging or deposition of materials) that would alter or disturb these waters. If the proposed structure or activity affects the course, location, condition, or capacity of the navigable water, even if the proposed activity is outside the boundaries of the stream in associated wetlands, a Section 10 permit from the USACE is required.

2.2 Massachusetts Department of Environmental Protection

The Massachusetts Wetlands Protection Act (WPA) (Section 40 of Chapter 131 of the General Laws of Massachusetts and regulated under 310 Code of Massachusetts Regulations [CMR] section 10.00) defines multiple coastal (310 CMR 10.25-10.37) and inland resource areas (310 CMR 10.54-10.59) and gives the Massachusetts Department of Environmental Protection (MassDEP) jurisdiction over these resource areas. In most cases, the WPA also gives MassDEP jurisdiction over buffer zone extending 100 feet from the edge of the resource area. In addition to MassDEP, local municipalities' Conservation Commissions are responsible for administering the WPA and any local wetlands ordinance or bylaw.

The WPA defines two types of Land Subject to Flooding (310 CMR 10.57): isolated and bordering. Isolated Land Subject to Flooding (ILSF) is defined as "an isolated depression or a closed basin which serves as a ponding area for run-off or high ground water which has risen above the ground surface." Bordering Land Subject to Flooding (BLSF) is defined as "an area with low, flat topography adjacent to and inundated by flood waters rising from creeks, rivers, streams, ponds or lakes. It extends from the banks of these waterways and water bodies; where a bordering vegetated wetland occurs, it extends from said wetland." The boundary of BLSF is further defined as "the estimated maximum lateral extent of flood water which will theoretically result from the statistical 100-year frequency storm" as shown on the most recently available flood profile data prepared for the community by the National Flood Insurance Program (NFIP), currently administered by the Federal Emergency Management Agency (FEMA), successor to the U.S. Department of Housing and Urban Development). Under the WPA, ILSF and BLSF do not have associated buffer zones.

The WPA defines Bordering Vegetated Wetland (BVW) under 310 CMR 10.55 as any freshwater wetland which borders on creeks, rivers, stream ponds or lakes. Under the WPA, a 100-foot buffer zone is associated with BVWs. Isolated wetlands (IWs) are not connected to a waterway or waterbody and, therefore, are not regulated under the WPA and do not have an associated buffer zone under the WPA. IWs may have an associated buffer zone or similar zone associated with them under the local ordinance or bylaw. In some cases, IWs may qualify as ILSF and, in those instances, are regulated under the WPA.



The WPA defines Bank (310 CMR 10.54) as the portion of the land surface which normally abuts and confines a waterbody, occurring between a waterbody and a BVW and adjacent floodplain, or between a waterbody and an upland. Under the WPA, a 100-foot buffer zone is associated with Banks.

The WPA defines Riverfront Area (310 CMR 10.58) as the 200-foot area of land measured horizontally from a river's Mean Annual High Water (MAHW) line. The section defines a river as any stream that is perennial and includes, but is not limited to, streams shown as perennial on current USGS maps or that have a watershed size greater than or equal to one square mile. Riverfront Area is not associated with intermittent streams as they do not flow throughout the year. Under the WPA, Riverfront Area does not have an associated buffer zone.

A Notice of Intent filing is required from the MassDEP for any disturbance, including the removal of vegetation or alteration to a Banks, BVW, ILSF, BLSF, Riverfront Area, or buffer zone.

2.3 Town of Shutesbury Conservation Commission

The Shutesbury Conservation Commission (SCC) administers a local wetlands bylaw and regulations in addition to the WPA. The SCC has jurisdiction over any freshwater wetland, marsh, wet meadow, bog, swamp, isolated wetland, lake, pond, river, and stream (surface or subsurface) and land within 100 feet of any of these areas. The SCC also has jurisdiction over land under waterbodies and land subject to flooding or inundation by groundwater, surface water, storm flowage, or within a 100-year floodplain.

3.0 Project Site Characteristics

TRC reviewed publicly available literature and materials used for the investigation, survey, and report preparation, including:

- MassGIS OLIVER¹, the National Hydrography Dataset;
- The Shutesbury, Massachusetts 7.5 Minute Quadrangle (USGS, 2018);
- The FEMA Flood Insurance Rate Map (FIRM) Panels 2501280015A (effective date June 18, 1980);
- The U.S. Fish and Wildlife Service (USFWS), National Wetlands Inventory (NWI);
- The U.S. Department of Agriculture (USDA), NRCS Web Soil Survey;
- Recent aerial orthoimagery.

The following sections summarize TRC's review of each of these resources.

3.1 Hydrology

The Site is gently sloping in the eastern half with some steep slopes in the western portion. The Site generally drains northward and eastward off-site to wetlands and tributaries to Dean Brook to the north.

Pratt South Project Wetland and Waterbody Delineation Report

¹ The MassDEP Wetlands Conservancy Program uses aerial photography and photo interpretation to delineate and map wetland boundaries. These boundaries are available via the Massachusetts Office of Geographic Information (MassGIS) online mapping tool, OLIVER. Desktop review consisted of utilizing MassGIS OLIVER to gather a general understanding of existing conditions and potential regulated resource areas.



3.1.1 Floodplains

Flood hazard areas identified on the FEMA's FIRMs are identified as Special Flood Hazard Areas (SFHAs). SFHAs are defined as the area that will be inundated by the flood event having a 1-percent chance of being equaled or exceeded in any given year. The 1-percent annual chance flood is also referred to as the base flood or 100-year flood. FEMA uses a variety of labels for SFHAs:

Zone A	Zone A99	Zone AR/A
Zone AO	Zone AR	Zone V
Zone AH	Zone AR/AE	Zone VE, and
Zones A1-A30	Zone AR/AO	Zones V1-V30

Zone AE Zone AR/A1-A30

Moderate flood hazard areas, labeled Zone B or Zone X (shaded on FEMA mapping) are also shown on the FIRM, and are the areas between the limits of the base flood and the 0.2-percent-annual-chance (or 500-year) flood. The areas of minimal flood hazard, which are the areas outside the SFHA and higher than the elevation of the 0.2-percent-annual-chance flood, are labeled Zone C or Zone X (unshaded on FEMA mapping).

According to the FEMA FIRM 2501280015A (effective date June 18, 1980) the Site is located within a Zone C area of minimal flood disturbance zone. Base flood elevations and flood hazard factors are not available for this area.

3.2 Federal and State Mapped Wetlands and Streams

The USFWS is the principal federal agency tasked with providing information to the public on the status and trends of wetlands on a national scale. The USFWS NWI is a publicly available resource that provides detailed information on the abundance, characteristics, and distribution of nationwide wetlands (where mapped). NWI mapping data is offered to promote the understanding, conservation, and restoration of wetlands. The online MassGIS OLIVER mapping tool was accessed to determine the extent of statemapped aquatic resources.

According to TRC's review of NWI and MassGIS OLIVER mapping, there are four wetlands onsite: one in the northern central portion of the Site, two in the center of the Site, and one in the southeast corner of the Site. The northern central wetland extends off-site to the north and the wetland in the southeast corner extends off-site to the south.

3.3 Mapped Soils

The NRCS's Web Soil Survey identifies nine soil map units within the Site. Map units can represent a type of soil, a combination of soils, or miscellaneous land cover types (e.g., water, rock outcrop, developed impervious surface). Map units are usually named for the predominant soil series or land types within the map unit. A summary of soil characteristics for soils mapped at the Site are included in Table 1, below. The following sections provide details about hydric ratings, drainage class, prime farmland, and hydrologic soil groups (HSGs). Details about soil map unit descriptions are provided in the NRCS Soil Report included as Appendix D.



Table 1: Mapped Soils

Symbol	Soil Name	Hydric Rating (%)	Drainage Class	Hydrologic Soil Group	Farmland Classification
71B	Ridgebury fine sandy loam, 3 to 8 percent slopes, extremely stony	88	Poorly drained	D	Not prime farmland
73A	Whitman fine sandy loam, 0 to 3 percent slopes, extremely stony	99	Very poorly drained	D	Not prime farmland
109C	Chatfield-Hollis complex 8 to 15 percent slopes, rocky	2	Chatfield: Well drained Hollis: Somewhat excessively drained	Chatfield: B Hollis: D	Not prime farmland
109D	Chatfield-Hollis complex 15 to 25 percent slopes, rocky	0	Chatfield: Well drained Hollis: Somewhat excessively drained	Chatfield: B Hollis: D	Not prime farmland
245B	Hinckley loamy sand, 3 to 8 percent slopes	0	Excessively drained	А	Farmland of statewide importance
245C	Hinckley loamy sand, 8 to 15 percent slopes	0	Excessively drained	А	Farmland of statewide importance
441C	Gloucester sandy loam, 8 to 15 percent slopes, very stony	1	Somewhat excessively drained	С	Farmland of statewide importance
441D	Gloucester sandy loam, 15 to 25 percent slopes, very stony	0	Somewhat excessively drained	С	Not prime farmland
441F	Gloucester sandy loam, 25 to 45 percent slopes, very stony	0	Somewhat excessively drained	С	Not prime farmland

3.3.1 Hydric Rating

The Corps of Engineers Wetlands Delineation Manual (Environmental Laboratory, 1987) (1987 Manual) defines a hydric soil as "...a soil that in its undrained condition, is saturated, flooded or ponded long enough during the growing season to develop anaerobic conditions that favor the growth and regeneration of hydrophytic vegetation."

Due to limitations imposed by the small scale of the soil survey mapping, it is not uncommon to identify wetlands within areas not mapped as hydric soil while areas mapped as hydric often do not support wetlands. This concept is emphasized by the NRCS:

Enlargement of maps beyond the scale of mapping can cause misunderstanding of the detail of mapping and accuracy of soil line placement. The maps do not show the small areas of contrasting soils that could have been shown at a more detailed scale.

Hydric Soil Rating (HSR) indicates the percentage of a map unit that meets the criteria for hydric soils.

Map unit 71B has an HSR of 88 percent, map unit 73A has an HSR of 99 percent, map unit 109C has an HSR of 2 percent, map unit 441C has an HSR of 1 percent, and map units 109D, 245B, 245C, 441D, and



441F have an HSR of 0 percent. For map unit 71B, the hydric components within the map unit are Ridgebury, extremely stony and Whitman, extremely stony. For map unit 73A, the hydric components within the map unit are Whitman, extremely stony; Ridgebury, extremely stony; Scarboro; and Swansea. For map unit 109C, the hydric component within the map unit is Leicester, very stony. For map unit 441C, the hydric component within the map unit is Ridgebury, very stony.

3.3.2 Natural Drainage Class

Natural drainage class refers to the frequency and duration of wet periods under conditions similar to those under which the soil developed. Anthropogenic alteration of the water regime, either through drainage or irrigation, is not a consideration unless the alterations have significantly changed the morphology of the soil.

Map unit 71B is rated as poorly drained. Map unit 73A is rated as very poorly drained. For map units 109C and 109D, the Chatfield component is rated as well drained and the Hollis component is rated as somewhat excessively drained. Map units 245B and 245C are rated as excessively drained. Map units 441C, 441D, and 441F are rated as somewhat excessively drained.

3.3.3 Prime Farmland

Prime farmland is land that has the best combination of physical and chemical characteristics for producing food, feed, forage, fiber, and oilseed crops, and is available for these uses (the land could be cropland, pastureland, rangeland, forestland, or other land, but not urban built-up land or water). Land used for a specific high-value food or fiber crop is classified as "unique farmland." Generally, additional "farmlands of statewide importance" include those that are nearly prime farmland and that economically produce high yields of crops when treated and managed according to acceptable farming methods. In some local areas, there is concern for certain additional farmlands, even though these lands are not identified as having national or statewide importance. These farmlands are identified as being of "local importance" through ordinances adopted by local government. The NRCS State Conservationist reviews and certifies lists of farmland of state and local importance. These lists, along with state and locally established Land Evaluation and Site Assessment (LESA) systems where applicable, are used by federal agencies to review and evaluate activities that may impact farmland. As defined in 7 CFR Part 657, important farmland encompasses prime and unique farmland, as well as farmland of statewide and local importance.

According to the NRCS, map units 71B, 73A, 109C, 109D, 441D, and 441F are classified as "not prime farmland" and map units 245B, 245C, and 441C are classified as "farmland of statewide importance."

3.3.4 Hydrologic Soil Groups

Soils are assigned to a HSG based on estimates of runoff potential. Soils are assigned to one of four groups according to the rate of water infiltration when the soils are not protected by vegetation, are thoroughly wet, and receive precipitation from long-duration storms.

The soils in the United States are assigned to four groups (A, B, C, and D) and three dual classes (A/D, B/D, and C/D). The groups are defined as follows:

Group A: Soils have a high infiltration rate (low runoff potential) when thoroughly wet. These consist mainly of deep, well drained to excessively drained sands or gravelly sands. These soils have a high rate of water transmission.



Group B: Soils have a moderate infiltration rate when thoroughly wet. These consist chiefly of moderately deep or deep, moderately well drained or well drained soils that have moderately fine texture to moderately coarse texture. These soils have a moderate rate of water transmission.

Group C: Soils have a slow infiltration rate when thoroughly wet. These consist chiefly of soils having a layer that impedes the downward movement of water or soils of moderately fine texture or fine texture. These soils have a slow rate of water transmission.

Group D: Soils have a very slow infiltration rate (high runoff potential) when thoroughly wet. Soils consist chiefly of clays that have a high shrink-swell potential, soils that have a high water table, soils that have a claypan or clay layer at or near the surface, and soils that are shallow over nearly impervious material. These soils have a very slow rate of water transmission.

If a soil is assigned to a dual hydrologic group (A/D, B/D, or C/D), the first letter is for drained areas and the second is for undrained areas. Only the soils that in their natural condition in Group D are assigned to dual classes.

Map units 71B and 73A are in HSG D. For map units 109C and 109D, the Chatfield component is in HSG B and the Hollis component is in HSG D. Map units 245B and 245C are in HSG A. Map units 441C, 441D, and 441F are in HSG C.

4.0 Wetland and Stream Delineation Methodology

In addition to the desktop review described in Section 3.0, TRC biologists performed field investigations at the Site to identify wetlands, waterbodies, and other surface waters on July 29, 30, 2020 and August 3, 2020.

4.1 Non-wetland Aquatic Resource Methodology

Streams and other non-wetland aquatic features within the Site were identified by the presence of an OHWM, which is the line established by the fluctuations of water (33 CFR 328.3). The OHWM line is indicated by physical characteristics, which can include: a clear, natural line impressed on the bank; shelving; changes in the character of soil; destruction of terrestrial vegetation; the presence of litter and debris; or other characteristics of the surrounding areas. For streams three feet or more in width, each stream bank was delineated with blue flagging. For smaller streams, the stream centerline is delineated with notes for the width. Flags were located with a handheld global positioning system (GPS) unit and the data post-processed to achieve sub-meter accuracy.

4.2 Wetland Delineation Methodologies

The delineation of wetlands was conducted in accordance with criteria set forth in the 1987 Manual, the Regional Supplement to the Corps of Engineers Wetland Delineation Manual: Northcentral and Northeast Region (Version 2.0) (USACE, 2012) (Supplement), and the Delineating Bordering Vegetated Wetlands Under the Massachusetts Wetlands Protection Act- A Handbook (MassDEP, 1995) (the MassDEP Handbook).

The three-parameter approach to identify and delineate wetlands presented in the 1987 Manual and the Supplement requires that, except for atypical and disturbed situations, wetlands possess hydrophytic vegetation, hydric soils, and wetland hydrology. A two-parameter approach that considers only vegetation and hydrology indicators is presented in the MassDEP Handbook. Per the MassDEP Handbook, hydric soil is included as evidence of wetland hydrology.



Wetland boundary flags were located with a handheld GPS unit and the data were post-processed to achieve sub-meter accuracy. Delineated resources were classified in accordance with the system presented in *The Classification of Wetlands and Deepwater Habitats of the United States, Second Edition* (Federal Geographic Data Committee, 2013).

4.2.1 Hydrophytic Vegetation Methodologies

Hydrophytic vegetation is defined in the 1987 Manual as:

...the sum total of macrophytic plant life that occurs in areas where the frequency and duration of inundation or soil saturation produce permanently or periodically saturated soils of sufficient duration to exert a controlling influence on the plant species present.

Plants are categorized according to their occurrence in wetlands. Scientific names and wetland indicator statuses for vegetation are those listed in *The National Wetland Plant List: 2016 Wetland Ratings* (NWPL) (Lichvar et al., 2016). The indicator statuses specific to the "Northcentral and Northeast Region" as defined by the USACE apply to the Site. For upland species that are not listed on the NWPL, the Integrated Taxonomic Information System was referenced for currently accepted scientific names. The official short definitions for wetland indicator statuses are as follows:

- Obligate Wetland (OBL): Almost always occur in wetlands;
- Facultative Wetland (FACW): Usually occur in wetlands, but may occur in non-wetlands;
- Facultative (FAC): Occur in wetlands and non-wetlands (50/50 mix);
- Facultative Upland (FACU): Usually occur in non-wetlands, but may occur in wetlands; and
- Upland (UPL): Almost never occur in wetlands.

Plants that are not found in a region, but are found in an adjacent region, take on the indicator status of that adjacent region for dominance calculations. Plants that are included on the NWPL, but not within the Site region or an adjacent region, are not included in dominance calculations. Plants that are not found in wetlands in any region are considered "UPL" for dominance calculations.

Vegetation community sampling was accomplished using the methodologies outlined in the 2012 Supplement. The "50/20 rule" was applied to determine whether a species was dominant in its stratum. In using the 50/20 rule, the plants that comprise each stratum are ranked from highest to lowest in percent cover. The species that cumulatively equal or exceed 50 percent of the total percent cover for each stratum are dominant species, and any additional species that individually provides 20 percent or more percent cover is also considered dominant species of its respective strata.

A hydrophytic vegetation community is present when: 1) all of the dominant species are FACW and/or OBL (Rapid Test for Hydrophytic Vegetation); 2) greater than 50 percent of the dominant species' (as determined by the 50/20 rule) indicator statuses are FAC, FACW, or OBL (Dominance Test); and/or 3) when the calculated Prevalence Index is equal to or less than 3.0. When applying the Prevalence Index, all plants are assigned a numeric value based on indicator status (OBL = 1, FACW = 2, FAC = 3, FACU = 4, and UPL = 5) and their abundance (absolute percent cover) is used to calculate the prevalence index.

Cover types are also assigned to each wetland and waterbody in accordance with the system presented in *The Classification of Wetlands and Deepwater Habitats of the United States, Second Edition* (Federal Geographic Data Committee, 2013).



4.2.2 Hydric Soil Methodologies

Hydric soil indicators described in *Field Indicators for Identifying Hydric Soils in New England, Version 4* (New England Hydric Soils Technical Committee, 2017) and in *Field Indicators of Hydric Soils in the United States, Version 8.2* (NRCS, 2018) were used to determine the presence of characteristic soil morphologies resulting from prolonged saturation and/or inundation. Soil color was described using standard color notations provided on Munsell® soil color charts (X-Rite, Inc., 2015). Soil texture was determined using the methods described by Thien (1979). Soil test pits were dug using a spade shovel to a depth of approximately 20 inches or more (if needed).

Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin (MLRA Handbook) (USDA NRCS, 2006) was referenced to determine the hydric soil indicators that apply to the Site. Per the MLRA Handbook, the Site is within Major Land Resource Area (MLRA) 144A (New England and Eastern New York Upland, Southern Part) of Land Resource Region (LRR) R (Northeastern Forage and Forest Region). Hydric soil indicators that do not apply to this MLRA were not considered on the wetland determination data forms.

The presence or absence of hydric soils was determined through examination of samples extracted with a hand shovel or hand auger from the upper horizons of the soil profile. Soils were examined to depths of approximately 18 to 20 inches, unless restrictive layers such as hard pan, rock, densely packed fill materials, etc. were encountered at shallower depths.

4.2.3 Wetland Hydrology Methodologies

Per the 1987 Manual:

The term "wetland hydrology" encompasses all hydrologic characteristics of areas that are periodically inundated or have soils saturated to the surface at some time during the growing season. Areas with evident characteristics of wetland hydrology are those where the presence of water has an overriding influence on characteristics of vegetation and soils due to anaerobic and reducing conditions, respectively. Such characteristics are usually present in areas that are inundated or have soils that are saturated to the surface for sufficient duration to develop hydric soils and support vegetation typically adapted for life in periodically anaerobic soil conditions. Hydrology is often the least exact of the parameters, and indicators of wetland hydrology are sometimes difficult to find in the field. However, it is essential to establish that a wetland area is periodically inundated or has saturated soils during the growing season. (Environmental Laboratory, 1987)

Wetland hydrology indicators are grouped into 18 primary and 11 secondary indicators presented in the Supplement. The USACE considers wetland hydrology to be present when at least one primary indicator or two secondary indicators are identified.

5.0 Results

5.1 Upland Areas

The upland areas consist of successional forests throughout most the Site. The dominant vegetation in the uplands consists of sweet birch (*Betula lenta*), red maple (*Acer rubrum*), eastern white pine (*Pinus strobus*), eastern hemlock (*Tsuga canadensis*), American witch-hazel (*Hamamelis virginiana*), maple-leaf arrowwood (*Viburnum acerifolium*), false lily-of-the-valley (*Maianthemum canadense*), princess-pine (*Dendrolycopodium obscurum*), yellow birch (*Betula allegheniensis*), partridge berry (*Mitchella ripens*),



northern lady fern (*Athyrium angustum*), glossy false buckthorn (*Frangula alnus*), northern white oak (*Quercus alba*), mountain-laurel (*Kalmia latifolia*), late lowbush blueberry (*Vaccinium angustifolium*), northern red oak (*Quercus rubra*), one-flower Indian-pipe (*Monotropa uniflora*), cinnamon fern (*Osmundastrum cinnamomeum*), and hay-scented fern (*Dennstaedtia punctilobula*). The terrain of the Site is steeply sloping to the north and east in the western portion of the Site and gently sloping west in the eastern portion of the Site. The soils observed throughout upland portions of the Site were generally classified as loamy sand and sandy loam.

5.2 Delineated Wetlands and Waterbodies

TRC identified five wetlands and two waterbodies within the Site during the July and August 2020 resource delineation effort (Figure 2 in Appendix A). Delineated areas are described in the following sections and summarized at the end of this section in Table 2. Refer to the photographs in Appendix B and the wetland determination data forms in Appendix C for further details about each delineated area.

5.2.1 Delineated Wetlands

Wetland W-MJR-1 is a palustrine forested (PFO) wetland associated with stream S-MJR-1. This wetland is located along the eastern edge of the Site and extends off-site to the north, east, and south. The dominant vegetation included red maple, eastern hemlock, yellow birch, highbush blueberry (*Vaccinium corymbosum*), mountain-laurel, cinnamon fern, and spotted touch-me-not (*Impatiens capensis*). Indicators of wetland hydrology included saturation, geomorphic position, microtopographic relief, and the FAC-neutral test. Soils were composed of a thick layer of dark muck on top of silty clay loam. This soil meets Hydric Soil Indicator A1 as described in *Field Indicators of Hydric Soils in the United States, Version 8.2* (Field Indicators) (USDA NRCS, 2018). *This wetland is MassDEP jurisdictional and it also falls under USACE jurisdiction, as it is likely connected to other WOUS*.

Wetland W-MJR-2 is a PFO wetland located on the southern boundary in the eastern half of the Site and extends off-site to the south. The dominant vegetation included eastern hemlock, red maple, yellow birch, American witch-hazel, and cinnamon fern. Indicators of wetland hydrology included geomorphic position and microtopographic relief. Soils were composed of a layer of dark sandy loam over grayish-brown sandy clay loam with redoximorphic concentrations in the matrix. This soil meets Hydric Soil Indicator F3 as described in *Field Indicators of Hydric Soils in the United States, Version 8.2* (Field Indicators) (USDA NRCS, 2018). This wetland is likely MassDEP jurisdictional and it also falls under USACE jurisdiction, as it is likely connected to other WOUS.

Wetland W-MJR-3 is partially a palustrine shrub/scrub (PSS) and partially a PFO wetland with two small sections of palustrine emergent (PEM) wetland skirting the northwestern and southeastern edges of the wetland. The wetland is in the center of the Site and extends off-site to the south. The dominant vegetation within the PEM portion of the wetland included white meadowsweet (*Spiraea alba*) and swamp smartweed (*Persicaria hydropiperoides*). The dominant vegetation within the PSS portion of the wetland included speckled alder (*Alnus incana*), glossy false buckthorn, fringed sedge (*Carex crinita*), and spotted touch-menot. The dominant vegetation within the PFO portion of the wetland included eastern hemlock, red maple, glossy false buckthorn, and spotted touch-me-not. Indicators of wetland hydrology within the PEM portion of the wetland included surface water, saturation, inundation visible on aerial imagery, geomorphic position, and the FAC-neutral test. Indicators of wetland hydrology within the PSS portion of the wetland included saturation, saturation visible on aerial imagery, geomorphic position, and the FAC-neutral test. Indicators of wetland hydrology within the PFO portion of the wetland included water0stained leaves and geomorphic position. Soils within the PEM portion were unobtainable due to inundation. Soils within the PSS portion



were composed of a thick layer of dark mucky peat. Soils within the PFO portion were composed of a layer of dark loamy sand on top of a thick layer of brown sandy loam with redoximorphic concentrations in the matrix. The soil within the PEM portion of the wetland was not able to be sampled and therefore did not meet any Hydric Soil Indicator; however, according to the NRCS Web Soil Survey, the wetland's soil map unit has a high HSR (i.e., 88 percent). The soil within the PSS portion of the wetland meets Hydric soil indicator A1 according to the Field Indicators (USDA NRCS, 2018). The soil within the PFO portion of the wetland did not meet any Hydric Soil Indicator; however, according to the NRCS Web Soil Survey, the wetland's soil map unit has a high HSR (i.e., 88 percent). *This wetland is likely MassDEP jurisdictional and it also falls under USACE jurisdiction, as it is likely connected to other WOUS*.

Wetland W-MJR-4 is a PFO wetland located in the center of the Site and extends off-site to the north and south. The dominant vegetation included eastern hemlock, red maple, mountain-laurel, yellow birch, cinnamon fern, and sensitive fern (*Onoclea sensibilis*). Indicators of wetland hydrology included saturation, water-stained leaves, drainage patterns, geomorphic position, and microtopographic relief. Soils were composed of a layer of dark mucky peat over dark-greenish gray sandy loam with redoximorphic concentrations in the matrix. This soil meets hydric soil indicators A11 and F3 as described in the Field Indicators (USDA NRCS, 2018). *This wetland is MassDEP jurisdictional and it also falls under USACE jurisdiction, as it is likely connected to other WOUS*.

Wetland W-MJR-5 is a PFO wetland located in the northwest corner of the Site and extends off-site to the north. The dominant vegetation within this wetland included red maple, yellow birch, eastern hemlock, northern spicebush (*Lindera benzoin*), American witch-hazel, cinnamon fern, and false lily-of-the-valley. Indicators of wetland hydrology included high water table and saturation. Soils were composed of a thick layer of dark muck over gray sandy clay loam with redoximorphic concentrations in the matrix. This soil meets Hydric Soil Indicators A2 and A11 as described in the Field Indicators (USDA NRCS, 2018). *This wetland is likely MassDEP jurisdictional and it also falls under USACE jurisdiction, as it is likely connected to other WOUS*.

5.2.2 Delineated Waterbodies

Stream S-MJR-1 is an intermittent stream (R4, NWI classification) that flows out of wetland W-MJR-1 offsite to the south. The streambed was comprised of silt and clay. TRC observed an average width of approximately 15 feet and a water depth of approximately 10 inches. Stream S-MJR-1 has defined banks slightly wider than the OHWM at approximately 17 feet wide. The bank was delineated on both sides of the stream.

The USGS maps stream S-MJR-1 as intermittent. The USGS StreamStats analysis in Appendix E shows that it has a watershed of less than 0.5 square miles and has a predicted flow rate of less than 0.01 cubic feet per second at the 99% flow duration. Therefore, this stream is considered intermittent. *This stream is MassDEP jurisdictional and falls under USACE jurisdiction, as it is likely connected to other WOUS.*

Stream S-MJR-2 is an intermittent stream (R4, NWI classification) that flows out of wetland W-MJR-4 off-site to the north. The streambed was comprised of silt and clay. TRC observed an average width of approximately 6 feet and a water depth of approximately 0 inches. Stream S-MJR-1 has defined banks slightly wider than the OHWM at approximately 7 feet wide. The bank was delineated on both sides of the stream.

While the USGS maps stream S-MJR-2 as perennial, the USGS StreamStats analysis in Appendix E shows that it has a predicted flow rate greater than 0.01 cubic feet per second at the 99% flow duration but has a



watershed that is less than 0.5 square miles. Therefore, this stream is considered intermittent. *This stream* is *MassDEP jurisdictional and falls under USACE jurisdiction, as it is likely connected to other WOUS.*

Table 2. Delineated Wetlands and Waterbodies

Wetland Field Designation	Field Designated NWI Classification ¹	Assumed Jurisdictional Status	Assumed Buffer/ Setback Requirements
W-MJR-1	PFO	USACE/MassDEP/Local	100-ft buffer zone
W-MJR-2	PFO	USACE/MassDEP/Local	100-ft buffer zone
W-MJR-3	PEM/PSS/PFO	USACE/MassDEP/Local	100-ft buffer zone
W-MJR-4	PFO	USACE/MassDEP/Local	100-ft buffer zone
W-MJR-5	PFO	USACE/MassDEP/Local	100-ft buffer zone
S-MJR-1	R4	USACE/MassDEP/Local	100-ft buffer zone
S-MJR-2	R4	USACE/MassDEP/Local	100-ft buffer zone

¹ The Classification of Wetlands and Deepwater Habitats of the United States, Second Edition (Federal Geographic Data Committee, 2013). Categories include: Palustrine Forested (PFO), Palustrine Shrub-Scrub (PSS), Palustrine Emergent (PEM), and Riverine Intermittent (R4).

6.0 Conclusions

It is TRC's opinion that delineated wetlands W-MJR-1, W-MJR-2, W-MJR-3, W-MJR-4, and W-MJR-5 are BVWs regulated by MassDEP and are also likely under USACE jurisdiction. There are no buffers or setbacks associated with USACE-regulated wetlands. However, there is a 100-foot buffer zone associated with MassDEP- and SCC-regulated wetlands.

R4 streams S-MJR-1 and S-MJR-2 are USACE jurisdictional, as they are hydrologically connected to WOUS. These streams are also regulated by the MassDEP, as they flow within, into, or out of a MassDEP-regulated wetland resource areas.

Final determination of jurisdictional status for on-site wetlands and waterbodies must be made by the regulators.





7.0 References

- Environmental Laboratory. 1987. *Corps of Engineers Wetland Delineation Manual*. Technical Report Y-87-1. U.S. Army Corps of Engineers: Waterways Experiment Station; Vicksburg, MS.
- Environmental Protection Agency (EPA). 2019. *Electronic Code of Federal Regulations*. Title 40, Chapter 1, Subchapter H, Part 230, Subpart A, Section 230.3. https://www.ecfr.gov/cgi-bin/text-idx?SID=c2ac4e35564a7e132276a5092222dded&mc=true&node=se40.27.230 13&rgn=div8. Accessed August 2020.
- Federal Geographic Data Committee. 2013. Classification of wetlands and deepwater habitats of the United States. FGDC-STD-004-2013. Second Edition. Wetlands Subcommittee, Federal Geographic Data Committee and U.S. Fish and Wildlife Service, Washington, DC.
- Lichvar, R.W., D.L. Banks, W.N. Kirchner, and N.C. Melvin. 2016. *The National Wetland Plant List*: 2016 wetland ratings. Phytoneuron 2016-30: 1-17. Published 28 April 2016. ISSN 2153 733X.
- MassDEP. 1995. *Delineating Bordering Vegetated Wetlands Under the Massachusetts Wetland Protection Act.* Publication No. 17668-1022000-2/95-2.75-C.R. Massachusetts Department of Environmental Protection, Division of Wetlands and Waterways. Boston, MA. Scott Jackson, author.
- New England Hydric Soils Technical Committee. 2017. *Version 4, Field Indicators for Identifying Hydric Soils in New England*. New England Interstate Water Pollution Control Commission, Lowell, MA.
- U.S. Army Corps of Engineers (USACE). 2012. Regional Supplement to the Corps of Engineers Wetland Delineation Manual: Northcentral and Northeast Region (Version 2.0). U.S. Army Engineer Research and Development Center, Vicksburg, MS, 162 pp.
- USDA NRCS. Web Soil Survey. http://websoilsurvey.nrcs.usda.gov/. Accessed August 2020.
- USDA NRCS. 2018. Field Indicators of Hydric Soils in the United States, Version 8.2 L.M. Vasilas, G.W. Hurt, and J.F. Berkowitz (eds.). USDA, NRCS, in cooperation with the National Technical Committee for Hydric Soils.
- USDA NRCS. 2006. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. USDA Handbook 296.
- U.S. Department of the Interior, Geological Survey (USGS). 2018. Shutesbury, Massachusetts Quadrangle. 7.5 Minute Series (Topographic).



Appendix A: Figures







Appendix B: Photographs

PRATT SOUTH PROJECT PRATT CORNER ROAD, SHUTESBURY, MASSACHUSETTS

Photograph: 1

Date: 7/29/2020

Direction: West

Description:

Representative

conditions at Wetland W-

MJR-1.



Photograph: 2

Date: 7/30/2020

Direction: East

Description:

Representative

conditions at Wetland W-

MJR-4.





PRATT SOUTH PROJECT PRATT CORNER ROAD, SHUTESBURY, MASSACHUSETTS

Photograph: 3

Date: 8/3/2020

Direction: South

Description:

Representative conditions in uplands near Wetland W-MJR-5.



Photograph: 4

Date: 8/3/2020

Direction: North

Description:

Representative

conditions at Wetland W-

MJR-5.





PRATT SOUTH PROJECT PRATT CORNER ROAD, SHUTESBURY, MASSACHUSETTS

Photograph: 5

Date: 7/29/2020

Direction: South

Description:

Representative conditions at Wetland W-

MJR-2.



Photograph: 6

Date: 7/30/2020

Direction: South

Description:

Representative conditions in PEM portion of Wetland W-

MJR-3.





PRATT SOUTH PROJECT PRATT CORNER ROAD, SHUTESBURY, MASSACHUSETTS

Photograph: 7

Date: 7/30/2020

Direction: South

Description:

Representative

conditions in PFO portion of Wetland W-MJR-3.



Photograph: 8

Date: 7/30/2020

Direction: North

Description:

Representative conditions in uplands in Right-of-Way near Wetland W-MJR-3.







Appendix C: Wetland Determination Data Forms

Project/Site: Pratt South	City/County: Shu	tesbury, Franklin	Sampling Date:	2020-July-29		
Applicant/Owner: W.D. Cowls, 1	Inc.	State: MA	Sampling Point:	W-PMO-01_PFO-1		
Investigator(s): Matt Regan, Mo	lly Lennon, Caroline Harrington	Section, Township,	Range:			
Landform(hillslope,terrace,etc.):	Swamp	Local relief (concave, conv	ex, none): Concave	Slope (%): 0 to 1		
Subregion(LRRorMLRA): MLF	RA 144A of LRR R	Lat: 42.410832331	6 Long: -72.4596094061	Datum: WGS84		
SoilMapUnitName: Hinckley lo	amy sand, 8 to 15 percent slopes		NWI classific	cation:		
Areclimatic/hydrologicconditions	onthesitetypicalforthistimeofyear?	Yes 🔽 No	(If no, explain in Rema	rks.)		
Are Vegetation, Soil,	or Hydrology significantly dis		al Circumstances" present?	Yes No		
Are Vegetation, Soil,	or Hydrology naturally probl	lematic? (If needed,	explain any answers in Rem	arks.)		
Summary of Findings – At	ttach site map showing sampli	ng point locations, trar	nsects, important featur	es, etc.		
Hydrophytic Vegetation Present?	Yes _ ✓ _ No					
Hydric Soil Present?	Yes No	Is the Sampled Area withi	in a Wetland?	Yes No		
		·		W-PMO-01		
Wetland Hydrology Present?	Yes No	If yes, optional Wetland S	ite id:	W-PMU-01		
•	cedures here or in a separate report	:)				
Covertype is PFO.						
LIVDDOLOCV						
HYDROLOGY						
Wetland Hydrology Indicators:						
Primary Indicators (minimum of c	one is required; check all that apply)		Secondary Indicators (minir	num of two required)		
Surface Water (A1)	Water-Stained Lea	aves (B9)	Surface Soil Cracks (B6)			
High Water Table (A2)	Aquatic Fauna (B1		Drainage Patterns (B10)			
✓ Saturation (A3)	Marl Deposits (B1	5)	Moss Trim Lines (B16)			
Water Marks (B1)	Hydrogen Sulfide	Odor (C1)	Dry-Season Water Table (C2)			
Sediment Deposits (B2)	Oxidized Rhizospł	heres on Living Roots (C3)	Crayfish Burrows (C8)			
			Saturation Visible on Ae	3 3 · ·		
Drift Deposits (B3)	Presence of Redu		Stunted or Stressed Plants (D1)			
Algal Mat or Crust (B4)		ction in Tilled Soils (C6)	✓ Geomorphic Position (D	2)		
Iron Deposits (B5)	Thin Muck Surface		Shallow Aquitard (D3)			
Inundation Visible on Aerial In	· · · · · · · · · · · · · · · · · · ·	Remarks)	<u>✓</u> Microtopographic Relief	(D4)		
Sparsely Vegetated Concave S	urface (B8)		<u>✓</u> FAC-Neutral Test (D5)			
Field Observations:						
Surface Water Present?	Yes No <u></u> Depth	(inches):	_			
Water Table Present?	Yes No Depth	(inches):	Wetland Hydrology Present	? Yes No		
Saturation Present?	Yes 🗸 No Depth	(inches): 0				
(includes capillary fringe)			-			
	gauge, monitoring well, aerial photo	s provious inspections) if	available:			
Describe Recorded Data (stream)	gauge, monitoring well, aeriai prioto:	s, previous irispections), ir	avallable.			
Remarks:						
Remarks:						
Remarks:						
Remarks:						
Remarks:						
Remarks:						
Remarks:						

<u>ree Stratum</u> (Plot size: <u>30 ft</u>)		Dominant Species?	Indicator Status	Dominance Test worksheet: Number of Dominant Species That	5	(4)
. Acer rubrum	10	Yes	FAC	Are OBL, FACW, or FAC:	3	(A)
2. Tsuga canadensis	10	Yes	FACU	Total Number of Dominant Species	7	(B)
Betula alleghaniensis	10	Yes	FAC	Across All Strata:	•	
Fraxinus pennsylvanica	5	No	FACW	Percent of Dominant Species That Are OBL, FACW, or FAC:	71.4	(A/B)
i. i.				Prevalence Index worksheet:		
				Total % Cover of:	Multiply B	<u>y:</u>
		= Total Cov		OBL species 20	x 1 =	20
Continue (Charles Charles (District) 45 ft	35	_= 10tal Cov	er	FACW species 90	x 2 =	180
iapling/Shrub Stratum (Plot size:15 ft)	25	\/	EA CIA/	FAC species 20	x 3 =	60
. Vaccinium corymbosum	35	Yes	FACW	FACU species 35	x 4 =	140
Kalmia latifolia	15	Yes	FACU	UPL species 0	x 5 =	0
				Column Totals 165	(A)	400 (B)
·				Prevalence Index = B/A =	2.4	
•				Hydrophytic Vegetation Indicators:		
				1- Rapid Test for Hydrophytic V	egetation	
·				✓ 2 - Dominance Test is >50%	egetation	
	50	= Total Cov	er	\checkmark 3 - Prevalence Index is \le 3.01		
<u>lerb Stratum</u> (Plot size: <u>5 ft</u>)				4 - Morphological Adaptations	(Provide s	unnorting
. Osmundastrum cinnamomeum	25	Yes	FACW	data in Remarks or on a separate sh		upporting
. Impatiens capensis	20	Yes	FACW	Problematic Hydrophytic Veget	-	lain)
Symplocarpus foetidus	15	No	OBL	¹Indicators of hydric soil and wetland		
. Kalmia latifolia	10	No	FACU	present, unless disturbed or probler	, 0,	y mast be
o. Onoclea sensibilis	5	No	FACW	Definitions of Vegetation Strata:		
5. Carex crinita		No	OBL	Tree – Woody plants 3 in. (7.6 cm) or	more in di	ameter a
7.				breast height (DBH), regardless of he		arrieter a
				Sapling/shrub – Woody plants less th	_	3H and
				greater than or equal to 3.28 ft (1 m)		
				Herb – All herbaceous (non-woody)	olants, rega	ardless of
				size, and woody plants less than 3.2	_	
2				Woody vines – All woody vines great	er than 3.2	8 ft in
Z		= Total Cov	or	height.		
Manda Mina Charles (District 20 ft)	80	_ 10tal Cov	er	Hydrophytic Vegetation Present? Y	es 🗸 No)
Noody Vine Stratum (Plot size: 30 ft)				y if ye ignit		
·						
). 						
3						
l .						
	0	= Total Cov	er			

	•	to the de	•			indicato	r or confirm the	absence of indicate	ors.)
Depth	Matrix		Redox				_		
(inches)	Color (moist)		Color (moist)	<u>%</u>	Type ¹	Loc ²	-	xture	Remarks
0 - 18	2.5Y 2.5/1	100		_			•	/luck	
18 - 24	10Y 5/1	100		_			Silty C	lay Loam	
				_					
				_					-
				_					-
				_					
				_					
				_					
								_	
¹Type: C = 0	Concentration, D =	Depletio	n, RM = Reduced	Mati	rix, MS =	Masked	Sand Grains.	² Location: PL = Pore	e Lining, M = Matrix.
Hydric Soil	Indicators:							Indicators for P	roblematic Hydric Soils³:
_✓ Histoso			Polyvalue Bel	ow S	urface (S	8) (LRR 1	R, MLRA 149B)	2 cm Muck i	(A10) (LRR K, L, MLRA 149B)
Histic Ep	oipedon (A2)		Thin Dark Sur	face	(S9) (LRR	R, MLR	A 149B)		e Redox (A16) (LRR K, L, R)
Black Hi	istic (A3)		Loamy Mucky	/ Min	eral (F1)	(LRR K, I	_)		/ Peat or Peat (S3) (LRR K, L, R)
	en Sulfide (A4)		Loamy Gleye						e (S7) (LRR K, L)
	d Layers (A5)		Depleted Mat					Polyvalue B	elow Surface (S8) (LRR K, L)
	d Below Dark Surf								urface (S9) (LRR K, L)
	ark Surface (A12) Jucky Mineral (S1)		Depleted Dar Redox Depre			1		Iron-Manga	nese Masses (F12) (LRR K, L, R)
-	Gleyed Matrix (S4)		Redox Depie	33101	15 (1-0)				loodplain Soils (F19) (MLRA 149B)
	Redox (S5)							Mesic Spod	ic (TA6) (MLRA 144A, 145, 149B)
-	d Matrix (S6)							Red Parent	
	u Matrix (36) Irface (S7) (LRR R, N	ΛΙ D Δ 1/0	OR)						w Dark Surface (TF12)
Dark 30	11 Idee (37) (ERR 16, 1	VILION 1-7.	,,,					Other (Expl	ain in Remarks)
3Indicators	of hydrophytic veg	getation a	and wetland hydr	olog	y must be	e preser	t, unless distur	bed or problematic	
Restrictive	Layer (if observed)	:							
	Type:		None			Hydric	Soil Present?		Yes/_ No
	Depth (inches):								
Remarks:									

Project/Site: Pratt South	City/County։ Տիւ	ıtesbury, Hampshire	Sampling Da	ite: 2020-July-29		
Applicant/Owner: W.D. Cowls,	Inc.	State: MA	Sampling Poin	t: W-PMO-01_UPL-1		
Investigator(s): Matt Regan, Mo	olly Lennon, Caroline Harrington	Section, Township,	Range:			
Landform(hillslope,terrace,etc.):	Hillslope	Local relief (concave, conv	ex, none): Concave	Slope (%): 1 to 3		
Subregion(LRRorMLRA): ML	_RA 144A of LRR R	Lat: 42.410772903	9 Long: -72.45973739	78 Datum: WGS84		
SoilMapUnitName: Hinckley lo	oamy sand, 8 to 15 percent slopes		NWI class	sification:		
Areclimatic/hydrologicconditions	sonthesitetypicalforthistimeofyear?	Yes 🟒 No	(If no, explain in Re	marks.)		
Are Vegetation, Soil,	or Hydrology significantly d	isturbed? Are "Norma	al Circumstances" presen	t? Yes 🟒 No		
Are Vegetation, Soil,	or Hydrology naturally prob	olematic? (If needed,	explain any answers in R	emarks.)		
SUMMARY OF FINDINGS – A	ttach site map showing sampl	ing point locations, trar	nsects, important feat	ures, etc.		
Hydrophytic Vegetation Present?	? Yes No					
Hydric Soil Present?	Yes No	Is the Sampled Area withi	n a Wetland?	Yes No/_		
	Yes No _ _ _	·				
Wetland Hydrology Present?	·	If yes, optional Wetland Si	te iD:			
· ·	ocedures here or in a separate repor	t)				
Covertype is UPL.						
111/2201051						
HYDROLOGY						
Wetland Hydrology Indicators:						
Primary Indicators (minimum of	one is required; check all that apply)		Secondary Indicators (m	inimum of two required)		
Surface Water (A1)	Water-Stained Le	eaves (B9)	Surface Soil Cracks (E	36)		
High Water Table (A2)	Aquatic Fauna (B		Drainage Patterns (B	10)		
Saturation (A3)	Marl Deposits (B	15)	Moss Trim Lines (B16)			
Water Marks (B1)	Hydrogen Sulfide	e Odor (C1)	Dry-Season Water Table (C2)			
Sediment Deposits (B2)	Oxidized Rhizosp	heres on Living Roots (C3)				
			Saturation Visible on			
Drift Deposits (B3)	Presence of Redu		Stunted or Stressed Plants (D1)			
Algal Mat or Crust (B4)		iction in Tilled Soils (C6)	Geomorphic Position (D2)			
Iron Deposits (B5) Inundation Visible on Aerial II	Thin Muck Surface		Shallow Aquitard (D3)			
	· · · · · · · · · · · · · · · · · · ·	Remarks)	Microtopographic Re			
Sparsely Vegetated Concave : Field Observations:	Surface (Do)		FAC-Neutral Test (D5)	1		
	Voc. No. (Donth	(inchas)				
Surface Water Present?	·	(inches):				
Water Table Present?		(inches):	Wetland Hydrology Pres	ent? Yes No		
Saturation Present?	Yes No _ _/ Depth	(inches):				
(includes capillary fringe)						
Describe Recorded Data (stream	gauge, monitoring well, aerial photo	os, previous inspections), if a	available:			
Remarks:						
nemarks.						

		Indicator Status	Dominance Test worksheet: Number of Dominant Specie		1	(A)
20	Yes	FACU	Are OBL, FACW, or FAC:	_	'	(A)
10	Yes	FAC	Total Number of Dominant	Species	9	(B)
10	Yes	FACU	Across All Strata:	_		
10	Yes	FACU	·	s That	11.1	(A/B)
						`_
	= Total Cov	er	' <u> </u>			0
	- 10101 COV	Ci		0 :	x 2 =	0
25	Voc	EACH	FAC species	10 :	x 3 =	30
			FACU species 1	15	x 4 =	460
			- UPL species	12 :	x 5 =	60
			- Column Totals 1	37	(A)	550 (B)
5	No	NI	Prevalence Index	= B/A =	4	
			Hydronhytic Vegetation Indi	icators:		
			, , ,		getation	
			· · ·		Secucion	
55	= Total Cov	er				
					Drovido ci	ınnortin
12	Yes	UPL	, ,			pportin
10	Yes	FACU	· ·			lain)
10	Yes	FACU		_		
			,		, 0,	must b
			-		auc	
			_			
						ameter a
			-		_)
			-			on and
			- I -			rdloss o
						ii uless o
						8 ft in
			-	ics gi cate	1 (11011 3.2	010111
37	_= Total Cov	er	·		- N-	
			Hydrophytic vegetation Pre	esent? Ye	S NC	
			_			
			_			
	= Total Cov	or	-			
	% Cover 20 10 10 10 10 50 25 15 10 5 110 5 10 5	% Cover Species?	20 Yes	Scover Species? Status	Number of Dominant Species That Are OBL, FACW, or FAC:	Number of Dominant Species That Are OBL, FACW, or FAC: 10 Yes FACU 10 Yes FACU 10 Yes FACU 20 Yes FACU 10 Yes FACU 20 Yes FACU 25 Yes FACU 25 No NI 25 No NI 20 Yes FACU 20 Yes Y

Profile Deso	cription: (Describe Matrix	to the d	epth needed to d Redox			indicato	or confirm the a	bsence of indi	icators.)
(inches)	Color (moist)	%	Color (moist)	%	Type ¹	Loc ²	Textu	re	Remarks
0 - 3	7.5YR 3/2	100	Color (IIIoist)		Турс		Loamy		Kemarks
3 - 20	10YR 5/3	100					Loamy S		
3 20	1011(3/3	100					Louiny	Juliu	
	•			- —					
				-					-
				-					-
				- —					
				- —					
				-					
				-					
				-					
	•			- —				-	_
1T C		D +: -	- DM Deduced			N 4 = = - = =	Court Curius 21	tiDI I	David links - M. Makski
	Concentration, D =	Depletio	on, KIVI = Reduced	iviati	rix, IVIS =	iviasked	Sand Grains. ² L		Pore Lining, M = Matrix.
Hydric Soil			Daharaha B	lavec	f /^	:0\ /I PP	D MI DA 4 40D)		or Problematic Hydric Soils³:
Histoso	i (A1) pipedon (A2)		Polyvalue Be				R, MLRA 149B) A 149B)		ıck (A10) (LRR K, L, MLRA 149B)
	istic (A3)		Loamy Muck						rairie Redox (A16) (LRR K, L, R)
	en Sulfide (A4)		Loamy Gleye			(Litting)	-,		ucky Peat or Peat (S3) (LRR K, L, R)
	d Layers (A5)		Depleted Ma						rface (S7) (LRR K, L)
Deplete	d Below Dark Surf	ace (A11						•	e Below Surface (S8) (LRR K, L) rk Surface (S9) (LRR K, L)
Thick Da	ark Surface (A12)		Depleted Dar	k Su	rface (F7))			nganese Masses (F12) (LRR K, L, R)
Sandy N	Mucky Mineral (S1)		Redox Depre	ssior	ıs (F8)				nt Floodplain Soils (F19) (MLRA 149B)
Sandy C	Gleyed Matrix (S4)								podic (TA6) (MLRA 144A, 145, 149B)
Sandy F	Redox (S5)								ent Material (F21)
Stripped	d Matrix (S6)								allow Dark Surface (TF12)
Dark Su	ırface (S7) (LRR R, N	MLRA 14	9B)					-	xplain in Remarks)
3Indicators	of hydrophytic veg	atation	and wetland hydr	വര	y must h	a nracar	nt unlace dieturhe	ad or problem:	· atic
	Layer (if observed)		and wettand nyan	0106.	y masc b	Preser	it, uriless distarbe	ed of problem	utc.
	Type:	•	None			Hydric	Soil Present?		Yes No⁄_
	Depth (inches):	-	None	-		liyanc	John Frederic.		103 <u> </u>
Remarks:	Берит (писпез).					l .			
Remarks.									
ĺ									

State: MA Sampling Point: W-PMO-02_PFO-1 State: MA Sampling Point: W-PMO-02_PFO-1 State: Accorded to the following point: W-PMO-02_PFO-1 State: MA Sampling Point: W-PMO-02_PFO-1 State: Accorded to the following point: M-PMO-02_PFO-1 State: Accorded
lief (concave, convex, none): Concave at: 42.4102973147 Long: -72.4614644051 Datum: WGS84 NWI classification: Yes _ No (If no, explain in Remarks.) Are "Normal Circumstances" present? Yes _ No (If needed, explain any answers in Remarks.) It locations, transects, important features, etc.
At: 42.4102973147 Long: -72.4614644051 Datum: WGS84 NWI classification: Yes _ No _ (If no, explain in Remarks.) Are "Normal Circumstances" present? Yes _ No _ (If needed, explain any answers in Remarks.) It locations, transects, important features, etc.
NWI classification: Yes No (If no, explain in Remarks.) Are "Normal Circumstances" present? Yes No (If needed, explain any answers in Remarks.) t locations, transects, important features, etc.
Yes No (If no, explain in Remarks.) Are "Normal Circumstances" present? Yes No (If needed, explain any answers in Remarks.) t locations, transects, important features, etc. ampled Area within a Wetland? Yes No
Are "Normal Circumstances" present? Yes No (If needed, explain any answers in Remarks.) t locations, transects, important features, etc. ampled Area within a Wetland? Yes No
(If needed, explain any answers in Remarks.) t locations, transects, important features, etc. ampled Area within a Wetland? Yes No
t locations, transects, important features, etc. ampled Area within a Wetland? Yes No
ampled Area within a Wetland? Yes No
ampled Area within a Wetland? Yes No
production rectains size is.
Secondary Indicators (minimum of two required)
Surface Soil Cracks (B6)
Drainage Patterns (B10)
Moss Trim Lines (B16)
Dry-Season Water Table (C2)
Living Roots (C3) — Crayfish Burrows (C8)
Saturation Visible on Aerial Imagery (C9)
(C4) Stunted or Stressed Plants (D1)
illed Soils (C6) Geomorphic Position (D2) Shallow Aguitard (D3)
✓ Microtopographic Relief (D4)
FAC-Neutral Test (D5)
Wetland Hydrology Present? Yes ∠∠ No

	Absolute	Dominant	Indicator	Dominance Test worksheet:		
<u>Tree Stratum</u> (Plot size: <u>30 ft</u>)	% Cover	Species?	Status	Number of Dominant Species That	3	(A)
1. Tsuga canadensis	30	Yes	FACU	Are OBL, FACW, or FAC:		(A)
2. Acer rubrum	10	Yes	FAC	Total Number of Dominant Species	5	(B)
3. <i>Quercus rubra</i>	5	No	FACU	Across All Strata:		
4.				Percent of Dominant Species That	60	(A/B)
5.				Are OBL, FACW, or FAC:		
6.				Prevalence Index worksheet:		_
7.				Total % Cover of:	Multiply	-
	45	= Total Cov	er	OBL species 0	x 1 = _	0
Sapling/Shrub Stratum (Plot size:15 ft)		-		FACW species 60	x 2 =	120
1. Betula alleghaniensis	30	Yes	FAC	FAC species 40	x 3 =	120
Hamamelis virginiana	20	Yes	FACU	FACU species 60	x 4 =	240
3. Osmundastrum cinnamomeum	0	No	FACW	UPL species 0	x 5 =	0
4.		110	17.000	Column Totals 160	(A)	480 (B)
5.				Prevalence Index = B/A =	3	
6.				Hydrophytic Vegetation Indicators:		
				1- Rapid Test for Hydrophytic \	/egetation	
7		T-t-LC-		2 - Dominance Test is >50%		
	50	= Total Cov	er	3 - Prevalence Index is ≤ 3.01		
Herb Stratum (Plot size:5 ft)	60	V	EA CIA/	4 - Morphological Adaptations	¹ (Provide	supporting
1. Osmundastrum cinnamomeum	60	Yes	FACW	data in Remarks or on a separate sh	neet)	
2. Acer pensylvanicum	5	No	FACU	Problematic Hydrophytic Vege	tation¹ (Ex	(plain)
3				landicators of hydric soil and wetlan	d hydrolo	gy must be
4				present, unless disturbed or proble	matic	
5				Definitions of Vegetation Strata:		
6.				Tree – Woody plants 3 in. (7.6 cm) or	r more in o	diameter at
7				breast height (DBH), regardless of h	eight.	
8.				Sapling/shrub – Woody plants less t		DBH and
9				greater than or equal to 3.28 ft (1 m		
10				Herb – All herbaceous (non-woody)		gardless of
11				size, and woody plants less than 3.2		
12.				Woody vines – All woody vines grea	ter than 3.	.28 ft in
	65	= Total Cov	er	height.		
Woody Vine Stratum (Plot size:30 ft)		_		Hydrophytic Vegetation Present?	Yes 🟒 N	lo
1.						
2.						
3.						
4.				•		
	0	= Total Cov	er	•		
		-				
Remarks: (Include photo numbers here or on a separat	e sheet.)					

	cription: (Describe	to the de	-			ndicator	or confirm the al	osence of indicator	s.)
Depth _	Matrix	04	Redox			1002	Tout	turo	Domorks
(inches)	Color (moist)	<u>%</u>	Color (moist)	<u>%</u>	Type ¹	Loc²	Text		Remarks
0 - 5	10YR 2/2	100					Sandy		
5 - 20	10YR 5/2	95	10YR 5/6	5	C	<u>M</u>	Sandy Cl	ay Loam	
				- —					
				_					
				_					
				_					
1Type: C = C	Concentration, D =	Denletio	n RM = Reduced	Mat	rix MS =	Masked	Sand Grains 21 (ocation: PL = Pore I	ining M = Matrix
Hydric Soil		Depictio	n, Kivi Kedacea	IVIGE	17, 1415	Maskea	Suria Grains.		oblematic Hydric Soils³:
-			Dobarduo Do	o C	urfaca (C	0) /I DD I	MIDA 140D)		•
Histoso	oipedon (A2)		Polyvalue Be Thin Dark Su				R, MLRA 149B)		10) (LRR K, L, MLRA 149B)
	istic (A3)		Loamy Muck						Redox (A16) (LRR K, L, R)
	en Sulfide (A4)		Loamy Gleye			(LIXIX IX, L	,	-	Peat or Peat (S3) (LRR K, L, R)
	d Layers (A5)		Depleted Ma					Dark Surface	
	d Below Dark Surfa	ace (A11						•	ow Surface (S8) (LRR K, L)
	ark Surface (A12)	•	Depleted Dar			1			face (S9) (LRR K, L)
Sandy N	lucky Mineral (S1)		Redox Depre	ssior	ıs (F8)				ese Masses (F12) (LRR K, L, R)
	Gleyed Matrix (S4)								odplain Soils (F19) (MLRA 149B)
-	Redox (S5)								(TA6) (MLRA 144A, 145, 149B)
-	d Matrix (S6)							Red Parent M	
	rface (S7) (LRR R, N	/LRA 149	9B)						Dark Surface (TF12)
	,		•					Other (Explain	n in Remarks)
-	of hydrophytic veg		and wetland hydr	olog	y must be	e presen	t, unless disturbe	d or problematic.	
	Layer (if observed): _								
	Type:		None	-		Hydric	Soil Present?		Yes/_ No
	Depth (inches):								-
Remarks:									
Ì									

Project/Site: Pratt South	City/County: Shu	utesbury, Franklin	Sampling Date: 2020-July-29			
Applicant/Owner: W.D. Cowls,	Inc.	State: MA	Sampling Point: W-PMO-02_UPL-1			
Investigator(s): Matt Regan, Mo	olly Lennon, Caroline Harrington	Section, Township,	Range:			
Landform(hillslope,terrace,etc.):	Flat	Local relief (concave, conv	ex, none): None	Slope (%): 0 to 1		
Subregion(LRRorMLRA): ML	RA 144A of LRR R	Lat: 42.410339224	2 Long: -72.4618152716	Datum: WGS84		
SoilMapUnitName: Hinckley lo	pamy sand, 8 to 15 percent slopes		NWI classifica	ation:		
Areclimatic/hydrologicconditions	onthesitetypicalforthistimeofyear?	Yes 🟒 No	(If no, explain in Remar	ks.)		
Are Vegetation, Soil,	or Hydrology significantly d		al Circumstances" present?	Yes No		
Are Vegetation, Soil,	or Hydrology naturally prob	olematic? (If needed,	explain any answers in Rema	rks.)		
Summary of Findings – A	ttach site map showing sampl	ing point locations, trar	nsects, important feature	es, etc.		
Hydrophytic Vegetation Present?	Yes No _ ✓					
Hydric Soil Present?	Yes No _ _ _	Is the Sampled Area withi	n a Wetland?	Yes No		
Wetland Hydrology Present?	Yes No _ _ ∠	If yes, optional Wetland S	ite ID:			
			ice is.			
Covertype is UPL.	ocedures here or in a separate repor	ij				
Covertype is OPL.						
HYDROLOGY						
HIDROLOGI						
Wetland Hydrology Indicators:						
Primary Indicators (minimum of	one is required; check all that apply)	1	Secondary Indicators (minim	um of two required)		
Surface Water (A1)	Water-Stained Le	eaves (B9)	Surface Soil Cracks (B6)			
High Water Table (A2)	Aquatic Fauna (B	13)	Drainage Patterns (B10)			
Saturation (A3)	Marl Deposits (B	15)	Moss Trim Lines (B16)			
Water Marks (B1)	Hydrogen Sulfide		Dry-Season Water Table (C2)			
Sediment Deposits (B2)	Oxidized Rhizosp	heres on Living Roots (C3)	Crayfish Burrows (C8)Saturation Visible on Aer	ial Imagony (CQ)		
Duift Danasita (D2)	Dunnan of Dod	d June 10 (C.4)				
Drift Deposits (B3) Algal Mat or Crust (B4)	Presence of Redu	icea iron (C4) iction in Tilled Soils (C6)	Stunted or Stressed Plants (D1)			
Iron Deposits (B5)	Thin Muck Surface		,			
Inundation Visible on Aerial II						
Sparsely Vegetated Concave S	· · ·	nemano,	FAC-Neutral Test (D5)	.5 1)		
Field Observations:						
Surface Water Present?	Yes No Depth	ı (inches):				
Water Table Present?		(inches):	- Wetland Hydrology Present?	Yes No		
		· · · · · · · · · · · · · · · · · · ·	- Wedand Hydrology Fresents	res 110 _ _/ _		
Saturation Present?	Yes No _ _∕ Depth	(inches):				
(includes capillary fringe)						
Describe Recorded Data (stream	gauge, monitoring well, aerial photo	os, previous inspections), if a	available:			
Remarks:						

	Dominant Species?	Indicator Status	Dominance Test worksheet: Number of Dominant Species That	: 1	(4)
30	Yes	FACU	Are OBL, FACW, or FAC:		(A)
20	Yes	FACU	Total Number of Dominant Species	5 7	(B)
10	No	FAC	Across All Strata:		
				28.6	(A/B)
					-
60	= Total Cov	er	· -	-	0
	-		· -		0
25	Yes	FAC			165
			· -	x 4 =	320
			UPL species 5	x 5 =	25
	110	TAC	Column Totals 140	(A)	510 (B)
			Prevalence Index = B/A =	3.6	
			Hydrophytic Vegetation Indicators:		
			1 , , , ,		
				J	
55	_= Total Cov	er	3 - Prevalence Index is $\leq 3.0^{\circ}$		
			4 - Morphological Adaptation	s¹ (Provide s	upporting
10	Yes	FACU			
10	Yes	FAC	•		olain)
5	Yes	UPL			
			,	, 0.	,
			· · · · · · · · · · · · · · · · · · ·		
			_	or more in d	iameter a
					arricter a
				_	RH and
			• II • • • • • • • • • • • • • • • • •		311 ana
			. ['		ardless of
					8 ft in
25	_= Total Cov	er		Voc No	
			Trydrophytic vegetation Fresent:	162 140	<i></i>
	30 20 10 60 25 20 10 55 10 10 5	20 Yes 10 No 60 = Total Cov 25 Yes 20 Yes 10 No 55 = Total Cov 10 Yes 10 Yes 5 Yes	30	30 Yes	30

Profile Des	cription: (Describe Matrix	to the d	epth needed to d Redox			indicato	r or confirm the a	bsence of indi	icators.)
(inches)	Color (moist)	<u></u> %	Color (moist)	%	Type ¹	Loc ²	Textu	re	Remarks
0 - 2	7.5YR 2.5/2	100	Color (moist)		Турс		Loamy		Kemarks
2 - 20	7.5YR 4/4	100		-			Loamy S		-
	7.511(4)4	100					Loaniy	Juliu	
	•		•	-			•		
				_					
				_					
				-					
				- —					
				_					-
				-					
	•			- —				-	
1T C		D l - +:	DM Dadward			N 4 l	Cond Coding 31	tiDI I	David Linius M. Makin
	Concentration, D =	Depletio	on, RM = Reduced	Mat	rix, MS =	Masked	Sand Grains. ² L		Pore Lining, M = Matrix.
Hydric Soil			Delia alice D	ا میں	f 10	:0) /I DD	D MI DA 4 400)		or Problematic Hydric Soils³:
Histoso	i (A1) pipedon (A2)		Polyvalue Be				R, MLRA 149B) a 149B)		uck (A10) (LRR K, L, MLRA 149B)
	istic (A3)		Loamy Muck						rairie Redox (A16) (LRR K, L, R)
	en Sulfide (A4)		Loamy Gleye			(Little)	-,		ucky Peat or Peat (S3) (LRR K, L, R)
	d Layers (A5)		Depleted Ma						rface (S7) (LRR K, L)
Deplete	d Below Dark Surf	ace (A11						•	ie Below Surface (S8) (LRR K, L) rk Surface (S9) (LRR K, L)
Thick D	ark Surface (A12)		Depleted Dar	k Su	rface (F7)			nganese Masses (F12) (LRR K, L, R)
Sandy N	Mucky Mineral (S1)		Redox Depre	ssior	ıs (F8)				nt Floodplain Soils (F19) (MLRA 149B)
Sandy C	Gleyed Matrix (S4)								podic (TA6) (MLRA 144A, 145, 149B)
Sandy F	Redox (S5)								ent Material (F21)
Strippe	d Matrix (S6)								allow Dark Surface (TF12)
Dark Su	ırface (S7) (LRR R, N	/ILRA 14	9B)					-	xplain in Remarks)
3Indicators	of hydrophytic veg	retation	and wetland hydr	olog	v must h	a nrasar	nt unlace disturba	ad or problem:	· atic
	Layer (if observed)		and wedana nyan	0108	y mast b		it, uriless distarbe	ed of problem	auc.
	Type:	•	None			Hydric	Soil Present?		Yes No
	Depth (inches):	-	None	•		liyanc	John Tesent.		103 <u> </u>
Remarks:	Берит (писпез).								
Remarks.									

Project/Site: Pratt South	City/County: Shut	esbury, Franklin	Sampling Date: 2020-July-30			
Applicant/Owner: W.D. Cow	s, Inc.	State: MA	Sampling Point:	W-PMO-03_PEM-1		
Investigator(s): Matt Regan, C	aroline Harrington	Section, Township,	Range:			
Landform (hillslope, terrace, etc	:.): Marsh	Local relief (concave, conv	/ex, none): Concave	Slope (%): 0 to 1		
Subregion (LRR or MLRA):	MLRA 144A of LRR R	Lat: 42.411600030	1 Long: -72.4644833152	Datum: WGS84		
Soil Map Unit Name: Ridgebu	ury fine sandy loam, 3 to 8 percer	nt slopes, extremely stony	NWI classif	fication:		
Are climatic/hydrologic conditio	ns on the site typical for this time		(If no, explain in Rem	arks.)		
Are Vegetation, Soil,			al Circumstances" present?	Yes No		
Are Vegetation, Soil,	or Hydrology naturally	problematic? (If needed,	explain any answers in Rer	narks.)		
SUMMARY OF FINDINGS -	Attach site map showing sa	mpling point locations, trai	nsects, important featu	ires, etc.		
Hydrophytic Vegetation Presen	it? Yes/_ No					
Hydric Soil Present?	Yes No	Is the Sampled Area with	in a Wetland?	Yes No		
		·		W-PMO-03		
Wetland Hydrology Present?	Yes No	If yes, optional Wetland S	ite iD.			
Remarks: (Explain alternative p	rocedures here or in a separate r	eport)				
Covertype is PEM.						
covertype is i Livi.						
HYDROLOGY						
NA/atland I budualant ladiaatawa						
Wetland Hydrology Indicators:	. Constitution of the color of the color		Carandan Indiantan (min	:		
Primary Indicators (minimum c	of one is required; check all that a	pply)	Secondary Indicators (min	•		
<u></u> Surface Water (A1)	Water-Stain	ed Leaves (B9)	Surface Soil Cracks (B6)			
High Water Table (A2)	Aquatic Fau	na (B13)	Drainage Patterns (B10)			
✓ Saturation (A3)	Marl Deposi		Moss Trim Lines (B16)			
Water Marks (B1)	, ,	ulfide Odor (C1)	Dry-Season Water Table (C2)			
Sediment Deposits (B2)	Oxidized Rh	izospheres on Living Roots (C3)				
	_		Saturation Visible on Aerial Imagery (C9)			
Drift Deposits (B3)		Reduced Iron (C4)	Stunted or Stressed Plants (D1)			
Algal Mat or Crust (B4)		Reduction in Tilled Soils (C6)	✓ Geomorphic Position (I	D2)		
Iron Deposits (B5)	Thin Muck S		Shallow Aquitard (D3)	5 (5.4)		
<u>✓</u> Inundation Visible on Aerial		nin in Remarks)	Microtopographic Relie	et (D4)		
Sparsely Vegetated Concave	e Surface (B8)		✓ FAC-Neutral Test (D5)			
Field Observations:						
Surface Water Present?	Yes No [Depth (inches): 6	_			
Water Table Present?	Yes No [Depth (inches): 0	Wetland Hydrology Preser	nt? Yes No		
Saturation Present?	Yes <u></u> No [Depth (inches): 0				
(includes capillary fringe)			•			
. , , , ,	m gauge, monitoring well, aerial _l	abatas provious inspections) if				
Describe Recorded Data (strea	m gauge, monitoring well, aerial	photos, previous inspections), ii	avallable:			
Remarks:						
nemans.						

·		<u> </u>		Daminanaa Taatuusukshaati		
<u>Tree Stratum</u> (Plot size: <u>30 ft</u>)		Dominant Species?		Dominance Test worksheet: Number of Dominant Species That	.+	
4	% Cover	3pecies:	Status	Are OBL, FACW, or FAC:	2	(A)
1.				Total Number of Dominant Specie		
2				Across All Strata:	. 2	(B)
3				Percent of Dominant Species Tha		
4				Are OBL, FACW, or FAC:	100	(A/B)
5				Prevalence Index worksheet:	-	
6				Total % Cover of:	Multiply I	Bv:
7				OBL species 85	x 1 =	85
	0	= Total Cov	er	FACW species 5	x 2 =	10
Sapling/Shrub Stratum (Plot size:15 ft)				FAC species 0	_ x3=	0
1. <i>Spiraea alba</i>	5	Yes	FACW	FACU species 0	_ x4=	0
2				UPL species 0	_ ^ x 5 =	0
3				Column Totals 90		95 (B)
4.					_ (A) _	93 (b)
5.				Prevalence Index = B/A		
6.				Hydrophytic Vegetation Indicator		
7.	· ——			1- Rapid Test for Hydrophyti	c Vegetation	
	5	= Total Cov	er	✓ 2 - Dominance Test is >50%		
Herb Stratum (Plot size: _ 5 ft)		-		\checkmark 3 - Prevalence Index is \le 3.0		
Persicaria hydropiperoides	65	Yes	OBL	4 - Morphological Adaptatio		supporting
Sparganium eurycarpum	10	No	OBL	data in Remarks or on a separate		
3. Scirpus atrovirens	10	No	OBL	Problematic Hydrophytic Ve		
4.			OBL	¹Indicators of hydric soil and wetl	, .	gy must be
-				present, unless disturbed or prob	lematic	
5.	· ——			Definitions of Vegetation Strata:		
6				Tree – Woody plants 3 in. (7.6 cm)		liameter at
7				breast height (DBH), regardless o		
8				Sapling/shrub – Woody plants les		BH and
9				greater than or equal to 3.28 ft (1		
10				Herb – All herbaceous (non-wood size, and woody plants less than 3		ardiess of
11				Woody vines – All woody vines gr		20 ft in
12				height.	eater triair 5.	20 11 111
	85	= Total Cov	er			
Woody Vine Stratum (Plot size: 30 ft)				Hydrophytic Vegetation Present?	Yes N	0
1						
2						
3.	. <u></u> _					
4.						
	0	= Total Cov	er			
Demonstrat (Include whete words are being an an an area		_				
Remarks: (Include photo numbers here or on a separate	te sneet.)					

	<u> </u>	Features		
(inches) Color (moist)	% Color (moist)	<u>% Type¹</u>	Loc² Texture	Remarks
Type: C = Concentration, D = De	pletion, RM = Reduce	d Matrix, MS =	Masked Sand Grains.	² Location: PL = Pore Lining, M = Matrix.
Hydric Soil Indicators: Histosol (A1) Histic Epipedon (A2) Black Histic (A3) Hydrogen Sulfide (A4) Stratified Layers (A5) Depleted Below Dark Surface Thick Dark Surface (A12) Sandy Mucky Mineral (S1) Sandy Gleyed Matrix (S4) Sandy Redox (S5) Stripped Matrix (S6) Dark Surface (S7) (LRR R, MLR	Thin Dark Su Loamy Muck Loamy Gleys Depleted Ma (A11) Redox Dark Depleted Da Redox Depre	orface (S9) (LRI cy Mineral (F1) ed Matrix (F2) atrix (F3) Surface (F6) rk Surface (F7 essions (F8))	Indicators for Problematic Hydric Soils³: 2 cm Muck (A10) (LRR K, L, MLRA 149B) Coast Prairie Redox (A16) (LRR K, L, R) 5 cm Mucky Peat or Peat (S3) (LRR K, L, R) Dark Surface (S7) (LRR K, L) Polyvalue Below Surface (S8) (LRR K, L) Thin Dark Surface (S9) (LRR K, L) Iron-Manganese Masses (F12) (LRR K, L, R) Piedmont Floodplain Soils (F19) (MLRA 149B) Mesic Spodic (TA6) (MLRA 144A, 145, 149B) Red Parent Material (F21) Very Shallow Dark Surface (TF12) Other (Explain in Remarks)
Restrictive Layer (if observed):	ition and wettand nyu	Tology must b	e present, unless disturi	bed of problematic.
Type: Depth (inches):	None		Hydric Soil Present?	Yes No
Remarks: Due to inundation a clear soil pro nundation, FACW and OBL veget			•	s were assumed to be hydric due to the presence of

Project/Site: Pratt South	Ci	i ty/County: Shutesbu	ry, Franklin	Sampling Dat	:e: 2020-July-30		
Applicant/Owner: W.D. Cowls	, Inc.		State: MA	Sampling Point:	: W-PMO-03_PFO-1		
Investigator(s): Matt Regan, Ca	roline Harring	ton	Section, Township,	Range:			
Landform (hillslope, terrace, etc.)	: Depress	ion	Local relief (concave, conv	vex, none): None	Slope (%): 0 to 1		
Subregion (LRR or MLRA):	ILRA 144A of L	RR R	Lat: 42.411789544	9 Long: -72.464756481	4 Datum: WGS84		
Soil Map Unit Name: Ridgebur	y fine sandy lo	am, 3 to 8 percent slo	pes, extremely stony	NWI classi	fication:		
Are climatic/hydrologic condition	s on the site ty	pical for this time of y	ear? Yes _✓_ No	(If no, explain in Rem	narks.)		
Are Vegetation, Soil,	or Hydrolo	gy significantly d		al Circumstances" present			
Are Vegetation, Soil,	or Hydrolo	gy naturally prob	olematic? (If needed,	explain any answers in Re	marks.)		
SUMMARY OF FINDINGS – A	ttach site m	an showing sampl	ing point locations trai	nsects important feati	ires etc		
	- Sitte III	ap showing sampi	Ting point locations, trai	iscets, important reatt			
Hydrophytic Vegetation Present	? Y	′es / _ No					
Hydric Soil Present?	Y	′es No	Is the Sampled Area with	in a Wetland?	Yes/_ No		
Wetland Hydrology Present?	Υ	′es No	If yes, optional Wetland S	ite ID:	W-PMO-03		
Remarks: (Explain alternative pr							
•	Jedui es Here	or iii a separate repor	·)				
Covertype is PFO.							
HYDROLOGY							
IIIDROLOGI							
Wetland Hydrology Indicators:							
Primary Indicators (minimum of	one is require	d; check all that apply)		Secondary Indicators (mir	nimum of two required)		
				Surface Soil Cracks (B6	•		
Surface Water (A1)		Water-Stained Le		Drainage Patterns (B1	·		
High Water Table (A2)		Aquatic Fauna (B		Moss Trim Lines (B16)			
Saturation (A3)		Marl Deposits (B					
Water Marks (B1)		Hydrogen Sulfide		Dry-Season Water Table (C2) ₃₁ Crayfish Burrows (C8)			
Sediment Deposits (B2)		Oxidized Rhizosp	oheres on Living Roots (C3)				
				Saturation Visible on A			
Drift Deposits (B3)		Presence of Red		Stunted or Stressed Plants (D1)			
Algal Mat or Crust (B4)		Recent Iron Redu	uction in Tilled Soils (C6)	✓ Geomorphic Position (
Iron Deposits (B5)		Thin Muck Surface	ce (C7)	Shallow Aquitard (D3)			
Inundation Visible on Aerial I	magery (B7)	Other (Explain in	Remarks)	Microtopographic Reli	ef (D4)		
Sparsely Vegetated Concave	Surface (B8)			FAC-Neutral Test (D5)			
Field Observations:							
Surface Water Present?	Yes N	No ./ Denth	ı (inches):				
		•	·	-	.a. V		
Water Table Present?	Yes N		(inches):	Wetland Hydrology Prese	ent? Yes No		
Saturation Present?	Yes N	No 🟒 Depth	ı (inches):				
(includes capillary fringe)				-			
		مساسه المساسم المساسم		- veileble.			
Describe Recorded Data (stream	i gauge, moniu	oring well, aerial photo	os, previous inspections), ii	avallable:			
Remarks:							
The state of the s							
l							

	Absoluto	Dominant	Indicator	Dominance Test worksheet:		
Tree Stratum (Plot size: 30 ft)		Dominant Species?	Status	Number of Dominant Species That		
1 Tours sandancia		<u> </u>		Are OBL, FACW, or FAC:	3	(A)
1. Tsuga canadensis	40	Yes	FACU	Total Number of Dominant Species		
2. Acer rubrum	25	Yes	FAC	Across All Strata:	5	(B)
3				Percent of Dominant Species That		
4				Are OBL, FACW, or FAC:	60	(A/B)
5				Prevalence Index worksheet:		
6.				Total % Cover of:	Multiply I	Bv:
7				OBL species 0	x 1 =	0
	65	= Total Cove	er	FACW species 10	x 2 =	20
Sapling/Shrub Stratum (Plot size:15 ft)				FAC species 45	x3=	135
1. Tsuga canadensis	25	Yes	FACU	FACU species 65	x 4 =	260
2. Frangula alnus	20	Yes	FAC	UPL species 0	x5=	0
3.					_	
4.				Column Totals 120	(A) _	415 (B)
5.	· ——			Prevalence Index = B/A =	3.5	
6.				Hydrophytic Vegetation Indicators:		
7.				1- Rapid Test for Hydrophytic	/egetation	
/·	45	= Total Cove	or	✓ 2 - Dominance Test is >50%		
Herb Stratum (Plot size: _ 5 ft)	43	_ TOTAL COV		3 - Prevalence Index is $\leq 3.0^{1}$		
	10	Voc	ΓΛ <i>C</i> \Λ/	4 - Morphological Adaptations	1 (Provide :	supporting
1. Impatiens capensis	10	Yes	FACW	data in Remarks or on a separate sl	neet)	
2.				Problematic Hydrophytic Vege	tation¹ (Ex	plain)
3				landicators of hydric soil and wetlar	d hydroloខ្	gy must be
4				present, unless disturbed or proble	matic	
5				Definitions of Vegetation Strata:		
6				Tree – Woody plants 3 in. (7.6 cm) o	r more in c	liameter at
7				breast height (DBH), regardless of h	eight.	
8.				Sapling/shrub – Woody plants less t	han 3 in. D	BH and
9.				greater than or equal to 3.28 ft (1 m	ı) tall.	
10.				Herb – All herbaceous (non-woody)	plants, reg	gardless of
11.	· ——			size, and woody plants less than 3.2	8 ft tall.	
12.				Woody vines – All woody vines grea	ter than 3.	28 ft in
	10	= Total Cove	ar	height.		
Woody Vine Stratum (Plot size:30 ft)		_ Total Cov	-1	Hydrophytic Vegetation Present?	YesN	0
1.						
·						
2.						
3						
4						
	0	= Total Cove	er			
Remarks: (Include photo numbers here or on a separat	e sheet.)					
	•					

	cription: (Describe	to the de	•			indicato	r or confirm the ab	sence of ind	icators.)
Depth _	Matrix		Redox				- .		
(inches)	Color (moist)	<u>%</u>	Color (moist)	<u>%</u>	Type ¹	Loc ²	Texture		Remarks
0 - 3	10YR 3/1	100	4000 745	· <u> </u>			Loamy Sa		
3 - 10	10YR 4/3	95	10YR 5/6	5	C	M	Sandy Lo	am	
				-					
				-					
				_					
				-					
				_					
				- —					
				- —					
				- —					
		- —							
		. —							
	Concentration, D =	Depletio	n, RM = Reduced	Mat	rix, MS =	Masked	Sand Grains. ² Lo	ocation: PL =	Pore Lining, M = Matrix.
Hydric Soil								Indicators fo	or Problematic Hydric Soils³:
Histoso			•				R, MLRA 149B)	2 cm Mu	uck (A10) (LRR K, L, MLRA 149B)
	pipedon (A2)		Thin Dark Su					Coast Pr	rairie Redox (A16) (LRR K, L, R)
	istic (A3) en Sulfide (A4)		Loamy Muck			(LKK K, I	L)	5 cm Mu	ucky Peat or Peat (S3) (LRR K, L, R)
	d Layers (A5)		Depleted Ma						rface (S7) (LRR K, L)
	d Below Dark Surfa	ace (A11						•	ue Below Surface (S8) (LRR K, L)
	ark Surface (A12)		Depleted Dar)			rk Surface (S9) (LRR K, L)
Sandy N	lucky Mineral (S1)		Redox Depre	ssior	ıs (F8)				nganese Masses (F12) (LRR K, L, R)
Sandy C	Gleyed Matrix (S4)								nt Floodplain Soils (F19) (MLRA 149B) podic (TA6) (MLRA 144A, 145, 149B)
Sandy F	Redox (S5)								ent Material (F21)
Stripped	d Matrix (S6)								allow Dark Surface (TF12)
Dark Su	rface (S7) (LRR R, N	/ILRA 149	9B)					-	Explain in Remarks)
Indicators	of hydrophytic veg	otation	and watland by	-olog	, must b	0 050505	at unlace disturba		
	Layer (if observed):		and wettand nydi	olog	y must b	e preser	it, uriless disturbed	u or problem	auc.
	Type:		None			Hydric	Soil Present?		Yes/_ No
	Depth (inches):	-	None	•		liyanc	John Frederic.		163 <u>-</u> 4 NO
Remarks:	Deptil (ilicites).								
According t	o the USDA NRCS t	the map	ped soil type is cl	assifi	ed as hy	dric.			

Project/Site: Pratt South	City/County: Shu	tesbury, Hampshire	Sampling Date: 2020-July-30			
Applicant/Owner: W.D. Cov	vls, Inc.	State: MA	Sampling Point	: W-PMO-03_PSS-1		
Investigator(s): Matt Regan,	Caroline Harrington	Section, Township,	Range:			
Landform(hillslope,terrace,etc	:.): Swamp	Local relief (concave, conv	ex, none): Concave	Slope (%): 0 to 1		
Subregion(LRRorMLRA):	MLRA 144A of LRR R	Lat: 42.410388887	Long: -72.464897465	Datum: WGS84		
SoilMapUnitName: Hinckle	y loamy sand, 8 to 15 percent slopes		NWI class	ification:		
Areclimatic/hydrologiccondition	onsonthesitetypicalforthistimeofyear?	Yes 🟒 No	(If no, explain in Ren	narks.)		
Are Vegetation, Soil	_, or Hydrology significantly di	sturbed? Are "Norm	al Circumstances" present	? Yes _ ✓ No		
Are Vegetation, Soil	_, or Hydrology naturally prob	lematic? (If needed,	explain any answers in Re	marks.)		
SUMMARY OF FINDINGS -	- Attach site map showing sampli	ng noint locations trai	nsects important feat	ures etc		
Hydrophytic Vegetation Prese						
Hydric Soil Present?	Yes _ 🗸 No	Is the Sampled Area withi	n a Wetland?	Yes No		
	Yes _ ✓ _ No	If yes, optional Wetland S		W-PMO-03		
Wetland Hydrology Present?	procedures here or in a separate report		ite iD.	<u>W-PIVIO-03</u>		
HYDROLOGY						
HYDROLOGY						
Wetland Hydrology Indicators	:					
Primary Indicators (minimum	of one is required; check all that apply)		Secondary Indicators (mir	nimum of two required)		
Surface Water (A1)	Water-Stained Lea	aves (B9)	Surface Soil Cracks (B6)			
High Water Table (A2)	Aquatic Fauna (B1		Drainage Patterns (B10)			
✓ Saturation (A3)	Marl Deposits (B1	5)	Moss Trim Lines (B16)			
Water Marks (B1)	Hydrogen Sulfide		Dry-Season Water Table (C2)			
Sediment Deposits (B2)	Oxidized Rhizospl	heres on Living Roots (C3)	 Crayfish Burrows (C8) Saturation Visible on Aerial Imagery (C9) 			
Drift Deposits (B3)	Presence of Redu	cod Iron (CA)				
Algal Mat or Crust (B4)		ction in Tilled Soils (C6)	Stunted or Stressed Plants (D1) Geomorphic Position (D2)			
Iron Deposits (B5)	Thin Muck Surface		Shallow Aquitard (D3)			
Inundation Visible on Aeria	al Imagery (B7) Other (Explain in I	Remarks)	Microtopographic Reli			
Sparsely Vegetated Concav	ve Surface (B8)		✓ FAC-Neutral Test (D5)			
Field Observations:						
Surface Water Present?	Yes No 🟒 Depth	(inches):				
Water Table Present?	Yes No 🟒 Depth	(inches):	Wetland Hydrology Prese	ent? Yes No		
Saturation Present?	Yes <u></u> ✓ No Depth	(inches): 0				
(includes capillary fringe)			-			
Describe Recorded Data (streated) Remarks:	am gauge, monitoring well, aerial photo	s, previous inspections), if	available:			

Tree Stratum (Plot size: 30 ft)		Dominant Species?	Indicator Status	Dominance Test worksheet Number of Dominant Speciare OBL, FACW, or FAC:		4	(A)
 2.				Total Number of Dominant Across All Strata:	Species	4	(B)
3.				Percent of Dominant Specie Are OBL, FACW, or FAC:	es That	100	(A/B)
5				Prevalence Index workshee	et:		
j				Total % Cover of:		Multiply E	<u>By:</u>
				OBL species	75	x 1 =	75
	0	= Total Cov	er	FACW species	75	x 2 =	150
apling/Shrub Stratum (Plot size: <u>15 ft</u>)				FAC species	25	x 3 =	75
. Alnus incana	30	Yes	FACW	FACU species	0	x 4 =	0
. Frangula alnus	25	Yes	FAC	UPL species	0	x 5 =	0
3. <u>Aronia arbutifolia</u>	10	No	FACW	Column Totals	175	(A)	300 (B)
. Spiraea alba	5	No	FACW	Prevalence Index		_	,
i				-			
i				Hydrophytic Vegetation Ind		ogotation	
'				2 - Dominance Test is		egetation	
	70	= Total Cov	er	✓ 3 - Prevalence Index is			
lerb Stratum (Plot size: <u>5 ft</u>)						(Provide c	unnorting
. Carex crinita	60	Yes	OBL	4 - Morphological Ada data in Remarks or on a se			upporting
. Impatiens capensis	30	Yes	FACW	Problematic Hydroph			nlain)
3. Scirpus atrovirens	10	No	OBL	Indicators of hydric soil an	_		
1. Typha latifolia	5	No	OBL	present, unless disturbed of		, .	y must be
5.				Definitions of Vegetation St		nacie	
-				Tree – Woody plants 3 in. (7		more in d	iameter a
				breast height (DBH), regard			idifficter d
				Sapling/shrub - Woody plan			BH and
				greater than or equal to 3.2			2
				Herb – All herbaceous (non			ardless of
11				size, and woody plants less			
				Woody vines – All woody vi			28 ft in
2		Tatal Car		height.			
	105	= Total Cov	er	Hydrophytic Vegetation Pr	esent? Y	es / No	1
Noody Vine Stratum (Plot size: 30 ft)				injuroprijas regetation r			
·							
2.							
3.							
4.		= Total Cov	er				

Profile Des Depth	cription: (Describe t Matrix	to the d	epth needed to d Redox			indicato	r or confirm the a	bsence of ind	licators.)
(inches)	Color (moist)		Color (moist)	%	Type ¹	Loc ²	Textur	e	Remarks
0 - 20	2.5Y 2.5/1	100		- <u>~</u> 	.,,,,,		Mucky P	-	
				_	-				
		_		_					
				-					
				_					
				-					
				_					
				- —					
1Type: C = 0	Concentration, D =	 Depletio	on, RM = Reduced	Mati	rix, MS =	 Masked	Sand Grains. ² L	ocation: PL =	Pore Lining, M = Matrix.
Hydric Soil	Indicators:	-							for Problematic Hydric Soils³:
/ Histoso			-				R, MLRA 149B)	2 cm M	uck (A10) (LRR K, L, MLRA 149B)
	oipedon (A2) istic (A3)		Thin Dark Su Loamy Muck						rairie Redox (A16) (LRR K, L, R)
	en Sulfide (A4)		Loamy Gleye			(=:::::	-,		ucky Peat or Peat (S3) (LRR K, L, R)
Stratifie	d Layers (A5)		Depleted Ma						ırface (S7) (LRR K, L) ue Below Surface (S8) (LRR K, L)
	d Below Dark Surfa	ace (A11) Redox Dark S	Surfa	ce (F6)				rk Surface (S9) (LRR K, L)
	ark Surface (A12)		Depleted Dar)			anganese Masses (F12) (LRR K, L, R)
	Mucky Mineral (S1)		Redox Depre	ssior	ıs (F8)				ont Floodplain Soils (F19) (MLRA 149B)
-	Gleyed Matrix (S4)								podic (TA6) (MLRA 144A, 145, 149B)
Sandy F	Redox (S5)								rent Material (F21)
Strippe	d Matrix (S6)								allow Dark Surface (TF12)
Dark Su	ırface (S7) (LRR R, M	1LRA 14	9B)					-	Explain in Remarks)
	of hydrophytic veg		and wetland hydi	olog	y must b	e preser	nt, unless disturbe	ed or problem	natic.
Restrictive	Layer (if observed):		Mana			U to contract on	C-11 B		Ver de Ne
	Type:		None	-		Hydric	Soil Present?		Yes No
	Depth (inches):								
Remarks:									

Project/Site: Pratt South	City/County:_ Shu	tesbury, Franklin	Sampling Date: 2020-July-30			
Applicant/Owner: W.D. Cowls, I	inc.	State: MA	Sampling Point: W-PMO-03_UPL-1			
Investigator(s): Matt Regan, Car	oline Harrington	Section, Township, I	Range: Hinckley loamy sand, 8 to 15 percent slo			
Landform(hillslope,terrace,etc.):	Hillslope	Local relief (concave, conve	ex, none): Convex Slope (%): 1 t			
Subregion(LRRorMLRA): MLF	RA 144A of LRR R	Lat: 42.4105282361	Long: -72.4644694851 Datum: WGS8			
SoilMapUnitName: 42.4105282	2361		NWI classification:			
Areclimatic/hydrologicconditions	onthesitetypicalforthistimeofyear?	Yes No	(If no, explain in Remarks.)			
Are Vegetation, Soil,	or Hydrology significantly di	sturbed? Are "Norma	al Circumstances" present? Yes 🟒 No			
Are Vegetation, Soil,	or Hydrology naturally prob	lematic? (If needed, e	explain any answers in Remarks.)			
SUMMARY OF FINDINGS – At	tach site map showing sampli	ng point locations, tran	sects, important features, etc.			
Hydrophytic Vegetation Present?	Yes No _ ✓					
Hydric Soil Present?	Yes No	Is the Sampled Area withir	n a Wetland? Yes No/			
	Yes No _ _ ✓	· ·				
Wetland Hydrology Present?		If yes, optional Wetland Sit	te iD:			
	cedures here or in a separate report)				
Covertype is UPL.						
HYDROLOCY						
HYDROLOGY						
Wetland Hydrology Indicators:						
Primary Indicators (minimum of o	one is required; check all that apply)		Secondary Indicators (minimum of two required			
Surface Water (A1)	Water-Stained Lea	aves (B9)	Surface Soil Cracks (B6)			
High Water Table (A2)	Aquatic Fauna (B1	13)	Drainage Patterns (B10)			
Saturation (A3)	Marl Deposits (B1	5)	Moss Trim Lines (B16)			
Water Marks (B1)	Hydrogen Sulfide	Odor (C1)	Dry-Season Water Table (C2)			
Sediment Deposits (B2)	Oxidized Rhizospl	heres on Living Roots (C3)	Crayfish Burrows (C8)			
- 10 - 11 (Saturation Visible on Aerial Imagery (C9)			
Drift Deposits (B3)	Presence of Redu		Stunted or Stressed Plants (D1)			
Algal Mat or Crust (B4)		ction in Tilled Soils (C6)	Geomorphic Position (D2)			
Iron Deposits (B5) Inundation Visible on Aerial Im	Thin Muck Surface		Shallow Aquitard (D3)			
Sparsely Vegetated Concave S	· · · · · · · · · · · · · · · · · · ·	remarks)	 Microtopographic Relief (D4) FAC-Neutral Test (D5)			
Field Observations:	diface (DO)					
Surface Water Present?	Yes No <u></u> Depth	(inches):				
		· —	Mathematikadasha - Durasa 2			
Water Table Present?	'		Wetland Hydrology Present? Yes No _			
Saturation Present?	Yes No Depth	(inches):				
(includes capillary fringe)						
Describe Recorded Data (stream	gauge, monitoring well, aerial photo	s, previous inspections), if a	vailable:			
Remarks:						

<u>Free Stratum</u> (Plot size: <u>30 ft</u>)		Dominant Species?	Indicator Status	Dominance Test worksh Number of Dominant S Are OBL, FACW, or FAC:	pecies That	2	(A)
 2.				Total Number of Domin		6	(B)
3.				Percent of Dominant Sp Are OBL, FACW, or FAC:		33.3	(A/B)
5				Prevalence Index works	heet:		
i				Total % Cover	<u>of:</u>	Multiply E	<u>Зу:</u>
'				OBL species	0	x 1 =	0
	0	= Total Cov	er	FACW species	0	x 2 =	0
apling/Shrub Stratum (Plot size: 15 ft)				FAC species	15	x 3 =	45
. Frangula alnus	10	Yes	FAC	- FACU species	80	x 4 =	320
. Acer rubrum	5	Yes	FAC	- UPL species	15	x 5 =	75
. Quercus alba	5	Yes	FACU	Column Totals	110	(A)	440 (B)
. Pinus strobus	5	Yes	FACU	Prevalence In	dex = B/A =	4	- ()
·				Hydrophytic Vegetation			
i				1- Rapid Test for H		lagatation	
<i>'</i>				·		regetation	
	25	= Total Cov	er	2 - Dominance Tes			
<u>lerb Stratum</u> (Plot size: <u>5 ft</u>)	-	_		3 - Prevalence Ind			
. Kalmia latifolia	30	Yes	FACU	4 - Morphological			supporting
2. Vaccinium angustifolium	30	Yes	FACU	data in Remarks or on a			-1-1-1
. Comptonia peregrina	15	No	UPL	Problematic Hydro			
l. Solidago canadensis	10	No	FACU	¹Indicators of hydric soi		, .	gy must be
5.		110	17100	present, unless disturbe		matic	
5.				Definitions of Vegetatio			
-				Tree – Woody plants 3 i			liameter a
7.				breast height (DBH), reg			D
3.				Sapling/shrub - Woody			BH and
)				greater than or equal to			
0				Herb – All herbaceous (ardiess of
1				size, and woody plants			20 ft in
2				Woody vines – All wood	y viries grea	ter triari 5	20 11 111
	85	= Total Cov	er	height.			
Noody Vine Stratum (Plot size: 30 ft)				Hydrophytic Vegetation	n Present? \	Yes N	0
l.							
<u> </u>							
3.				•			
1.				•			
··· -		= Total Cov	or	•			
		_ Total Cov	CI				

Profile Desc Depth	cription: (Describe Matrix	to the d	epth needed to do Redox			indicato	r or confirm the	absence of ind	icators.)
(inches)	Color (moist)	%	Color (moist)		Type ¹	Loc ²	Tev	ture	Remarks
0 - 3	10YR 3/2	100	Color (moist)	70	туре	LOC	-	/ Loam	Remarks
				_					
3 - 10	10YR 4/4	100		_			Loam	y Sand	
				_					
				_					
				_					
				_					
				_					
				_					
				_					
				_					
¹Type: C = C	Concentration, D =	Depletic	on, RM = Reduced	Mati	rix, MS =	Masked	Sand Grains.	² Location: PL =	Pore Lining, M = Matrix.
Hydric Soil		•							or Problematic Hydric Soils ³ :
Histosol			Polyvalue Bel	ow S	urface (S	8) (LRR	R. MLRA 149B)		·
	pipedon (A2)		Thin Dark Sur		-				uck (A10) (LRR K, L, MLRA 149B)
Black Hi			Loamy Mucky						rairie Redox (A16) (LRR K, L, R)
	en Sulfide (A4)		Loamy Gleyed	d Ma	trix (F2)				ucky Peat or Peat (S3) (LRR K, L, R) rface (S7) (LRR K, L)
Stratifie	d Layers (A5)		Depleted Mat	rix (I	- 3)				ie Below Surface (S8) (LRR K, L)
Deplete	d Below Dark Surf	ace (A11) Redox Dark S	urfa	ce (F6)				rk Surface (S9) (LRR K, L)
	ark Surface (A12)		Depleted Dar	k Sui	rface (F7)			nganese Masses (F12) (LRR K, L, R)
	lucky Mineral (S1)		Redox Depres	ssior	ıs (F8)				nt Floodplain Soils (F19) (MLRA 149B)
Sandy G	Gleyed Matrix (S4)								podic (TA6) (MLRA 144A, 145, 149B)
Sandy R	tedox (S5)								ent Material (F21)
Stripped	d Matrix (S6)								allow Dark Surface (TF12)
Dark Su	rface (S7) (LRR R, N	/ILRA 14	9B)					-	explain in Remarks)
21	- 6 lea educe de lea 48 e a ce								•
	of hydrophytic veg		and welland nydr	olog	y must b	e preser	it, uniess distur	bed or problem	auc.
	Layer (if observed): _								
	Type:		None			Hydric	Soil Present?		Yes No⁄_
-	Depth (inches):								
Remarks:									

Project/Site: Pratt South	City/County: Shut	esbury, Franklin	Sampling Date: 202	20-July-30
Applicant/Owner: W.D. Cowls	, Inc.	State: MA	Sampling Point: W-PN	ИО-03_UPL-2
Investigator(s): Matt Regan, Ca	aroline Harrington, Caroline Harringto	n Section, Township,	Range:	
Landform(hillslope,terrace,etc.):	Hillslope	Local relief (concave, conv	ex, none): Convex	Slope (%): 1 to 3
Subregion(LRRorMLRA): MI	LRA 144A of LRR R	Lat: 42.411710335	9 Long: -72.4644889311	Datum: WGS84
SoilMapUnitName:			NWI classificatio	n:
Areclimatic/hydrologicconditions	sonthesitetypicalforthistimeofyear?	Yes No	(If no, explain in Remarks.)	
Are Vegetation, Soil,	or Hydrology significantly dis	sturbed? Are "Norm	al Circumstances" present?	Yes No
Are Vegetation, Soil,	or Hydrology naturally probl	ematic? (If needed,	explain any answers in Remarks.)
SUMMARY OF FINDINGS – A	Attach site map showing samplir	ng point locations, trar	nsects, important features, o	etc.
Hydrophytic Vegetation Present	? Yes No ⁄ _			
Hydric Soil Present?	Yes No _∠ _	Is the Sampled Area withi	n a Wetland? Yes	i No _∠
		If yes, optional Wetland S		
Wetland Hydrology Present?	Yes No _ _ _	3 . 1	ite ib.	
·	ocedures here or in a separate report)			
Covertype is UPL.				
HYDROLOGY				
Wetland Hydrology Indicators:				
Primary Indicators (minimum of	one is required; check all that apply)		Secondary Indicators (minimum	of two required)
Surface Water (A1)	Water-Stained Lea	ives (B9)	Surface Soil Cracks (B6)	
High Water Table (A2)	Aquatic Fauna (B1	3)	Drainage Patterns (B10)	
Saturation (A3)	Marl Deposits (B1	5)	Moss Trim Lines (B16)	
Water Marks (B1)	Hydrogen Sulfide		Dry-Season Water Table (C2)	
Sediment Deposits (B2)	Oxidized Rhizosph	neres on Living Roots (C3)	Crayfish Burrows (C8)	(60)
			Saturation Visible on Aerial I	
Drift Deposits (B3)	Presence of Reduc		Stunted or Stressed Plants ([01)
Algal Mat or Crust (B4)		tion in Tilled Soils (C6)	Geomorphic Position (D2)	
Iron Deposits (B5)	Thin Muck Surface		Shallow Aquitard (D3)	
Inundation Visible on Aerial I		Remarks)	Microtopographic Relief (D4)	1
Sparsely Vegetated Concave	Surface (B8)		FAC-Neutral Test (D5)	
Field Observations:				
Surface Water Present?	•	(inches):		
Water Table Present?	Yes No Depth ((inches):	Wetland Hydrology Present?	Yes No / _
Saturation Present?	Yes No Depth ((inches):		
(includes capillary fringe)				
Describe Recorded Data (stream	n gauge, monitoring well, aerial photos	s, previous inspections), if a	available:	
	- 88-,	,, p		
Remarks:				

ree Stratum (Plot size: <u>30 ft</u>)		Dominant Species?	Indicator Status	Dominance Test worksheet: Number of Dominant Species Tha	at 1	(4)
. Pinus strobus	20	Yes	FACU	Are OBL, FACW, or FAC:		(A)
. Quercus rubra	15	Yes	FACU	Total Number of Dominant Specie	es 8	(B)
. Acer rubrum	10	Yes	FAC	Across All Strata:		(D)
				Percent of Dominant Species Tha	t 12.5	(A/B)
5.				Are OBL, FACW, or FAC:		`
				Prevalence Index worksheet:		
				Total % Cover of:	<u>Multiply E</u>	-
	45	= Total Cov	er	- OBL species 0	_ x 1 = _	0
apling/Shrub Stratum (Plot size:15 ft)			·.	FACW species 0	x 2 =	0
. Tsuga canadensis	25	Yes	FACU	FAC species 10	_ x 3 = _	30
Betula lenta	10	Yes	FACU	FACU species 100	x 4 =	400
. Quercus rubra		No	FACU	- UPL species0	x 5 =	0
		INU	FACU	Column Totals 110	(A)	430 (B)
•				Prevalence Index = B/A	= 3.9	
•				Hydrophytic Vegetation Indicator	s:	
				1- Rapid Test for Hydrophyti		
•				2 - Dominance Test is > 50%		
	40	= Total Cov	er	3 - Prevalence Index is ≤ 3.0	1	
lerb Stratum (Plot size: <u>5 ft</u>)				4 - Morphological Adaptatio		upporting
. Maianthemum canadense	10	Yes	FACU	data in Remarks or on a separate		аррот с
. Vaccinium angustifolium	10	Yes	FACU	Problematic Hydrophytic Ve		olain)
. Dendrolycopodium obscurum	5	Yes	FACU	Indicators of hydric soil and wetl	-	
				present, unless disturbed or prob	, .	,asc sc
i				Definitions of Vegetation Strata:		
				Tree – Woody plants 3 in. (7.6 cm)	or more in d	ismotor s
,				breast height (DBH), regardless o		iairietei a
				Sapling/shrub – Woody plants les	_	RH and
·				greater than or equal to 3.28 ft (1		Dirana
				Herb – All herbaceous (non-wood		ardless of
0				size, and woody plants less than 3		ai aic35 0i
1				Woody vines – All woody vines gr		28 ft in
2				height.		
	25	= Total Cov	er		. Voc. N	
Voody Vine Stratum (Plot size: <u>30 ft</u>)				Hydrophytic Vegetation Present?	res IN	J
				_		
·				_		
				1		
				-		

	cription: (Describe	to the de	-			indicato	r or confirm the at	osence of indi	icators.)
Depth _	Matrix		Redox						
(inches)	Color (moist)	<u>%</u>	Color (moist)	<u>%</u>	Type ¹	Loc ²	Textur		Remarks
0 - 2	7.5YR 3/2	100		_			Loamy S		
2 - 8	10YR 4/4	100		_			Loamy S	and	
				_					
				_					
				_					
				_					
				_					
				_					
¹Type: C = 0	Concentration, D =	Depletio	n, RM = Reduced	Mat	rix, MS =	Masked	Sand Grains. ² Lo	ocation: PL = I	Pore Lining, M = Matrix.
Hydric Soil	Indicators:							Indicators fo	or Problematic Hydric Soils³:
Histoso			Polyvalue Be	low S	urface (S	8) (LRR	R, MLRA 149B)		ick (A10) (LRR K, L, MLRA 149B)
Histic E	oipedon (A2)		Thin Dark Su						rairie Redox (A16) (LRR K, L, R)
Black Hi	istic (A3)		Loamy Muck	y Mir	eral (F1)	(LRR K,	L)		icky Peat or Peat (S3) (LRR K, L, R)
	en Sulfide (A4)		Loamy Gleye						face (S7) (LRR K, L)
	d Layers (A5)		Depleted Ma						e Below Surface (S8) (LRR K, L)
	d Below Dark Surfa	ace (A11)	· 					-	k Surface (S9) (LRR K, L)
	ark Surface (A12)		Depleted Dar)			nganese Masses (F12) (LRR K, L, R)
	Mucky Mineral (S1)		Redox Depre	ssior	is (F8)			Piedmor	nt Floodplain Soils (F19) (MLRA 149B)
-	Gleyed Matrix (S4)							Mesic Sp	oodic (TA6) (MLRA 144A, 145, 149B)
_	Redox (S5)							Red Pare	ent Material (F21)
	d Matrix (S6)							Very Sha	allow Dark Surface (TF12)
Dark Su	ırface (S7) (LRR R, N	/ILRA 149	9B)					Other (E	xplain in Remarks)
3Indicators	of hydrophytic veg	etation a	and wetland hydr	olog	y must b	e preser	nt, unless disturbe	d or problema	atic.
Restrictive	Layer (if observed):								
	Type:		Rock			Hydric	Soil Present?		Yes No/_
	Depth (inches):		8						
Remarks:									
Ì									

Project/Site: Pratt South	City/County: Shu	tesbury, Franklin	Sampling Date: 202	20-July-30
Applicant/Owner: W.D. Cowls	, Inc.	State: MA	Sampling Point: W-PI	MO-03_UPL-3
Investigator(s): Matt Regan, Ca	aroline Harrington	Section, Township,	Range:	
Landform(hillslope,terrace,etc.):	Hilltop	Local relief (concave, conv	ex, none): Convex	Slope (%): 1 to 3
Subregion(LRRorMLRA): M	LRA 144A of LRR R	Lat: 42.412035008	9 Long: -72.4646558986	_Datum: WGS84
SoilMapUnitName: Hinckley l	oamy sand, 8 to 15 percent slopes		NWI classificatio	n:
Areclimatic/hydrologicconditions	sonthesitetypicalforthistimeofyear?	Yes 🔽 No	(If no, explain in Remarks.)	
Are Vegetation, Soil,	or Hydrology significantly di		· ·	Yes No
Are Vegetation, Soil,	or Hydrology naturally prob	lematic? (If needed,	explain any answers in Remarks	.)
Summary of Findings – A	Attach site map showing sampli	ng point locations, trar	nsects, important features,	etc.
Hydrophytic Vegetation Present	? Yes No _ _/ _			
Hydric Soil Present?	Yes No _ _ _	Is the Sampled Area withi	n a Wetland? Yes	s No⁄_
Wetland Hydrology Present?	Yes No	If yes, optional Wetland S		
			itte iD.	
	ocedures here or in a separate report	I)		
Covertype is UPL.				
HYDROLOGY				
Wetland Hydrology Indicators:				
Primary Indicators (minimum of	one is required; check all that apply)		Secondary Indicators (minimum	of two required)
Surface Water (A1)	Water-Stained Le		Surface Soil Cracks (B6)	
High Water Table (A2)	Aquatic Fauna (B´		Drainage Patterns (B10)	
Saturation (A3)	Marl Deposits (B1		Moss Trim Lines (B16) Dry-Season Water Table (C2)	
Water Marks (B1)	Hydrogen Sulfide		Crayfish Burrows (C8)	1
Sediment Deposits (B2)	Oxidized Rhizosp	heres on Living Roots (C3)	Saturation Visible on Aerial I	magery (C9)
Drift Deposits (B3)	Presence of Redu	ced Iron (C4)	Stunted or Stressed Plants (I	
Algal Mat or Crust (B4)		ction in Tilled Soils (C6)	Geomorphic Position (D2)	51,
Iron Deposits (B5)	Thin Muck Surfac		Shallow Aquitard (D3)	
Inundation Visible on Aerial I			Microtopographic Relief (D4)
Sparsely Vegetated Concave	Surface (B8)		FAC-Neutral Test (D5)	
Field Observations:				
Surface Water Present?	Yes No _ _/ Depth	(inches):		
Water Table Present?	Yes No <u></u> ✓ Depth	(inches):	- Wetland Hydrology Present?	Yes No ∠
Saturation Present?		(inches):		
(includes capillary fringe)	163 140 _ Beptil		-	
		\(\dot\) in\(\dot\) if (
Describe Recorded Data (stream	n gauge, monitoring well, aerial photo	s, previous inspections), ii a	avallable:	
Remarks:				

	Ahsolute	Dominant	Indicator	Dominance Test worksheet:		
Tree Stratum (Plot size: 30 ft)		Species?	Status	Number of Dominant Species That	_	
1. Pinus strobus	25	Yes	FACU	Are OBL, FACW, or FAC:	2	(A)
2. Tsuga canadensis	15	Yes	FACU	Total Number of Dominant Species	6	(D)
3. Acer rubrum	10	Yes	FAC	Across All Strata:		(B)
4.		103	1710	Percent of Dominant Species That	33.3	(A/B)
5.				Are OBL, FACW, or FAC:		(// b)
6.				Prevalence Index worksheet:		
7.				Total % Cover of:	Multiply	<u>Ву:</u>
/		= Total Cov		OBL species 0	x 1 =	0
Continue/Charab Characterize (Diet sings 15 ft)	50	_ 10tal Cov	er	FACW species 0	x 2 =	0
Sapling/Shrub Stratum (Plot size:15 ft)	50	V	FACIL	FAC species 20	x 3 =	60
1. Tsuga canadensis	50	Yes	FACU	FACU species 120	x 4 =	480
2. Betula lenta	30	Yes	FACU	- UPL species 0	x 5 =	0
3				- Column Totals 140	(A)	540 (B)
4				Prevalence Index = B/A =	3.9	
5				Hydrophytic Vegetation Indicators:		
6				1- Rapid Test for Hydrophytic \	/egetation	
7				2 - Dominance Test is > 50%		
	80	= Total Cov	er	3 - Prevalence Index is $\leq 3.0^{\circ}$		
Herb Stratum (Plot size:5 ft)				4 - Morphological Adaptations	1 (Provide	sunnorting
1. Athyrium angustum	10	Yes	FAC	data in Remarks or on a separate sh		sapporting
2				Problematic Hydrophytic Vege		plain)
3				Indicators of hydric soil and wetlan		-
4				present, unless disturbed or proble		
5.				Definitions of Vegetation Strata:		
6.				Tree – Woody plants 3 in. (7.6 cm) o	r more in o	diameter at
7.				breast height (DBH), regardless of h		
8.				Sapling/shrub – Woody plants less t	han 3 in. D	BH and
9.				greater than or equal to 3.28 ft (1 m) tall.	
10.				Herb – All herbaceous (non-woody)	plants, reg	gardless of
11.				size, and woody plants less than 3.2	8 ft tall.	
12.				Woody vines – All woody vines grea	ter than 3.	28 ft in
	10	= Total Cov	er	height.		
Woody Vine Stratum (Plot size:30 ft)				Hydrophytic Vegetation Present?	Yes N	0
1.						
2.				-		
3.				-		
4.				-		
*		= Total Cov	or	-		
		_ TOTAL COV	<u>تا</u>			
Remarks: (Include photo numbers here or on a separa	te sheet.)					

Profile Des	cription: (Describe Matrix	to the d	epth needed to d Redox			indicato	r or confirm the a	bsence of indi	cators.)
(inches)	Color (moist)	%	Color (moist)	%	Type ¹	Loc ²	Textu	re	Remarks
0 - 2	10YR 3/2	100	Color (moist)		Турс		Loamy S		- Normania
2 - 8	10YR 4/4	100		_			Loamy S		_
				_					
				_					
			-	_	-				
				_					
				_					-
				_					-
				_					
				_					
				_					
				_					
1Type: C = C	Concentration, D =	Denletic	n RM = Reduced	Mat	rix MS =	Masked	Sand Grains 21	ocation: PL = F	Pore Lining, M = Matrix.
Hydric Soil		Depletic	ni, Kivi – Keduced	iviat	11, 1013 -	Maskea	Sana Grains. L		or Problematic Hydric Soils ³ :
Histoso			Polyvalua Bo	OW/ S	urface (9	(8) (I DD I	R, MLRA 149B)		·
	oipedon (A2)		Thin Dark Su						ick (A10) (LRR K, L, MLRA 149B)
Black Hi	•		Loamy Muck						rairie Redox (A16) (LRR K, L, R)
Hydroge	en Sulfide (A4)		Loamy Gleye	d Ma	trix (F2)				icky Peat or Peat (S3) (LRR K, L, R) face (S7) (LRR K, L)
Stratifie	d Layers (A5)		Depleted Ma	trix (l	- 3)				e Below Surface (S8) (LRR K, L)
	d Below Dark Surf	ace (A11							k Surface (S9) (LRR K, L)
	ark Surface (A12)		Depleted Dar)			nganese Masses (F12) (LRR K, L, R)
	lucky Mineral (S1)		Redox Depre	ssior	ıs (F8)				nt Floodplain Soils (F19) (MLRA 149B)
-	Gleyed Matrix (S4)								podic (TA6) (MLRA 144A, 145, 149B)
_	ledox (S5)							Red Pare	ent Material (F21)
	d Matrix (S6)								allow Dark Surface (TF12)
Dark Su	rface (S7) (LRR R, N	MLRA 14	9B)					Other (E	xplain in Remarks)
3Indicators	of hydrophytic veg	getation	and wetland hydr	olog	y must b	e preser	nt, unless disturbe	ed or problema	atic.
Restrictive	Layer (if observed)	:				Ī		•	
	Type:		Rock			Hydric	Soil Present?		Yes No/_
	Depth (inches):		8						
Remarks:									
i									
1									
1									
1									

Project/Site: Pratt South	City/County: Shutesbur	y, Franklin	Sampling Date:	2020-Aug-03
Applicant/Owner: W.D. Cowls,	Inc.	State: MA	Sampling Point: V	V-PMO-04_PFO-1
Investigator(s): Matt Regan, Ca	roline Harrington	Section, Township,	, Range:	·
Landform (hillslope, terrace, etc.):	Depression	Local relief (concave, conv	vex, none): Concave	Slope (%): 0 to 1
Subregion (LRR or MLRA): M	ILRA 144A of LRR R	Lat: 42.410410470	04 Long: -72.4669386261	Datum: WGS84
Soil Map Unit Name: Whitman	fine sandy loam, 0 to 3 percent slope	es, extremely stony	NWI classifica	ation:
Are climatic/hydrologic conditions	s on the site typical for this time of ye	ear? Yes <u>√</u> No	(If no, explain in Remar	ks.)
Are Vegetation, Soil,	or Hydrology significantly di		al Circumstances" present?	Yes No
Are Vegetation, Soil,	or Hydrology naturally prob	lematic? (If needed,	explain any answers in Rema	rks.)
Summary of Findings – A	ttach site map showing sampli	ng point locations, trai	nsects, important feature	es, etc.
Hydrophytic Vegetation Present?	? Yes _ ✓ _ No			
Hydric Soil Present?	Yes _ ✓ _ No	Is the Sampled Area withi	in a Wetland?	∕es No
Wetland Hydrology Present?	Yes No	If yes, optional Wetland S		W-PMO-04
			iite ib.	W-F WO-04
·	ocedures here or in a separate report	t)		
Covertype is PFO.				
I				
HYDROLOGY				
Wetland Hydrology Indicators:				
Primary Indicators (minimum of	one is required; check all that apply)		Secondary Indicators (minim	um of two required)
Surface Water (A1)	<u></u> Water-Stained Lea	aves (B9)	Surface Soil Cracks (B6)	
High Water Table (A2)	Aquatic Fauna (B1		✓ Drainage Patterns (B10)	
✓ Saturation (A3)	Marl Deposits (B1		Moss Trim Lines (B16)	(C3)
Water Marks (B1)	Hydrogen Sulfide		Dry-Season Water Table (Crayfish Burrows (C8)	(C2)
Sediment Deposits (B2)	Oxidized Rhizospl	heres on Living Roots (C3)	Saturation Visible on Aeri	ial Imagery (C9)
Drift Deposits (B3)	Processo of Radu	icad Iran (CA)		
Algal Mat or Crust (B4)	Presence of Redu	ction in Tilled Soils (C6)	Stunted or Stressed Plant Geomorphic Position (D2	
Iron Deposits (B5)	Thin Muck Surface		Shallow Aquitard (D3))
Inundation Visible on Aerial I			✓ Microtopographic Relief (D4)
Sparsely Vegetated Concave S		nemana)	FAC-Neutral Test (D5)	, J. 1,
Field Observations:				
Surface Water Present?	Yes No _ _/ Depth	(inches):		
Water Table Present?		(inches):	- Wetland Hydrology Present?	Yes No
Saturation Present?		(inches): 0	- Treatment for orogy i reserve.	
	Tes _ √ _ No Deptil	(inches). 0	-	
(includes capillary fringe)				
Describe Recorded Data (stream	gauge, monitoring well, aerial photo	s, previous inspections), if	available:	
Remarks:				

<u>Tree Stratum</u> (Plot size: <u>30 ft</u>)		Dominant Species?	Indicator Status	Dominance Test worksheet: Number of Dominant Species That	4	(A)
1. Tsuga canadensis	35	Yes	FACU	Are OBL, FACW, or FAC:		
2. Acer rubrum	20	Yes	FAC	Total Number of Dominant Species	8	(B)
3. Betula alleghaniensis	10	No	FAC	Across All Strata:		
k				Percent of Dominant Species That Are OBL, FACW, or FAC:	50	(A/B)
5				Prevalence Index worksheet:		
i				- Total % Cover of:	Multiply	Bv.
·				- OBL species 5	x 1 =	5
	65	= Total Cov	er	FACW species 95	x 2 =	190
apling/Shrub Stratum (Plot size: <u>15 ft</u>)				FAC species 40	x3=	120
. Kalmia latifolia	15	Yes	FACU	FACU species 60	x 4 =	240
Tsuga canadensis	10	Yes	FACU	- UPL species 0	x 5 =	0
. Betula alleghaniensis	10	Yes	FAC	Column Totals 200	-	
i.					(A) _	555 (B)
i.				Prevalence Index = B/A =		
				Hydrophytic Vegetation Indicators:		
				1- Rapid Test for Hydrophytic	Vegetation	
· -	35	= Total Cov	er	2 - Dominance Test is > 50%		
Herb Stratum (Plot size: 5 ft)		-		\checkmark 3 - Prevalence Index is ≤ 3.0 ¹		
. Osmundastrum cinnamomeum	65	Yes	FACW	4 - Morphological Adaptation		supporting
Onoclea sensibilis	30	Yes	FACW	data in Remarks or on a separate s		
3. Not Listed Plant	30	Yes	NI	Problematic Hydrophytic Veg		
I. Symplocarpus foetidus	5	No	OBL	Indicators of hydric soil and wetla	, ,	gy must be
i. Symptocarpus toctidus			ODL	present, unless disturbed or problem	emauc	
5.				Definitions of Vegetation Strata:		
7.				Tree – Woody plants 3 in. (7.6 cm) of breast height (DBH), regardless of		ilameter a
·				Sapling/shrub – Woody plants less		NPU and
).				greater than or equal to 3.28 ft (1 r		Di i and
				Herb – All herbaceous (non-woody		ardless of
0				size, and woody plants less than 3.		gai aicss oi
1				Woody vines – All woody vines gre		28 ft in
2				height.		
	130	_= Total Cov	er	Hydrophytic Vegetation Present?	Voc / N	lo.
Voody Vine Stratum (Plot size: 30 ft)				riyuropriyuc vegetation rresent:	162 1	
				-		
·				-		
				_		
2. 3.						
2				_		

	Matrix		Redox						
(inches)	Color (moist)	<u>%</u>	Color (moist)	<u>%</u>	Type ¹	Loc²	Texture		Remarks
0 - 6	7.5YR 2.5/1	100	10/5 5/5				Mucky P	-	
6 - 8	10Y 4/1	95	10YR 6/6	<u>5</u>	C	M	Sandy Lo	am	
		· —		_					
				-					
				-					
				_					
				-					
				_					
				-					
				-					
¹Type: C = C	Concentration, D =	Depletic	n, RM = Reduced	Mat	rix, MS =	Masked S	and Grains. ² Lo	ocation: PL = F	Pore Lining, M = Matrix.
Hydric Soil I			,		,				or Problematic Hydric Soils ³ :
Histosol			Polyvalue Be	low S	urface (S	8) (LRR R,	MLRA 149B)		ck (A10) (LRR K, L, MLRA 149B)
Histic Ep	oipedon (A2)		Thin Dark Su						airie Redox (A16) (LRR K, L, R)
Black Hi			Loamy Muck			(LRR K, L)			cky Peat or Peat (S3) (LRR K, L, R)
	en Sulfide (A4)		Loamy Gleye						face (S7) (LRR K, L)
	d Layers (A5) d Below Dark Surfa	(111	Depleted Ma					Polyvalu	e Below Surface (S8) (LRR K, L)
	d Below Dark Suria ark Surface (A12)	ace (ATT	Depleted Dark		` '				k Surface (S9) (LRR K, L)
	fucky Mineral (S1)		Redox Depre						nganese Masses (F12) (LRR K, L, R)
	ileyed Matrix (S4)				(,				nt Floodplain Soils (F19) (MLRA 149B)
-	edox (S5)								odic (TA6) (MLRA 144A, 145, 149B)
-	d Matrix (S6)								ent Material (F21)
	rface (S7) (LRR R, M	1LRA 149	9B)					-	llow Dark Surface (TF12) xplain in Remarks)
21	- £ h								
	of hydrophytic veg		and wetland nydi	rolog	y must be	e present, T	uniess disturbe	a or problem	atic.
	ayer (if observed): Type:		None			Hydric S	oil Present?		Yes/ No
	Depth (inches):		None	-		nyuric 3	on Fresent?		res NO
	Deptil (iliches).	_							
Remarks:									

Project/Site: Pratt South	City/County: Shu	tesbury, Franklin	Sampling Date: 2020-Aug-03		
Applicant/Owner: W.D. Cowls	, Inc.	State: MA	Sampling Point: W	/-PMO-04_UPL-1	
Investigator(s): Matt Regan, Ca	aroline Harrington	Section, Township,	Range:		
Landform(hillslope,terrace,etc.):	Flat	Local relief (concave, conv	ex, none): None	Slope (%): 0 to 1	
Subregion(LRRorMLRA): M	LRA 144A of LRR R	Lat: 42.410401082	6 Long: -72.4667832256	Datum: WGS84	
SoilMapUnitName: Hinckley l	oamy sand, 8 to 15 percent slopes		NWI classifica	ation:	
Areclimatic/hydrologiccondition:	sonthesitetypicalforthistimeofyear?	Yes <u></u> ✓ No	(If no, explain in Remarl	ks.)	
Are Vegetation, Soil,	or Hydrology significantly di	sturbed? Are "Norma	al Circumstances" present?	Yes No	
Are Vegetation, Soil,	or Hydrology naturally prob	lematic? (If needed,	explain any answers in Rema	rks.)	
SUMMARY OF FINDINGS – A	Attach site map showing sampli	ng point locations, tran	nsects, important feature	s, etc.	
Hydrophytic Vegetation Present	? Yes No _√				
Hydric Soil Present?	Yes No _ ∠ _	Is the Sampled Area within	n a Wetland?	Yes No/_	
	Yes No _ _	·			
Wetland Hydrology Present?		If yes, optional Wetland Si	ite iD:		
	ocedures here or in a separate report	t)			
Covertype is UPL.					
LIVEROLOGY					
HYDROLOGY					
Wetland Hydrology Indicators:					
Primary Indicators (minimum of	one is required; check all that apply)		Secondary Indicators (minim	um of two required)	
Surface Water (A1)	Water-Stained Le	aves (B9)	Surface Soil Cracks (B6)		
High Water Table (A2)	Aquatic Fauna (B [.]		Drainage Patterns (B10)		
Saturation (A3)	Marl Deposits (B1	5)	Moss Trim Lines (B16)		
Water Marks (B1)	Hydrogen Sulfide	Odor (C1)	Dry-Season Water Table (C2)	
Sediment Deposits (B2)	Oxidized Rhizosp	heres on Living Roots (C3)	Crayfish Burrows (C8)		
			Saturation Visible on Aeri		
Drift Deposits (B3)	Presence of Redu		Stunted or Stressed Plant		
Algal Mat or Crust (B4)		ction in Tilled Soils (C6)	Geomorphic Position (D2)	
Iron Deposits (B5)	Thin Muck Surfac		Shallow Aquitard (D3)	(D.4)	
Inundation Visible on Aerial	· · · · · · · · · · · · · · · · · · ·	Remarks)	Microtopographic Relief (D4)	
Sparsely Vegetated Concave	Surface (B8)		FAC-Neutral Test (D5)		
Field Observations:	Van Na (Danth	(in ab a a).			
Surface Water Present?		(inches):			
Water Table Present?		(inches):	Wetland Hydrology Present?	Yes No _ ∠ _	
Saturation Present?	Yes No _ _/ Depth	(inches):			
(includes capillary fringe)					
Describe Recorded Data (stream	n gauge, monitoring well, aerial photo	s, previous inspections), if a	available:	<u> </u>	
Remarks:					
nemarks.					

VEGETATION -- Use scientific names of plants.

	Absolute	Dominant	Indicator	Dominance Test worksheet:		
<u>Tree Stratum</u> (Plot size: <u>30 ft</u>)		Species?	Status	Number of Dominant Species That		445
1. Tsuga canadensis	50	Yes	FACU	Are OBL, FACW, or FAC:	0	(A)
Quercus rubra		Yes	FACU	Total Number of Dominant Species	6	
3.		103	17100	Across All Strata:		(B)
4.				Percent of Dominant Species That	0	(A /D)
				Are OBL, FACW, or FAC:		(A/B)
5.				Prevalence Index worksheet:		
6.				Total % Cover of:	<u>Multiply</u>	<u>By:</u>
7				OBL species 0	x 1 =	0
	70	= Total Cov	er	FACW species 0	x 2 =	0
Sapling/Shrub Stratum (Plot size: 15 ft)				FAC species 0	x 3 =	0
1. <i>Kalmia latifolia</i>	20	Yes	FACU	FACU species 115	x 4 =	460
2. Tsuga canadensis	15	Yes	FACU	UPL species 0	x 5 =	0
3.				Column Totals 115	(A)	460 (B)
4.					· · · -	460 (b)
5.				Prevalence Index = B/A =	4	
6.				Hydrophytic Vegetation Indicators:		
7.				1- Rapid Test for Hydrophytic \	egetation/	1
/·	 35	= Total Cov	or	2 - Dominance Test is > 50%		
Hart Street van (Diet siese E.S.		_ 10tal C0V	ei	3 - Prevalence Index is $\leq 3.0^{1}$		
Herb Stratum (Plot size:5 ft)	-	V	FACIL	4 - Morphological Adaptations	¹ (Provide	supporting
1. <i>Monotropa uniflora</i>		Yes	FACU	data in Remarks or on a separate sh	neet)	
2. Tsuga canadensis	5	Yes	FACU	Problematic Hydrophytic Vege	tation¹ (Ex	(plain)
3				¹ Indicators of hydric soil and wetlan	d hydrolo	gy must be
4				present, unless disturbed or proble	matic	
5				Definitions of Vegetation Strata:		
6.				Tree – Woody plants 3 in. (7.6 cm) o	r more in	diameter at
7.				breast height (DBH), regardless of h		
8.				Sapling/shrub – Woody plants less t	han 3 in. [DBH and
9.				greater than or equal to 3.28 ft (1 m		
40	-			Herb – All herbaceous (non-woody)		gardless of
-				size, and woody plants less than 3.2	8 ft tall.	
11.				Woody vines – All woody vines grea	ter than 3	.28 ft in
12				height.		
	10	= Total Cov	er	Hydrophytic Vegetation Present?	Voc N	do /
Woody Vine Stratum (Plot size: 30 ft)				Trydrophytic vegetation Fresent:	163 1	NO _ -
1						
2						
3						
4.						
	0	= Total Cov	er			
Barrandar (In dayle abota mark mark and barrandar and a		-				
Remarks: (Include photo numbers here or on a sepa	rate sneet.)					

	cription: (Describe	to the de	-			indicato	r or confirm the ab	sence of indi	icators.)
Depth	Matrix		Redox						
(inches)	Color (moist)	<u>%</u>	Color (moist)	<u>%</u>	Type ¹	Loc ²	Textur		Remarks
0 - 4	7.5YR 2.5/3	100		_			Loamy S		
4 - 8	7.5YR 3/4	100		_			Loamy S	and	
				_			_		
				_					
				_					
				_					
				_					
	-								
				_					
¹Type: C = 0	Concentration, D =	Depletio	n, RM = Reduced	Mat	rix, MS =	Masked	Sand Grains. ² Lo	ocation: PL = F	Pore Lining, M = Matrix.
Hydric Soil	Indicators:							Indicators fo	or Problematic Hydric Soils³:
Histoso			Polyvalue Be	ow S	urface (S	8) (LRR	R, MLRA 149B)	2 cm Mu	ick (A10) (LRR K, L, MLRA 149B)
Histic E _l	oipedon (A2)		Thin Dark Su	rface	(S9) (LRF	R R, MLR	A 149B)		rairie Redox (A16) (LRR K, L, R)
	istic (A3)		Loamy Muck	y Mir	eral (F1)	(LRR K,	L)		icky Peat or Peat (S3) (LRR K, L, R)
-	en Sulfide (A4)		Loamy Gleye						face (S7) (LRR K, L)
	d Layers (A5)		Depleted Ma						e Below Surface (S8) (LRR K, L)
	d Below Dark Surfa	ace (A11)						-	k Surface (S9) (LRR K, L)
	ark Surface (A12) Mucky Mineral (S1)		Depleted Dar)		Iron-Mar	nganese Masses (F12) (LRR K, L, R)
			Redox Depre	SSIOI	IS (F8)			Piedmor	nt Floodplain Soils (F19) (MLRA 149B)
-	Gleyed Matrix (S4)							Mesic Sp	oodic (TA6) (MLRA 144A, 145, 149B)
_	Redox (S5)							Red Pare	ent Material (F21)
	d Matrix (S6)							Very Sha	illow Dark Surface (TF12)
Dark Su	ırface (S7) (LRR R, N	/ILKA 149	9B)					Other (E	xplain in Remarks)
3Indicators	of hydrophytic veg	etation a	and wetland hydr	olog	y must b	e preser	nt, unless disturbed	d or problema	atic.
Restrictive	Layer (if observed):	:							
	Type:		Rock			Hydric	Soil Present?		Yes No <u>_</u> ✓
	Depth (inches):		8						
Remarks:	•								
İ									
l									
Ì									

WETLAND DETERMINATION DATA FORM – Northcentral and Northeast Region

Project/Site: Pratt South	City/County: Shu	tesbury, Franklin	Sampling Date: 2020-Aug-03		
Applicant/Owner: W.D. Cowl	s, Inc.	State: MA	Sampling Point:	W-PMO-05_PFO-1	
Investigator(s): Matt Regan, C	aroline Harrington	Section, Township,	Range:		
Landform(hillslope,terrace,etc.)	: Hillslope	Local relief (concave, conv	ex, none): Concave	Slope (%): 1 to 3	
Subregion(LRRorMLRA): $\underline{\mathbb{N}}$	ILRA 144A of LRR R	Lat: 42.413357799	Long: -72.4713890814	Datum: WGS84	
Soil Map Unit Name: Chatfiel	d-Hollis complex, 8 to 15 percent slope	es, rocky	NWI classific	cation:	
Are climatic/hydrologic condition	ns on the site typical for this time of ye	ear? Yes <u>✓</u> No	(If no, explain in Rema	rks.)	
Are Vegetation, Soil,			al Circumstances" present?	Yes No	
Are Vegetation, Soil,	or Hydrology naturally prob	lematic? (If needed,	explain any answers in Rem	arks.)	
SUMMARY OF FINDINGS - A	Attach site map showing sampli	ng point locations, trai	nsects, important featur	es, etc.	
Hydrophytic Vegetation Presen	t? Yes 🗸 No				
Hydric Soil Present?	Yes _ . ✓_ No	Is the Sampled Area withi	n a Wetland?	Yes/_ No	
Wetland Hydrology Present?	Yes _ ✓ _ No	If yes, optional Wetland S		W-PMO-05	
	<u> </u>		ite ib.	- VV-P1VIO-U3	
·	rocedures here or in a separate report)			
Covertype is PFO.					
HYDROLOGY					
Wetland Hydrology Indicators:	fono is required, shock all that apply		Cocondan Indicators (minin	num of two required)	
•	f one is required; check all that apply)		Secondary Indicators (minin	num of two requirea)	
Surface Water (A1)	Water-Stained Lea		Surface Soil Cracks (B6)		
✓ High Water Table (A2)	Aquatic Fauna (B1		Drainage Patterns (B10) Moss Trim Lines (B16)		
✓ Saturation (A3)	Marl Deposits (B1		Dry-Season Water Table (C2)		
Water Marks (B1)	Hydrogen Sulfide	neres on Living Roots (C3)	- C - (C)		
Sediment Deposits (B2)	Oxidized Kriizospi	ieres on Living Roots (C3)	Saturation Visible on Aerial Imagery (C9)		
Drift Deposits (B3)	Presence of Redu	ced Iron (C4)	Stunted or Stressed Plar		
Algal Mat or Crust (B4)		ction in Tilled Soils (C6)	Geomorphic Position (D.		
Iron Deposits (B5)	Thin Muck Surface		Shallow Aquitard (D3)		
Inundation Visible on Aerial			Microtopographic Relief	(D4)	
Sparsely Vegetated Concave	Surface (B8)		FAC-Neutral Test (D5)		
Field Observations:					
Surface Water Present?	Yes No Depth	(inches):			
Water Table Present?	Yes No Depth	(inches): 0	Wetland Hydrology Present	? Yes No	
Saturation Present?	·	(inches): 0			
(includes capillary fringe)	res <u>y</u> 110	(11101103).	-		
			1.11		
Describe Recorded Data (Stream	m gauge, monitoring well, aerial photo	s, previous inspections), if a	available:		
Remarks:					
1					

VEGETATION -- Use scientific names of plants.

·	Ahsolute	Dominant	Indicator	Dominance Test worksheet:		
<u>Tree Stratum</u> (Plot size: <u>30 ft</u>)		Species?	Status	Number of Dominant Species That		
1. Acer rubrum	25	Yes	FAC	Are OBL, FACW, or FAC:	4	(A)
Betula alleghaniensis	15	Yes	FAC	Total Number of Dominant Species	7	(D)
3. Tsuga canadensis	10	Yes	FACU	Across All Strata:		(B)
4.		103	17100	Percent of Dominant Species That	57.1	(A/B)
5.				Are OBL, FACW, or FAC:		(747 D)
6.				Prevalence Index worksheet:		
7.				Total % Cover of:	Multiply E	<u>Зу:</u>
/·		= Total Cov		OBL species 10	x 1 =	10
Conding/Shaulb Streeture (Diet sine) 45 ft	50	- 10tal COV	er	FACW species 90	x 2 =	180
Sapling/Shrub Stratum (Plot size: 15 ft)	40	V	EA CIA/	FAC species 40	x 3 =	120
1. Lindera benzoin	40	Yes	FACW	FACU species 40	x 4 =	160
2. Hamamelis virginiana	10	Yes	FACU	- UPL species 0	x 5 =	0
3				Column Totals 180	(A)	470 (B)
4				Prevalence Index = B/A =	2.6	
5				Hydrophytic Vegetation Indicators:		
6				1- Rapid Test for Hydrophytic	/egetation	
7				✓ 2 - Dominance Test is >50%	- cgctation	
	50	= Total Cov	er	\checkmark 3 - Prevalence Index is ≤ 3.0 ¹		
Herb Stratum (Plot size:5 ft)				4 - Morphological Adaptations	1 (Provide s	sunnorting
1. Osmundastrum cinnamomeum	50	Yes	FACW	data in Remarks or on a separate sl		apporting
2. <i>Maianthemum canadense</i>	20	Yes	FACU	Problematic Hydrophytic Vege		olain)
3. <i>Carex crinita</i>	10	No	OBL	¹Indicators of hydric soil and wetlar		
4.				present, unless disturbed or proble		,,
5.				Definitions of Vegetation Strata:		
6.				Tree – Woody plants 3 in. (7.6 cm) o	r more in d	liameter at
7.				breast height (DBH), regardless of h		
8.				Sapling/shrub – Woody plants less t	han 3 in. D	BH and
9.				greater than or equal to 3.28 ft (1 m	ı) tall.	
10.				Herb – All herbaceous (non-woody)	plants, reg	ardless of
11.				size, and woody plants less than 3.2	8 ft tall.	
12.				Woody vines – All woody vines grea	ter than 3.2	28 ft in
	80	= Total Cov	er	height.		
Woody Vine Stratum (Plot size:30 ft)				Hydrophytic Vegetation Present?	Yes 🟒 N	0
1.						
2.				•		
3.				-		
4.						
<u> </u>	0	= Total Cov	or			
		- Total Cov	C1			
Remarks: (Include photo numbers here or on a separat	e sheet.)					

(inches) 0 - 10	Matrix				tures	12	T	.	Damada
0 - 10	Color (moist)	<u>%</u>	Color (moist)	<u>%</u>	Type ¹	Loc ²		ture	Remarks
10 - 16	2.5Y 2.5/1	100	10VD F /C	· <u>-</u>				uck	
10 - 16	2.5Y 5/1	95	10YR 5/6	5	C	<u>M</u> _	Sandy C	lay Loam	
				- —					
				-					
				- —					
				-				 -	
				-					
				_				·	
				_					
				_					
				_					
Type: C = C	oncentration, D =	Depletio	n, RM = Reduced	Mat	rix, MS =	Masked Sa	nd Grains. ² Le	ocation: PL = Pore Li	ning, M = Matrix.
Hydric Soil I	ndicators:							Indicators for Prob	olematic Hydric Soils³:
Histosol			Polyvalue Bel					2 cm Muck (A1	0) (LRR K, L, MLRA 149B)
	pipedon (A2)		Thin Dark Su				49B)		edox (A16) (LRR K, L, R)
Black Hi			Loamy Mucky			(LRR K, L)		5 cm Mucky Pe	at or Peat (S3) (LRR K, L, R)
	en Sulfide (A4) d Layers (A5)		Loamy Gleye Depleted Ma					Dark Surface (S	
	d Below Dark Surfa	ce (A11)						-	w Surface (S8) (LRR K, L)
	ark Surface (A12)	` '	Depleted Dar	k Su	rface (F7)			Thin Dark Surf	
Sandy N	lucky Mineral (S1)		Redox Depre	ssior	ns (F8)				se Masses (F12) (LRR K, L, R) dplain Soils (F19) (MLRA 149B)
-	ileyed Matrix (S4)								ГА6) (MLRA 144A, 145, 149B)
-	edox (S5)							Red Parent Ma	
	d Matrix (S6)								ark Surface (TF12)
Dark Su	rface (S7) (LRR R, M	ILRA 149	9B)					Other (Explain	in Remarks)
Indicators (of hydrophytic veg	etation a	and wetland hydr	olog	y must be	present, u	ınless disturbe	d or problematic.	
Restrictive L	ayer (if observed):								
	Type:		None			Hydric So	il Present?	`	/es/_ No
	Depth (inches):								
Remarks:									
Remarks:									
Remarks:									
Remarks:									
Remarks:									
emarks:									
emarks:									
emarks:									
Remarks:									
emarks:									
emarks:									
emarks:									
Remarks:									
Remarks:									
Remarks:									
Remarks:									
emarks:									

WETLAND DETERMINATION DATA FORM – Northcentral and Northeast Region

Project/Site: Pratt South	City/County: Shu	tesbury, Franklin	Sampling Date: 2020-Aug-03			
Applicant/Owner: W.D. Cowls,	Inc.	State: MA	Sampling Point: W	/-PMO-05_UPL-1		
Investigator(s): Matt Regan, Car	roline Harrington	Section, Township,	Range:			
Landform(hillslope,terrace,etc.):	Hillslope	Local relief (concave, conve	ex, none): Convex	Slope (%): 1 to 3		
Subregion (LRR or MLRA): M	LRA 144A of LRR R	Lat: 42.4131767499	D Long: -72.4711473473	Datum: WGS84		
Soil Map Unit Name: Chatfield-	-Hollis complex, 8 to 15 percent slope	es, rocky	NWI classifica	ation:		
•	s on the site typical for this time of ye		(If no, explain in Remark	KS.)		
Are Vegetation, Soil,	or Hydrology significantly di		al Circumstances" present?	Yes _ ∠ No		
Are Vegetation, Soil,	or Hydrology naturally prob	lematic? (If needed, o	explain any answers in Rema	rks.)		
SUMMARY OF FINDINGS – A	ttach site map showing sampli	ng point locations, tran	sects, important feature	es, etc.		
Hydrophytic Vegetation Present?	Yes No					
Hydric Soil Present?	Yes No	Is the Sampled Area within	n a Wetland?	Yes No		
Wetland Hydrology Present?	Yes No _ _ ∠	If yes, optional Wetland Sit	te ID:			
Covertype is UPL.	ocedures here or in a separate report	.)				
Covertype is OPL.						
HYDROLOGY						
HIDKOLOGI						
Wetland Hydrology Indicators:						
Primary Indicators (minimum of o	one is required; check all that apply)		Secondary Indicators (minim	um of two required)		
Surface Water (A1)	Water-Stained Lea	aves (B9)	Surface Soil Cracks (B6)			
High Water Table (A2)	Aquatic Fauna (B1	13)	Drainage Patterns (B10)			
Saturation (A3)	Marl Deposits (B1	5)	Moss Trim Lines (B16)			
Water Marks (B1)	Hydrogen Sulfide		Dry-Season Water Table (C2)			
Sediment Deposits (B2)	Oxidized Rhizospl	heres on Living Roots (C3)	3) Crayfish Burrows (C8) Saturation Visible on Aerial Imagery (C9)			
Drift Deposits (B3)	Dracance of Dodu	and Iran (CA)				
Algal Mat or Crust (B4)	Presence of Redu	ced fron (C4) ction in Tilled Soils (C6)	Stunted or Stressed Plant	• •		
Algai Mat of Crust (B4) Iron Deposits (B5)	Thin Muck Surface		Geomorphic Position (D2) Shallow Aquitard (D3)			
Inundation Visible on Aerial Ir			Microtopographic Relief (D4)			
Sparsely Vegetated Concave S			FAC-Neutral Test (D5)	<i>-</i> .,		
Field Observations:						
Surface Water Present?	Yes No <u></u> Depth	(inches):				
Water Table Present?	,	·	Wetland Hydrology Present?	Yes No ∠		
Saturation Present?		· · · · ——	Wedana Hydrology i resent.	163110		
	res No _ /	(inches):				
(includes capillary fringe)				 -		
Describe Recorded Data (stream	gauge, monitoring well, aerial photo	s, previous inspections), if a	vailable:			
Remarks:						

VEGETATION -- Use scientific names of plants.

	Absoluto	Dominant	Indicator	Dominance Test worksheet:		
<u>Tree Stratum</u> (Plot size: <u>30 ft</u>)		Species?	Status	Number of Dominant Species That		
1. Betula lenta	25	<u> </u>	FACU	Are OBL, FACW, or FAC:	1	(A)
-		Yes		Total Number of Dominant Species		
2. Quercus rubra	20	Yes	FACU	Across All Strata:	6	(B)
3.				Percent of Dominant Species That		
4				Are OBL, FACW, or FAC:	16.7	(A/B)
5				Prevalence Index worksheet:		
6.				Total % Cover of:	Multiply E	By:
7				OBL species 0	x 1 =	0
	45	= Total Cov	er	FACW species 25	x 2 =	50
Sapling/Shrub Stratum (Plot size:15 ft)				FAC species 0	x 3 =	0
1. Hamamelis virginiana	20	Yes	FACU	FACU species 85	x 4 =	340
2. Betula lenta	15	Yes	FACU	UPL species 10	x 5 =	50
3.					_	
4.					(A) _	440 (B)
5.				Prevalence Index = B/A =	3./	
6.				Hydrophytic Vegetation Indicators:		
7.				1- Rapid Test for Hydrophytic	/egetation	
·	35	= Total Cov	ar	2 - Dominance Test is > 50%		
Herb Stratum (Plot size: _ 5 ft)		_ Total Cov	-1	3 - Prevalence Index is $\leq 3.0^{1}$		
1. Osmundastrum cinnamomeum	25	Yes	FACW	4 - Morphological Adaptations		supporting
-			UPL	data in Remarks or on a separate sh		
2. Dennstaedtia punctilobula	10	Yes		Problematic Hydrophytic Vege	tation¹ (Exp	olain)
3. <i>Maianthemum canadense</i>	5	No	FACU	Indicators of hydric soil and wetlan	d hydrolog	y must be
4				present, unless disturbed or proble	matic	
5				Definitions of Vegetation Strata:		
6				Tree – Woody plants 3 in. (7.6 cm) o	r more in d	iameter at
7				breast height (DBH), regardless of h	eight.	
8				Sapling/shrub – Woody plants less t		BH and
9.				greater than or equal to 3.28 ft (1 m) tall.	
10.				Herb – All herbaceous (non-woody)		ardless of
11.				size, and woody plants less than 3.2		
12.				Woody vines – All woody vines grea	ter than 3.2	28 ft in
	40	= Total Cov	er	height.		
Woody Vine Stratum (Plot size:30 ft)		-		Hydrophytic Vegetation Present?	Yes N	0
1.						
2.						
3.						
4						
	0	= Total Cov	er			
Remarks: (Include photo numbers here or on a separat	e sheet.)					
·						

Profile Des Depth	cription: (Describe Matrix	to the d	epth needed to d Redox			indicato	r or confirm the al	bsence of indi	icators.)
(inches)	Color (moist)	%	Color (moist)	%	Type ¹	Loc ²	Textu	re	Remarks
0 - 6	7.5YR 3/2	100					Loamy S		
6 - 10	7.5YR 4/3	100		_			Loamy S		
				_					
				_					
				_					
				_					
				_					
				_					
				_					
				_					
¹Type: C = 0	Concentration, D =	Depletic	n, RM = Reduced	Mati	rix, MS =	Masked	Sand Grains. ² Lo	ocation: PL = F	Pore Lining, M = Matrix.
Hydric Soil		•							or Problematic Hydric Soils ³ :
Histoso			Polyvalue Bel	low S	urface (S	8) (LRR 1	R, MLRA 149B)		ıck (A10) (LRR K, L, MLRA 149B)
Histic E	oipedon (A2)		Thin Dark Su						rairie Redox (A16) (LRR K, L, R)
	istic (A3)		Loamy Mucky			(LRR K, I	_)		icky Peat or Peat (S3) (LRR K, L, R)
, .	en Sulfide (A4)		Loamy Gleye						rface (S7) (LRR K, L)
	d Layers (A5)	(111	Depleted Mar					Polyvalu	e Below Surface (S8) (LRR K, L)
	d Below Dark Surfa ark Surface (A12)	ace (ATT	Depleted Dark					Thin Dar	rk Surface (S9) (LRR K, L)
	Mucky Mineral (S1)		Redox Depre			'		Iron-Mai	nganese Masses (F12) (LRR K, L, R)
-	Gleyed Matrix (S4)		Redox Bepre	55101	15 (1 0)				nt Floodplain Soils (F19) (MLRA 149B)
-	Redox (S5)								oodic (TA6) (MLRA 144A, 145, 149B)
-	d Matrix (S6)								ent Material (F21)
	urface (S7) (LRR R, N	/ILRA 14	9B)					-	allow Dark Surface (TF12)
	,		•					Other (E	xplain in Remarks)
	of hydrophytic veg		and wetland hydr	olog	y must be	e preser	t, unless disturbe	d or problem	atic.
Restrictive	Layer (if observed): _	:				l			
	Type:		Rock	-		Hydric	Soil Present?		Yes No⁄_
	Depth (inches):		10						
Remarks:									



Appendix D: NRCS Soil Report



NRCS

Natural Resources Conservation Service A product of the National Cooperative Soil Survey, a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local participants

Custom Soil Resource Report for Franklin County, Massachusetts



Preface

Soil surveys contain information that affects land use planning in survey areas. They highlight soil limitations that affect various land uses and provide information about the properties of the soils in the survey areas. Soil surveys are designed for many different users, including farmers, ranchers, foresters, agronomists, urban planners, community officials, engineers, developers, builders, and home buyers. Also, conservationists, teachers, students, and specialists in recreation, waste disposal, and pollution control can use the surveys to help them understand, protect, or enhance the environment.

Various land use regulations of Federal, State, and local governments may impose special restrictions on land use or land treatment. Soil surveys identify soil properties that are used in making various land use or land treatment decisions. The information is intended to help the land users identify and reduce the effects of soil limitations on various land uses. The landowner or user is responsible for identifying and complying with existing laws and regulations.

Although soil survey information can be used for general farm, local, and wider area planning, onsite investigation is needed to supplement this information in some cases. Examples include soil quality assessments (http://www.nrcs.usda.gov/wps/portal/nrcs/main/soils/health/) and certain conservation and engineering applications. For more detailed information, contact your local USDA Service Center (https://offices.sc.egov.usda.gov/locator/app?agency=nrcs) or your NRCS State Soil Scientist (http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/contactus/?cid=nrcs142p2 053951).

Great differences in soil properties can occur within short distances. Some soils are seasonally wet or subject to flooding. Some are too unstable to be used as a foundation for buildings or roads. Clayey or wet soils are poorly suited to use as septic tank absorption fields. A high water table makes a soil poorly suited to basements or underground installations.

The National Cooperative Soil Survey is a joint effort of the United States Department of Agriculture and other Federal agencies, State agencies including the Agricultural Experiment Stations, and local agencies. The Natural Resources Conservation Service (NRCS) has leadership for the Federal part of the National Cooperative Soil Survey.

Information about soils is updated periodically. Updated information is available through the NRCS Web Soil Survey, the site for official soil survey information.

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require

alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination, write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

Contents

Preface	2
How Soil Surveys Are Made	
Soil Map	
Soil Map	
Legend	
Map Unit Legend	. 11
Map Unit Descriptions	11
Franklin County, Massachusetts	. 13
71B—Ridgebury fine sandy loam, 3 to 8 percent slopes, extremely	
stony	. 13
73A—Whitman fine sandy loam, 0 to 3 percent slopes, extremely stony	. 14
109C—Chatfield-Hollis complex, 8 to 15 percent slopes, rocky	16
109D—Chatfield-Hollis complex, 15 to 25 percent slopes, rocky	18
245B—Hinckley loamy sand, 3 to 8 percent slopes	
245C—Hinckley loamy sand, 8 to 15 percent slopes	.22
441C—Gloucester sandy loam, 8 to 15 percent slopes, very stony	24
441D—Gloucester sandy loam, 15 to 25 percent slopes, very stony	26
441F—Gloucester sandy loam, 25 to 45 percent slopes, very stony	. 27
References	.29

How Soil Surveys Are Made

Soil surveys are made to provide information about the soils and miscellaneous areas in a specific area. They include a description of the soils and miscellaneous areas and their location on the landscape and tables that show soil properties and limitations affecting various uses. Soil scientists observed the steepness, length, and shape of the slopes; the general pattern of drainage; the kinds of crops and native plants; and the kinds of bedrock. They observed and described many soil profiles. A soil profile is the sequence of natural layers, or horizons, in a soil. The profile extends from the surface down into the unconsolidated material in which the soil formed or from the surface down to bedrock. The unconsolidated material is devoid of roots and other living organisms and has not been changed by other biological activity.

Currently, soils are mapped according to the boundaries of major land resource areas (MLRAs). MLRAs are geographically associated land resource units that share common characteristics related to physiography, geology, climate, water resources, soils, biological resources, and land uses (USDA, 2006). Soil survey areas typically consist of parts of one or more MLRA.

The soils and miscellaneous areas in a survey area occur in an orderly pattern that is related to the geology, landforms, relief, climate, and natural vegetation of the area. Each kind of soil and miscellaneous area is associated with a particular kind of landform or with a segment of the landform. By observing the soils and miscellaneous areas in the survey area and relating their position to specific segments of the landform, a soil scientist develops a concept, or model, of how they were formed. Thus, during mapping, this model enables the soil scientist to predict with a considerable degree of accuracy the kind of soil or miscellaneous area at a specific location on the landscape.

Commonly, individual soils on the landscape merge into one another as their characteristics gradually change. To construct an accurate soil map, however, soil scientists must determine the boundaries between the soils. They can observe only a limited number of soil profiles. Nevertheless, these observations, supplemented by an understanding of the soil-vegetation-landscape relationship, are sufficient to verify predictions of the kinds of soil in an area and to determine the boundaries.

Soil scientists recorded the characteristics of the soil profiles that they studied. They noted soil color, texture, size and shape of soil aggregates, kind and amount of rock fragments, distribution of plant roots, reaction, and other features that enable them to identify soils. After describing the soils in the survey area and determining their properties, the soil scientists assigned the soils to taxonomic classes (units). Taxonomic classes are concepts. Each taxonomic class has a set of soil characteristics with precisely defined limits. The classes are used as a basis for comparison to classify soils systematically. Soil taxonomy, the system of taxonomic classification used in the United States, is based mainly on the kind and character of soil properties and the arrangement of horizons within the profile. After the soil

scientists classified and named the soils in the survey area, they compared the individual soils with similar soils in the same taxonomic class in other areas so that they could confirm data and assemble additional data based on experience and research.

The objective of soil mapping is not to delineate pure map unit components; the objective is to separate the landscape into landforms or landform segments that have similar use and management requirements. Each map unit is defined by a unique combination of soil components and/or miscellaneous areas in predictable proportions. Some components may be highly contrasting to the other components of the map unit. The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The delineation of such landforms and landform segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, onsite investigation is needed to define and locate the soils and miscellaneous areas.

Soil scientists make many field observations in the process of producing a soil map. The frequency of observation is dependent upon several factors, including scale of mapping, intensity of mapping, design of map units, complexity of the landscape, and experience of the soil scientist. Observations are made to test and refine the soil-landscape model and predictions and to verify the classification of the soils at specific locations. Once the soil-landscape model is refined, a significantly smaller number of measurements of individual soil properties are made and recorded. These measurements may include field measurements, such as those for color, depth to bedrock, and texture, and laboratory measurements, such as those for content of sand, silt, clay, salt, and other components. Properties of each soil typically vary from one point to another across the landscape.

Observations for map unit components are aggregated to develop ranges of characteristics for the components. The aggregated values are presented. Direct measurements do not exist for every property presented for every map unit component. Values for some properties are estimated from combinations of other properties.

While a soil survey is in progress, samples of some of the soils in the area generally are collected for laboratory analyses and for engineering tests. Soil scientists interpret the data from these analyses and tests as well as the field-observed characteristics and the soil properties to determine the expected behavior of the soils under different uses. Interpretations for all of the soils are field tested through observation of the soils in different uses and under different levels of management. Some interpretations are modified to fit local conditions, and some new interpretations are developed to meet local needs. Data are assembled from other sources, such as research information, production records, and field experience of specialists. For example, data on crop yields under defined levels of management are assembled from farm records and from field or plot experiments on the same kinds of soil.


Predictions about soil behavior are based not only on soil properties but also on such variables as climate and biological activity. Soil conditions are predictable over long periods of time, but they are not predictable from year to year. For example, soil scientists can predict with a fairly high degree of accuracy that a given soil will have a high water table within certain depths in most years, but they cannot predict that a high water table will always be at a specific level in the soil on a specific date.

After soil scientists located and identified the significant natural bodies of soil in the survey area, they drew the boundaries of these bodies on aerial photographs and

identified each as a specific map unit. Aerial photographs show trees, buildings, fields, roads, and rivers, all of which help in locating boundaries accurately.

Soil Map

The soil map section includes the soil map for the defined area of interest, a list of soil map units on the map and extent of each map unit, and cartographic symbols displayed on the map. Also presented are various metadata about data used to produce the map, and a description of each soil map unit.



MAP LEGEND

Area of Interest (AOI)

Area of Interest (AOI)

Soils

Soil Map Unit Polygons

-

Soil Map Unit Lines

Soil Map Unit Points

Special Point Features

(o)

Blowout

 \boxtimes

Borrow Pit

Ж

Clay Spot

 \Diamond

Closed Depression

Š

Gravel Pit

..

Gravelly Spot

0

Landfill Lava Flow

٨

Marsh or swamp

2

Mine or Quarry

X.

Miscellaneous Water

0

Perennial Water
Rock Outcrop

Saline Spot

• •

Sandy Spot

0

Severely Eroded Spot

Sinkhole

&

Slide or Slip

Ø

Sodic Spot



Spoil Area Stony Spot



Very Stony Spot



Wet Spot Other



Special Line Features

Water Features

_

Streams and Canals

Transportation

ransp

Rails

~

Interstate Highways

~

US Routes

 \sim

Major Roads

~

Local Roads

Background

1

Aerial Photography

MAP INFORMATION

The soil surveys that comprise your AOI were mapped at 1:12.000.

Warning: Soil Map may not be valid at this scale.

Enlargement of maps beyond the scale of mapping can cause misunderstanding of the detail of mapping and accuracy of soil line placement. The maps do not show the small areas of contrasting soils that could have been shown at a more detailed scale.

Please rely on the bar scale on each map sheet for map measurements.

Source of Map: Natural Resources Conservation Service Web Soil Survey URL:

Coordinate System: Web Mercator (EPSG:3857)

Maps from the Web Soil Survey are based on the Web Mercator projection, which preserves direction and shape but distorts distance and area. A projection that preserves area, such as the Albers equal-area conic projection, should be used if more accurate calculations of distance or area are required.

This product is generated from the USDA-NRCS certified data as of the version date(s) listed below.

Soil Survey Area: Franklin County, Massachusetts Survey Area Data: Version 15, Jun 9, 2020

Soil map units are labeled (as space allows) for map scales 1:50.000 or larger.

Date(s) aerial images were photographed: Apr 9, 2011—May 12, 2011

The orthophoto or other base map on which the soil lines were compiled and digitized probably differs from the background imagery displayed on these maps. As a result, some minor shifting of map unit boundaries may be evident.

Map Unit Legend

Map Unit Symbol	Map Unit Name	Acres in AOI	Percent of AOI
71B	Ridgebury fine sandy loam, 3 to 8 percent slopes, extremely stony	6.9	7.5%
73A	Whitman fine sandy loam, 0 to 3 percent slopes, extremely stony	3.2	3.5%
109C	Chatfield-Hollis complex, 8 to 15 percent slopes, rocky	3.5	3.7%
109D	Chatfield-Hollis complex, 15 to 25 percent slopes, rocky	6.5	7.0%
245B	Hinckley loamy sand, 3 to 8 percent slopes	10.1	10.9%
245C	Hinckley loamy sand, 8 to 15 percent slopes	23.0	24.8%
441C	Gloucester sandy loam, 8 to 15 percent slopes, very stony	15.5	16.8%
441D	Gloucester sandy loam, 15 to 25 percent slopes, very stony	17.8	19.2%
441F	Gloucester sandy loam, 25 to 45 percent slopes, very stony	6.1	6.6%
Totals for Area of Interest	'	92.6	100.0%

Map Unit Descriptions

The map units delineated on the detailed soil maps in a soil survey represent the soils or miscellaneous areas in the survey area. The map unit descriptions, along with the maps, can be used to determine the composition and properties of a unit.

A map unit delineation on a soil map represents an area dominated by one or more major kinds of soil or miscellaneous areas. A map unit is identified and named according to the taxonomic classification of the dominant soils. Within a taxonomic class there are precisely defined limits for the properties of the soils. On the landscape, however, the soils are natural phenomena, and they have the characteristic variability of all natural phenomena. Thus, the range of some observed properties may extend beyond the limits defined for a taxonomic class. Areas of soils of a single taxonomic class rarely, if ever, can be mapped without including areas of other taxonomic classes. Consequently, every map unit is made up of the soils or miscellaneous areas for which it is named and some minor components that belong to taxonomic classes other than those of the major soils.

Most minor soils have properties similar to those of the dominant soil or soils in the map unit, and thus they do not affect use and management. These are called noncontrasting, or similar, components. They may or may not be mentioned in a particular map unit description. Other minor components, however, have properties and behavioral characteristics divergent enough to affect use or to require different

management. These are called contrasting, or dissimilar, components. They generally are in small areas and could not be mapped separately because of the scale used. Some small areas of strongly contrasting soils or miscellaneous areas are identified by a special symbol on the maps. If included in the database for a given area, the contrasting minor components are identified in the map unit descriptions along with some characteristics of each. A few areas of minor components may not have been observed, and consequently they are not mentioned in the descriptions, especially where the pattern was so complex that it was impractical to make enough observations to identify all the soils and miscellaneous areas on the landscape.

The presence of minor components in a map unit in no way diminishes the usefulness or accuracy of the data. The objective of mapping is not to delineate pure taxonomic classes but rather to separate the landscape into landforms or landform segments that have similar use and management requirements. The delineation of such segments on the map provides sufficient information for the development of resource plans. If intensive use of small areas is planned, however, onsite investigation is needed to define and locate the soils and miscellaneous areas

An identifying symbol precedes the map unit name in the map unit descriptions. Each description includes general facts about the unit and gives important soil properties and qualities.

Soils that have profiles that are almost alike make up a *soil series*. Except for differences in texture of the surface layer, all the soils of a series have major horizons that are similar in composition, thickness, and arrangement.

Soils of one series can differ in texture of the surface layer, slope, stoniness, salinity, degree of erosion, and other characteristics that affect their use. On the basis of such differences, a soil series is divided into *soil phases*. Most of the areas shown on the detailed soil maps are phases of soil series. The name of a soil phase commonly indicates a feature that affects use or management. For example, Alpha silt loam, 0 to 2 percent slopes, is a phase of the Alpha series.

Some map units are made up of two or more major soils or miscellaneous areas. These map units are complexes, associations, or undifferentiated groups.

A *complex* consists of two or more soils or miscellaneous areas in such an intricate pattern or in such small areas that they cannot be shown separately on the maps. The pattern and proportion of the soils or miscellaneous areas are somewhat similar in all areas. Alpha-Beta complex, 0 to 6 percent slopes, is an example.

An *association* is made up of two or more geographically associated soils or miscellaneous areas that are shown as one unit on the maps. Because of present or anticipated uses of the map units in the survey area, it was not considered practical or necessary to map the soils or miscellaneous areas separately. The pattern and relative proportion of the soils or miscellaneous areas are somewhat similar. Alpha-Beta association, 0 to 2 percent slopes, is an example.

An *undifferentiated group* is made up of two or more soils or miscellaneous areas that could be mapped individually but are mapped as one unit because similar interpretations can be made for use and management. The pattern and proportion of the soils or miscellaneous areas in a mapped area are not uniform. An area can be made up of only one of the major soils or miscellaneous areas, or it can be made up of all of them. Alpha and Beta soils, 0 to 2 percent slopes, is an example.

Some surveys include *miscellaneous areas*. Such areas have little or no soil material and support little or no vegetation. Rock outcrop is an example.

Franklin County, Massachusetts

71B—Ridgebury fine sandy loam, 3 to 8 percent slopes, extremely stony

Map Unit Setting

National map unit symbol: 2w69c

Elevation: 0 to 1,290 feet

Mean annual precipitation: 36 to 71 inches Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 140 to 240 days

Farmland classification: Not prime farmland

Map Unit Composition

Ridgebury, extremely stony, and similar soils: 80 percent

Minor components: 20 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Ridgebury, Extremely Stony

Setting

Landform: Drainageways, hills, ground moraines, depressions, drumlins

Landform position (two-dimensional): Toeslope, footslope Landform position (three-dimensional): Base slope, head slope

Down-slope shape: Concave Across-slope shape: Concave

Parent material: Coarse-loamy lodgment till derived from gneiss, granite, and/or

schist

Typical profile

Oe - 0 to 1 inches: moderately decomposed plant material

A - 1 to 6 inches: fine sandy loam Bw - 6 to 10 inches: sandy loam

Bg - 10 to 19 inches: gravelly sandy loam Cd - 19 to 66 inches: gravelly sandy loam

Properties and qualities

Slope: 3 to 8 percent

Surface area covered with cobbles, stones or boulders: 9.0 percent Depth to restrictive feature: 15 to 35 inches to densic material

Drainage class: Poorly drained

Runoff class: Very high

Capacity of the most limiting layer to transmit water (Ksat): Very low to moderately

low (0.00 to 0.14 in/hr)

Depth to water table: About 0 to 6 inches

Frequency of flooding: None Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 1.9 mmhos/cm) Available water capacity: Low (about 3.0 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 7s

Hydrologic Soil Group: D

Ecological site: F144AY009CT - Wet Till Depressions

Hydric soil rating: Yes

Minor Components

Woodbridge, extremely stony

Percent of map unit: 10 percent

Landform: Drumlins, hills, ground moraines

Landform position (two-dimensional): Footslope, summit, backslope

Landform position (three-dimensional): Crest, side slope

Down-slope shape: Convex Across-slope shape: Linear Hydric soil rating: No

Whitman, extremely stony

Percent of map unit: 8 percent Landform: Depressions Down-slope shape: Concave Across-slope shape: Concave

Hydric soil rating: Yes

Paxton, extremely stony

Percent of map unit: 2 percent

Landform: Hills, ground moraines, drumlins

Landform position (two-dimensional): Summit, shoulder, backslope

Landform position (three-dimensional): Crest, side slope

Down-slope shape: Convex, linear Across-slope shape: Convex, linear

Hydric soil rating: No

73A—Whitman fine sandy loam, 0 to 3 percent slopes, extremely stony

Map Unit Setting

National map unit symbol: 2w695

Elevation: 0 to 1,580 feet

Mean annual precipitation: 36 to 71 inches
Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 140 to 240 days

Farmland classification: Not prime farmland

Map Unit Composition

Whitman, extremely stony, and similar soils: 81 percent

Minor components: 19 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Whitman, Extremely Stony

Setting

Landform: Ground moraines, drumlins, depressions, drainageways, hills

Landform position (two-dimensional): Toeslope Landform position (three-dimensional): Base slope

Down-slope shape: Concave Across-slope shape: Concave

Parent material: Coarse-loamy lodgment till derived from gneiss, granite, and/or schist

Typical profile

Oi - 0 to 1 inches: peat

A - 1 to 10 inches: fine sandy loam

Bg - 10 to 17 inches: gravelly fine sandy loam Cdg - 17 to 61 inches: fine sandy loam

Properties and qualities

Slope: 0 to 3 percent

Surface area covered with cobbles, stones or boulders: 9.0 percent Depth to restrictive feature: 7 to 38 inches to densic material

Drainage class: Very poorly drained

Runoff class: Negligible

Capacity of the most limiting layer to transmit water (Ksat): Very low to moderately

low (0.00 to 0.14 in/hr)

Depth to water table: About 0 to 6 inches

Frequency of flooding: None Frequency of ponding: Frequent

Maximum salinity: Nonsaline (0.0 to 1.9 mmhos/cm) Available water capacity: Low (about 3.0 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 7s

Hydrologic Soil Group: D

Ecological site: F144AY041MA - Very Wet Till Depressions

Hydric soil rating: Yes

Minor Components

Ridgebury, extremely stony

Percent of map unit: 10 percent

Landform: Drainageways, hills, ground moraines, depressions, drumlins

Landform position (two-dimensional): Toeslope, footslope Landform position (three-dimensional): Base slope, head slope

Down-slope shape: Concave Across-slope shape: Concave Hydric soil rating: Yes

Scarboro

Percent of map unit: 5 percent

Landform: Outwash deltas, outwash terraces, depressions, drainageways

Landform position (three-dimensional): Tread

Down-slope shape: Concave Across-slope shape: Concave

Hydric soil rating: Yes

Swansea

Percent of map unit: 3 percent Landform: Swamps, bogs, marshes

Down-slope shape: Concave Across-slope shape: Concave

Hydric soil rating: Yes

Woodbridge, extremely stony

Percent of map unit: 1 percent

Landform: Hills, ground moraines, drumlins

Landform position (two-dimensional): Backslope, footslope, summit

Landform position (three-dimensional): Side slope, crest

Down-slope shape: Concave Across-slope shape: Linear Hydric soil rating: No

109C—Chatfield-Hollis complex, 8 to 15 percent slopes, rocky

Map Unit Setting

National map unit symbol: 2w69l Elevation: 110 to 1.320 feet

Mean annual precipitation: 36 to 71 inches
Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 140 to 240 days

Farmland classification: Not prime farmland

Map Unit Composition

Chatfield, very stony, and similar soils: 55 percent Hollis, very stony, and similar soils: 30 percent

Minor components: 15 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Chatfield, Very Stony

Setting

Landform: Hills, ridges

Landform position (two-dimensional): Summit, backslope, shoulder Landform position (three-dimensional): Crest, side slope, nose slope

Down-slope shape: Convex

Across-slope shape: Linear, convex

Parent material: Coarse-loamy melt-out till derived from granite, gneiss, and/or

schist

Typical profile

Oi - 0 to 1 inches: slightly decomposed plant material

A - 1 to 2 inches: fine sandy loam

Bw - 2 to 30 inches: gravelly fine sandy loam

2R - 30 to 40 inches: bedrock

Properties and qualities

Slope: 8 to 15 percent

Surface area covered with cobbles, stones or boulders: 1.6 percent

Depth to restrictive feature: 20 to 41 inches to lithic bedrock

Drainage class: Well drained

Runoff class: High

Capacity of the most limiting layer to transmit water (Ksat): Very low (0.00 to 0.00

in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None

Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 1.9 mmhos/cm) Available water capacity: Low (about 4.3 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 6s

Hydrologic Soil Group: B

Ecological site: F144AY034CT - Well Drained Till Uplands

Hydric soil rating: No

Description of Hollis, Very Stony

Setting

Landform: Hills, ridges

Landform position (two-dimensional): Backslope, shoulder, summit Landform position (three-dimensional): Crest, side slope, nose slope

Down-slope shape: Convex

Across-slope shape: Linear, convex

Parent material: Coarse-loamy melt-out till derived from granite, gneiss, and/or

schist

Typical profile

Oi - 0 to 2 inches: slightly decomposed plant material

A - 2 to 7 inches: gravelly fine sandy loam Bw - 7 to 16 inches: gravelly fine sandy loam

2R - 16 to 26 inches: bedrock

Properties and qualities

Slope: 8 to 15 percent

Surface area covered with cobbles, stones or boulders: 1.6 percent

Depth to restrictive feature: 8 to 23 inches to lithic bedrock

Drainage class: Somewhat excessively drained

Runoff class: Very high

Capacity of the most limiting layer to transmit water (Ksat): Very low (0.00 to 0.00

in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 1.9 mmhos/cm) Available water capacity: Very low (about 2.7 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 6s

Hydrologic Soil Group: D

Ecological site: F144AY033MA - Shallow Dry Till Uplands

Hydric soil rating: No

Minor Components

Charlton, very stony

Percent of map unit: 8 percent

Landform: Hills, ridges

Landform position (two-dimensional): Backslope Landform position (three-dimensional): Side slope

Down-slope shape: Linear, convex

Across-slope shape: Convex Hydric soil rating: No

Paxton, very stony

Percent of map unit: 4 percent

Landform: Hills, ground moraines, drumlins Landform position (two-dimensional): Backslope Landform position (three-dimensional): Side slope

Down-slope shape: Linear, convex Across-slope shape: Convex, linear

Hydric soil rating: No

Leicester, very stony

Percent of map unit: 2 percent

Landform: Hills, ground moraines, depressions, drainageways Landform position (two-dimensional): Toeslope, footslope Landform position (three-dimensional): Base slope

Down-slope shape: Concave, linear Across-slope shape: Concave Hydric soil rating: Yes

Rock outcrop

Percent of map unit: 1 percent Landform: Ridges, hills Hydric soil rating: No

109D—Chatfield-Hollis complex, 15 to 25 percent slopes, rocky

Map Unit Setting

National map unit symbol: 1hvbd Elevation: 190 to 1,130 feet

Mean annual precipitation: 38 to 52 inches Mean annual air temperature: 35 to 58 degrees F

Frost-free period: 127 to 178 days

Farmland classification: Not prime farmland

Map Unit Composition

Chatfield, rocky, and similar soils: 60 percent Hollis, rocky, and similar soils: 34 percent

Minor components: 6 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Chatfield, Rocky

Setting

Landform: Ground moraines

Landform position (two-dimensional): Backslope Landform position (three-dimensional): Side slope

Down-slope shape: Convex Across-slope shape: Linear

Parent material: Loamy supraglacial till derived from gneiss and/or schist

Typical profile

Oe - 0 to 1 inches: moderately decomposed plant material

A - 1 to 4 inches: fine sandy loam

Bw1 - 4 to 9 inches: gravelly fine sandy loam Bw2 - 9 to 19 inches: cobbly fine sandy loam

BC - 19 to 30 inches: sandy loam

C1 - 30 to 34 inches: gravelly sandy loam C2 - 34 to 37 inches: gravelly sandy loam

R - 37 to 65 inches: bedrock

Properties and qualities

Slope: 15 to 25 percent

Surface area covered with cobbles, stones or boulders: 2.1 percent

Depth to restrictive feature: 20 to 40 inches to lithic bedrock

Drainage class: Well drained Runoff class: Very high

Capacity of the most limiting layer to transmit water (Ksat): Moderately low to high

(0.14 to 6.00 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Available water capacity: Low (about 5.9 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 6s

Hydrologic Soil Group: B

Ecological site: F144AY034CT - Well Drained Till Uplands

Hydric soil rating: No

Description of Hollis, Rocky

Settina

Landform: Upland slopes

Landform position (two-dimensional): Backslope Landform position (three-dimensional): Side slope

Down-slope shape: Linear Across-slope shape: Convex

Parent material: Loamy supraglacial till derived from gneiss and/or schist

Typical profile

Oi - 0 to 1 inches: slightly decomposed plant material Oa - 1 to 3 inches: highly decomposed plant material

A - 3 to 4 inches: fine sandy loam

Bw - 4 to 15 inches: cobbly fine sandy loam

R - 15 to 65 inches: bedrock

Properties and qualities

Slope: 15 to 25 percent

Surface area covered with cobbles, stones or boulders: 2.1 percent

Depth to restrictive feature: 10 to 20 inches to lithic bedrock

Drainage class: Somewhat excessively drained

Runoff class: Very high

Capacity of the most limiting layer to transmit water (Ksat): Moderately low to

moderately high (0.14 to 0.60 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Available water capacity: Very low (about 2.8 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 6s

Hydrologic Soil Group: D

Ecological site: F144AY033MA - Shallow Dry Till Uplands

Hydric soil rating: No

Minor Components

Charlton, rocky

Percent of map unit: 2 percent

Landform: Valley sides on moraines, toes on moraines Landform position (two-dimensional): Backslope Landform position (three-dimensional): Side slope

Down-slope shape: Linear Across-slope shape: Linear Hydric soil rating: No

Montauk, very stony

Percent of map unit: 1 percent

Landform: Drumlins, ground moraines

Landform position (two-dimensional): Backslope Landform position (three-dimensional): Side slope

Down-slope shape: Convex Across-slope shape: Linear Hydric soil rating: No

Paxton, very stony

Percent of map unit: 1 percent

Landform: Ground moraines, drumlins

Landform position (two-dimensional): Backslope Landform position (three-dimensional): Side slope

Down-slope shape: Convex Across-slope shape: Linear Hydric soil rating: No

Canton, rocky

Percent of map unit: 1 percent

Landform: Ground moraines, valley sides, hillslopes Landform position (two-dimensional): Backslope Landform position (three-dimensional): Side slope

Down-slope shape: Linear Across-slope shape: Convex Hydric soil rating: No

Rock outcrop

Percent of map unit: 1 percent Hydric soil rating: Unranked

245B—Hinckley loamy sand, 3 to 8 percent slopes

Map Unit Setting

National map unit symbol: 2svm8

Elevation: 0 to 1.430 feet

Mean annual precipitation: 36 to 53 inches Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 140 to 250 days

Farmland classification: Farmland of statewide importance

Map Unit Composition

Hinckley and similar soils: 85 percent Minor components: 15 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Hinckley

Setting

Landform: Kames, outwash terraces, outwash deltas, outwash plains, eskers, moraines, kame terraces

Landform position (two-dimensional): Summit, backslope, footslope, shoulder Landform position (three-dimensional): Nose slope, side slope, base slope, crest, riser, tread

Down-slope shape: Linear, convex, concave Across-slope shape: Convex, linear, concave

Parent material: Sandy and gravelly glaciofluvial deposits derived from gneiss and/or granite and/or schist

Typical profile

Oe - 0 to 1 inches: moderately decomposed plant material

A - 1 to 8 inches: loamy sand

Bw1 - 8 to 11 inches: gravelly loamy sand Bw2 - 11 to 16 inches: gravelly loamy sand BC - 16 to 19 inches: very gravelly loamy sand

C - 19 to 65 inches: very gravelly sand

Properties and qualities

Slope: 3 to 8 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Excessively drained

Runoff class: Very low

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to very

high (1.42 to 99.90 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 1.9 mmhos/cm) Available water capacity: Very low (about 3.0 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 3s

Hydrologic Soil Group: A

Ecological site: F144AY022MA - Dry Outwash

Hydric soil rating: No

Minor Components

Windsor

Percent of map unit: 8 percent

Landform: Eskers, moraines, outwash terraces, outwash deltas, kame terraces, outwash plains, kames

Landform position (two-dimensional): Summit, shoulder, backslope, footslope Landform position (three-dimensional): Nose slope, side slope, base slope, crest, riser, tread

Down-slope shape: Linear, convex, concave Across-slope shape: Convex, linear, concave

Hydric soil rating: No

Sudbury

Percent of map unit: 5 percent

Landform: Outwash deltas, kame terraces, outwash plains, moraines, outwash terraces

Landform position (two-dimensional): Backslope, footslope

Landform position (three-dimensional): Side slope, base slope, head slope, tread

Down-slope shape: Concave, linear Across-slope shape: Linear, concave

Hydric soil rating: No

Agawam

Percent of map unit: 2 percent

Landform: Outwash terraces, outwash deltas, kame terraces, outwash plains, kames, eskers, moraines

Landform position (two-dimensional): Summit, shoulder, backslope, footslope Landform position (three-dimensional): Nose slope, side slope, base slope, crest, riser, tread

Down-slope shape: Linear, convex, concave Across-slope shape: Convex, linear, concave

Hydric soil rating: No

245C—Hinckley loamy sand, 8 to 15 percent slopes

Map Unit Setting

National map unit symbol: 2svm9

Elevation: 0 to 1,480 feet

Mean annual precipitation: 36 to 71 inches
Mean annual air temperature: 39 to 55 degrees F

Frost-free period: 140 to 240 days

Farmland classification: Farmland of statewide importance

Map Unit Composition

Hinckley and similar soils: 85 percent Minor components: 15 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Hinckley

Setting

Landform: Kames, eskers, moraines, outwash terraces, outwash deltas, kame terraces, outwash plains

Landform position (two-dimensional): Shoulder, toeslope, footslope, backslope Landform position (three-dimensional): Nose slope, side slope, crest, head slope, riser

Down-slope shape: Linear, concave, convex Across-slope shape: Convex, linear, concave

Parent material: Sandy and gravelly glaciofluvial deposits derived from gneiss and/or granite and/or schist

Typical profile

Oe - 0 to 1 inches: moderately decomposed plant material

A - 1 to 8 inches: loamy sand

Bw1 - 8 to 11 inches: gravelly loamy sand Bw2 - 11 to 16 inches: gravelly loamy sand BC - 16 to 19 inches: very gravelly loamy sand

C - 19 to 65 inches: very gravelly sand

Properties and qualities

Slope: 8 to 15 percent

Depth to restrictive feature: More than 80 inches

Drainage class: Excessively drained

Runoff class: Very low

Capacity of the most limiting layer to transmit water (Ksat): Moderately high to very

high (1.42 to 99.90 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Maximum salinity: Nonsaline (0.0 to 1.9 mmhos/cm) Available water capacity: Low (about 3.1 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 4e

Hydrologic Soil Group: A

Ecological site: F144AY022MA - Dry Outwash

Hydric soil rating: No

Minor Components

Merrimac

Percent of map unit: 5 percent

Landform: Moraines, outwash terraces, outwash plains, kames, eskers

Landform position (two-dimensional): Shoulder, backslope, footslope, toeslope

Landform position (three-dimensional): Side slope, crest, head slope, nose slope,

riser

Down-slope shape: Convex Across-slope shape: Convex

Hydric soil rating: No

Windsor

Percent of map unit: 5 percent

Landform: Outwash terraces, outwash deltas, kames, eskers, moraines, kame

terraces, outwash plains

Landform position (two-dimensional): Shoulder, backslope, footslope, toeslope Landform position (three-dimensional): Nose slope, side slope, crest, head slope,

riser

Down-slope shape: Linear, concave, convex Across-slope shape: Convex, linear, concave

Hydric soil rating: No

Sudbury

Percent of map unit: 5 percent

Landform: Kame terraces, outwash plains, moraines, outwash deltas, outwash

terraces

Landform position (two-dimensional): Backslope, footslope Landform position (three-dimensional): Base slope, tread

Down-slope shape: Concave, linear Across-slope shape: Linear, concave

Hydric soil rating: No

441C—Gloucester sandy loam, 8 to 15 percent slopes, very stony

Map Unit Setting

National map unit symbol: 9c7p Elevation: 380 to 1,040 feet

Mean annual precipitation: 38 to 50 inches
Mean annual air temperature: 35 to 58 degrees F

Frost-free period: 127 to 178 days

Farmland classification: Farmland of statewide importance

Map Unit Composition

Gloucester, very stony, and similar soils: 87 percent

Minor components: 13 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Gloucester, Very Stony

Settina

Landform: Moraines, upland slopes

Landform position (two-dimensional): Backslope Landform position (three-dimensional): Side slope

Down-slope shape: Linear Across-slope shape: Convex

Parent material: Sandy and gravelly supraglacial till derived from gneiss

Typical profile

Oa - 0 to 2 inches: highly decomposed plant material

A - 2 to 6 inches: sandy loam

Bw1 - 6 to 15 inches: gravelly sandy loam

Bw2 - 15 to 29 inches: very gravelly loamy coarse sand C - 29 to 65 inches: very gravelly loamy coarse sand

Properties and qualities

Slope: 8 to 15 percent

Surface area covered with cobbles, stones or boulders: 2.1 percent

Depth to restrictive feature: More than 80 inches Drainage class: Somewhat excessively drained

Runoff class: Low

Capacity of the most limiting layer to transmit water (Ksat): Moderately high (0.20

to 0.60 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Available water capacity: Very low (about 2.4 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 6s

Hydrologic Soil Group: C

Ecological site: F144AY032NH - Dry Till Uplands

Hydric soil rating: No

Minor Components

Canton, very stony

Percent of map unit: 5 percent

Landform: Ground moraines, valley sides, hillslopes Landform position (two-dimensional): Backslope Landform position (three-dimensional): Side slope

Down-slope shape: Linear Across-slope shape: Convex Hydric soil rating: No

Montauk, very stony

Percent of map unit: 5 percent

Landform: Drumlins, ground moraines

Landform position (two-dimensional): Summit, backslope

Landform position (three-dimensional): Side slope

Down-slope shape: Convex Across-slope shape: Linear Hydric soil rating: No

Newfields, very stony

Percent of map unit: 2 percent

Landform: Depressions on ground moraines, swales on ground moraines

Landform position (two-dimensional): Backslope Landform position (three-dimensional): Side slope

Down-slope shape: Linear Across-slope shape: Concave

Hydric soil rating: No

Ridgebury, very stony

Percent of map unit: 1 percent

Landform: Depressions on drumlins, depressions on ground moraines

Landform position (two-dimensional): Footslope

Landform position (three-dimensional): Side slope

Down-slope shape: Concave Across-slope shape: Linear, convex

Hydric soil rating: Yes

441D—Gloucester sandy loam, 15 to 25 percent slopes, very stony

Map Unit Setting

National map unit symbol: 9c7q Elevation: 360 to 1,040 feet

Mean annual precipitation: 38 to 50 inches
Mean annual air temperature: 35 to 58 degrees F

Frost-free period: 127 to 178 days

Farmland classification: Not prime farmland

Map Unit Composition

Gloucester, very stony, and similar soils: 90 percent

Minor components: 10 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Gloucester, Very Stony

Setting

Landform: Moraines, upland slopes

Landform position (two-dimensional): Backslope Landform position (three-dimensional): Side slope

Down-slope shape: Linear Across-slope shape: Convex

Parent material: Sandy and gravelly supraglacial till derived from gneiss

Typical profile

Oa - 0 to 2 inches: highly decomposed plant material

A - 2 to 6 inches: sandy loam

Bw1 - 6 to 15 inches: gravelly sandy loam

Bw2 - 15 to 29 inches: very gravelly loamy coarse sand C - 29 to 65 inches: very gravelly loamy coarse sand

Properties and qualities

Slope: 15 to 25 percent

Surface area covered with cobbles, stones or boulders: 2.1 percent

Depth to restrictive feature: More than 80 inches Drainage class: Somewhat excessively drained

Runoff class: Medium

Capacity of the most limiting layer to transmit water (Ksat): Moderately high (0.20

to 0.60 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Available water capacity: Very low (about 2.4 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 6s

Hydrologic Soil Group: C

Ecological site: F144AY032NH - Dry Till Uplands

Hydric soil rating: No

Minor Components

Canton, very stony

Percent of map unit: 5 percent

Landform: Valley sides, hillslopes, ground moraines Landform position (two-dimensional): Backslope Landform position (three-dimensional): Side slope

Down-slope shape: Linear Across-slope shape: Convex

Hydric soil rating: No

Montauk, very stony

Percent of map unit: 5 percent

Landform: Drumlins, ground moraines

Landform position (two-dimensional): Backslope Landform position (three-dimensional): Side slope

Down-slope shape: Convex Across-slope shape: Linear Hydric soil rating: No

441F—Gloucester sandy loam, 25 to 45 percent slopes, very stony

Map Unit Setting

National map unit symbol: 9cd4 Elevation: 370 to 1,010 feet

Mean annual precipitation: 38 to 50 inches
Mean annual air temperature: 35 to 58 degrees F

Frost-free period: 127 to 178 days

Farmland classification: Not prime farmland

Map Unit Composition

Gloucester, very stony, and similar soils: 90 percent

Minor components: 10 percent

Estimates are based on observations, descriptions, and transects of the mapunit.

Description of Gloucester, Very Stony

Setting

Landform: Moraines, upland slopes

Landform position (two-dimensional): Backslope Landform position (three-dimensional): Side slope

Down-slope shape: Linear Across-slope shape: Convex

Custom Soil Resource Report

Parent material: Sandy and gravelly supraglacial till derived from gneiss

Typical profile

Oa - 0 to 2 inches: highly decomposed plant material

A - 2 to 6 inches: sandy loam

Bw1 - 6 to 15 inches: gravelly sandy loam

Bw2 - 15 to 29 inches: very gravelly loamy coarse sand C - 29 to 65 inches: very gravelly loamy coarse sand

Properties and qualities

Slope: 25 to 45 percent

Surface area covered with cobbles, stones or boulders: 2.1 percent

Depth to restrictive feature: More than 80 inches Drainage class: Somewhat excessively drained

Runoff class: Medium

Capacity of the most limiting layer to transmit water (Ksat): Moderately high (0.20

to 0.60 in/hr)

Depth to water table: More than 80 inches

Frequency of flooding: None Frequency of ponding: None

Available water capacity: Very low (about 2.4 inches)

Interpretive groups

Land capability classification (irrigated): None specified

Land capability classification (nonirrigated): 7s

Hydrologic Soil Group: C

Ecological site: F144AY032NH - Dry Till Uplands

Hydric soil rating: No

Minor Components

Canton, very stony

Percent of map unit: 5 percent

Landform: Ground moraines, valley sides, hillslopes Landform position (two-dimensional): Backslope Landform position (three-dimensional): Side slope

Down-slope shape: Linear Across-slope shape: Convex

Hydric soil rating: No

Montauk, very stony

Percent of map unit: 5 percent

Landform: Drumlins, ground moraines

Landform position (two-dimensional): Backslope Landform position (three-dimensional): Side slope

Down-slope shape: Convex Across-slope shape: Linear Hydric soil rating: No

References

American Association of State Highway and Transportation Officials (AASHTO). 2004. Standard specifications for transportation materials and methods of sampling and testing. 24th edition.

American Society for Testing and Materials (ASTM). 2005. Standard classification of soils for engineering purposes. ASTM Standard D2487-00.

Cowardin, L.M., V. Carter, F.C. Golet, and E.T. LaRoe. 1979. Classification of wetlands and deep-water habitats of the United States. U.S. Fish and Wildlife Service FWS/OBS-79/31.

Federal Register. July 13, 1994. Changes in hydric soils of the United States.

Federal Register. September 18, 2002. Hydric soils of the United States.

Hurt, G.W., and L.M. Vasilas, editors. Version 6.0, 2006. Field indicators of hydric soils in the United States.

National Research Council. 1995. Wetlands: Characteristics and boundaries.

Soil Survey Division Staff. 1993. Soil survey manual. Soil Conservation Service. U.S. Department of Agriculture Handbook 18. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2 054262

Soil Survey Staff. 1999. Soil taxonomy: A basic system of soil classification for making and interpreting soil surveys. 2nd edition. Natural Resources Conservation Service, U.S. Department of Agriculture Handbook 436. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2 053577

Soil Survey Staff. 2010. Keys to soil taxonomy. 11th edition. U.S. Department of Agriculture, Natural Resources Conservation Service. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2 053580

Tiner, R.W., Jr. 1985. Wetlands of Delaware. U.S. Fish and Wildlife Service and Delaware Department of Natural Resources and Environmental Control, Wetlands Section.

United States Army Corps of Engineers, Environmental Laboratory. 1987. Corps of Engineers wetlands delineation manual. Waterways Experiment Station Technical Report Y-87-1.

United States Department of Agriculture, Natural Resources Conservation Service. National forestry manual. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/home/?cid=nrcs142p2 053374

United States Department of Agriculture, Natural Resources Conservation Service. National range and pasture handbook. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/landuse/rangepasture/?cid=stelprdb1043084

Custom Soil Resource Report

United States Department of Agriculture, Natural Resources Conservation Service. National soil survey handbook, title 430-VI. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/scientists/?cid=nrcs142p2_054242

United States Department of Agriculture, Natural Resources Conservation Service. 2006. Land resource regions and major land resource areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296. http://www.nrcs.usda.gov/wps/portal/nrcs/detail/national/soils/?cid=nrcs142p2_053624

United States Department of Agriculture, Soil Conservation Service. 1961. Land capability classification. U.S. Department of Agriculture Handbook 210. http://www.nrcs.usda.gov/Internet/FSE_DOCUMENTS/nrcs142p2_052290.pdf



Appendix E: USGS StreamStats Report

S-MJR-1 StreamStats Report

Region ID: MA

Workspace ID: MA20200828023112111000

Clicked Point (Latitude, Longitude): 42.41106, -72.45968

Time: 2020-08-27 22:31:28 -0400



Basin Characteristics					
Parameter Code	Parameter Description	Value	Unit		
DRNAREA	Area that drains to a point on a stream	0.0906	square miles		
ELEV	Mean Basin Elevation	970	feet		
LC06STOR	Percentage of water bodies and wetlands determined from the NLCD 2006	0	percent		
BSLDEM250	Mean basin slope computed from 1:250K DEM	4.012	percent		
DRFTPERSTR	Area of stratified drift per unit of stream length	-100000	square mile per mile		

Parameter Code	Parameter Description	Value	Unit
MAREGION	Region of Massachusetts 0 for Eastern 1 for Western	1	dimensionless
BSLDEM10M	Mean basin slope computed from 10 m DEM	11.842	percent
PCTSNDGRV	Percentage of land surface underlain by sand and gravel deposits	0	percent
FOREST	Percentage of area covered by forest	99.23	percent
ACRSDFT	Area underlain by stratified drift	0	square miles
CENTROIDX	Basin centroid horizontal (x) location in state plane coordinates	121484	meters
CENTROIDY	Basin centroid vertical (y) location in state plane units	907158.8	meters
CRSDFT	Percentage of area of coarse-grained stratified drift	0	percent
LAKEAREA	Percentage of Lakes and Ponds	0	percent
LC11DEV	Percentage of developed (urban) land from NLCD 2011 classes 21-24	0	percent
LC11IMP	Average percentage of impervious area determined from NLCD 2011 impervious dataset	0	percent
MAXTEMPC	Mean annual maximum air temperature over basin area, in degrees Centigrade	13.4	feet per mi
OUTLETX	Basin outlet horizontal (x) location in state plane coordinates	121005	feet
OUTLETY	Basin outlet vertical (y) location in state plane coordinates	907175	feet
PRECPRIS00	Basin average mean annual precipitation for 1971 to 2000 from PRISM	48.4	inches
STRMTOT	total length of all mapped streams (1:24,000-scale) in the basin	0	miles
WETLAND	Percentage of Wetlands	0	percent

Peak-Flow Statistics Parameters[Peak Statewide 2016 5156]					
Parameter Code	Parameter Name	Value	Units	Min Limit	Max Limit

Parameter Code	Parameter Name	Value	Units	Min Limit	Max Limit
DRNAREA	Drainage Area	0.0906	square miles	0.16	512
ELEV	Mean Basin Elevation	970	feet	80.6	1948
LC06STOR	Percent Storage from NLCD2006	0	percent	0	32.3

Peak-Flow Statistics Disclaimers[Peak Statewide 2016 5156]

One or more of the parameters is outside the suggested range. Estimates were extrapolated with unknown errors

Peak-Flow Statistics Flow Report[Peak Statewide 2016 5156]

Statistic	Value	Unit
2 Year Peak Flood	9.33	ft^3/s
5 Year Peak Flood	16.5	ft^3/s
10 Year Peak Flood	22.7	ft^3/s
25 Year Peak Flood	32	ft^3/s
50 Year Peak Flood	40.1	ft^3/s
100 Year Peak Flood	48.9	ft^3/s
200 Year Peak Flood	58.8	ft^3/s
500 Year Peak Flood	73.5	ft^3/s

Peak-Flow Statistics Citations

Zarriello, P.J.,2017, Magnitude of flood flows at selected annual exceedance probabilities for streams in Massachusetts: U.S. Geological Survey Scientific Investigations Report 2016-5156, 99 p. (https://dx.doi.org/10.3133/sir20165156)

Low-Flow Statistics Parameters [Statewide Low Flow WRIR00 4135]

Parameter Code	Parameter Name	Value	Units	Min Limit	Max Limit
DRNAREA	Drainage Area	0.0906	square miles	1.61	149
BSLDEM250	Mean Basin Slope from 250K DEM	4.012	percent	0.32	24.6

Parameter Code	Parameter Name	Value	Units	Min Limit	Max Limit
DRFTPERSTR	Stratified Drift per Stream Length	-100000	square mile per mile	0	1.29
MAREGION	Massachusetts Region	1	dimensionless	0	1

Low-Flow Statistics Flow Report[Statewide Low Flow WRIR00 4135]

Statistic Value Unit

Low-Flow Statistics Citations

Sauer, Vernon B.; Thomas, W. O., Jr.; Stricker, V. A.; Wilson, K. V.,1983, Flood characteristics of urban watersheds in the United States: U.S. Geological Survey Water-Supply Paper 2207, 63 p. (http://pubs.er.usgs.gov/publication/wsp2207)
()

Anderson, B.T.,2020, Magnitude and frequency of floods in Alabama, 2015: U.S. Geological Survey Scientific Investigations Report 2020–5032, 148 p.

(https://doi.org/10.3133/sir20205032)

Hedgecock, T.S.,2004, Magnitude and Frequency of Floods on Small Rural Streams in Alabama: U. S. Geological Survey Scientific Investigations Report 2004-5135, 10 p. (http://pubs.usgs.gov/sir/2004/5135/)

Hedgecock, T.S.,2010, Magnitude and Frequency of Floods for Urban Streams in Alabama, 2007: U.S Geological Survey Scientific Investigations Report 2010-5012, 17p. (https://pubs.usgs.gov/sir/2010/5012/)

Wiley, J.B., and Curran, J.H.,2003, Estimating annual high-flow statistics and monthly and seasonal low-flow statistics for ungaged sites on streams in Alaska and conterminous basins in Canada: U.S. Geological Survey Water-Resources Investigations Report 03-4114, 61 p. (http://water.usgs.gov/pubs/wri/wri034114/pdf/wri034114_v1.10.pdf)
Brabets, Timothy P.,1996, Evaluation of the streamflow-gaging network of Alaska in providing regional streamflow information: U.S. Geological Survey Water-Resources Investigations Report 96-4001, 98 p. (http://pubs.er.usgs.gov/usgspubs/wri/wri964001)
Curran, J.H., Barth, N.A., Veilleux, A.G., and Ourso, R.T.,2016, Estimating Flood Magnitude and Frequency at Gaged and Ungaged Sites on Streams in Alaska and Conterminous Basins in Canada, Based on Data through Water Year 2012: U.S. Geological Survey Scientific Investigations Report 2016-5024, 47 p.

(http://dx.doi.org/10.3133/sir20165024http://dx.doi.org/10.3133/sir20165024)
Southard, R.E.,2010, Estimation of the Magnituude and Frequency of Floods in Urban
Basins in Missouri: U.S. Geological Survey Scientific Investigations Report 2010-5073, 27
p. (http://pubs.usgs.gov/sir/2010/5073/)

Waltemeyer, S.D., Analysis of the Magnitude and Frequency of Peak Discharges for the Navajo Nation in Arizona, Utah, Colorado, and New Mexico: U. S. Geological Survey Scientific Investigations Report2006-5306, 42 p. (http://pubs.usgs.gov/sir/2006/5306/) Paretti, N.V., Kennedy, J.R., Turney, L.A., and Veilleux, A.G.,2014, Methods for estimating magnitude and frequency of floods in Arizona, developed with unregulated and rural peakflow data through water year 2010: U.S. Geological Survey Scientific Investigations Report

2014-5211, 61 p., http://dx.doi.org/10.3133/sir20145211.

(http://pubs.usgs.gov/sir/2014/5211/)

Kennedy, J.R., Paretti, N.V., and Veilleux, A.G.,2014, Methods for estimating magnitude and frequency of 1-, 3-, 7-, 15-, and 30-day flood-duration flows in Arizona: U.S. Geological Survey Scientific Investigations Report 2014–5109, 35 p.

(http://pubs.usgs.gov/sir/2014/5109/)

Report 2009-5136, 32 p.

Funkhouser, J.E., Eng, Ken, and Moix, M.W.,2008, Low-Flow Characteristics and Regionalization of Low Flow Characteristics for Selected Streams in Arkansas: U. S. Geological Survey Scientific Investigations Report 2008-5065, 161 p.

(http://pubs.usgs.gov/sir/2008/5065/pdf/SIR2008-5065.pdf)

Breaker, B.K.,2015, Dry season mean monthly flow and harmonic mean flow regression equations for selected ungaged basins in Arkansas: U.S. Geological Survey Scientific Investigations Report 2015–5031, 25 p. (http://pubs.usgs.gov/sir/2015/5031/) Wagner, D.M., Krieger, J.D., and Veilleux, A.G.,2016, Methods for estimating annual

exceedance probability discharges for streams in Arkansas, based on data through water year 2013: U.S. Geological Survey Scientific Investigations Report 2016–5081, 136 p. (http://dx.doi.org/10.3133/sir20165081)

Thomas, B.E, Hjalmarson, H.W., and Waltemeyer, S.D.,1997, Methods for Estimating Magnitude and Frequency of Floods in the Southwestern United States: U.S. Water-Supply Paper 2433, 196 p. (http://pubs.er.usgs.gov/publication/wsp2433)

Gotvald, A.J., Barth, N.A., Veilleux, A.G., and Parrett, Charles, 2012, Methods for determining magnitude and frequency of floods in California, based on data through water year 2006: U.S. Geological Survey Scientific Investigations Report 2012–5113, 38 p., 1 pl. (http://pubs.usgs.gov/sir/2012/5113/)

Sanocki, C.A., Williams-Sether, T., Steeves, P.A., and Christensen, V.G.,2019, Techniques for Estimating the Magnitude and Frequency of Peak Flows on Small Streams in the Binational U.S. and Canadian Lake of the Woods-Rainy River Basin Upstream from Kenora, Ontario, Canada, Based on Data through Water Year 2013: U.S. Geological Survey Scientific Investigations Report 2019–5012, 17 p. (https://doi.org/10.3133/sir20195012)
Capesius, J.P., and Stephens, V. C.,2009, Regional Regression Equations for Estimation of Natural Streamflow Statistics in Colorado: U. S. Geological Survey Scientific Investigations

(http://pubs.usgs.gov/sir/2009/5136/http://pubs.usgs.gov/sir/2009/5136/)

Kohn, M.S., Stevens, M.R., Harden, T.M., Godaire, J.E., Klinger, R.E., and Mommandi, A.,2016, Paleoflood investigations to improve peak-streamflow regional-regression equations for natural streamflow in eastern Colorado, 2015: U.S. Geological Survey Scientific Investigations Report 2016–5099, 58 p. (http://dx.doi.org/10.3133/sir20165099) Ahearn, E.A.,2004, Regression Equations for Estimating Flood Flows for the 2-, 10-, 25-, 50-, 100-, and 500-Year Recurrence Intervals in Connecticut: U.S. Geological Survey SRI 2004-5160, 62 p. (http://water.usgs.gov/pubs/sir/2004/5160/)

Ahearn, E.A.,2010, Regional regression equations to estimate flow-duration statistics in Connecticut: U. S. Geological Survey Scientific Investigations Report 2010-5052, 45 p. (http://pubs.usgs.gov/sir/2010/5052/)

Ries, K.G., III, and Dillow, J.J.A.,2006, Magnitude and frequency of floods in Delaware: Scientific Investigations Report 2006-5146, 59 p. (http://pubs.usgs.gov/sir/2006/5146/) Carpenter, D.H., and Hayes, D.C.,1996, Low-flow characteristics of streams in Maryland and Delaware: U.S. Geological Survey Water-Resources Investigations Report 94-4020, 113 p., 10 plates (http://pubs.er.usgs.gov/usgspubs/wri/wri944020)

Franklin, M.A. and Losey, G.T.,1984, Magnitude and Frequency of Floods from Urban Streams in Leon County, Florida: U.S. Geological Survey Water-Resources Investigations Report 84-4004, 37 p. (http://pubs.er.usgs.gov/publication/wri844004)

Lopez, M.A. and Woodham, W. M.,1983, Magnitude and frequency of flooding on small urban watersheds in the Tampa Bay area, west-central Florida: U.S. Geological Survey Water-Resources Investigations Report 82-42, 52 p.

(https://pubs.er.usgs.gov/publication/wri8242)

(https://doi.org/10.3133/sir20175001)

Rumenik, R. P.; Grubbs, J. W.,1996, Methods for estimating low-flow characteristics of ungaged streams in selected areas, northern Florida: U.S. Geological Survey Water-Resources Investigations Report 96-4124, 28 p.

(https://doi.org/10.3133/wri964124https://doi.org/10.3133/wri964124)

Verdi, R.J., and Dixon, J.F.,2011, Magnitude and Frequency of Floods for Rural Streams in Florida, 2006: U.S. Geological Survey Scientific Investigations Report 2011–5034, 69 p., 1 pl. (http://pubs.usgs.gov/sir/2011/5034/)

Inman, E.J., 2000, Lagtime relations for urban streams in Georgia: U.S. Geological Survey Water-Resources Investigations Report 00-4049, 12 p.

(http://ga.water.usgs.gov/pubs/wrir/wrir004049/pdf/wrir00-4049.pdf)

Gotvald, A.J., Feaster, T.D., and Weaver, J.C.,2009, Magnitude and Frequency of Rural Floods in the Southeastern United States, 2006: Volume 1, Georgia: U.S. Geological Survey Scientific Investigations Report 2009-5043, 120 p. (http://pubs.usgs.gov/sir/2009/5043/) Feaster, T.D., Gotvald, A.J., and Weaver, J.C.,2014, Methods for estimating the magnitude and frequency of floods for urban and small, rural streams in Georgia, South Carolina, and North Carolina, 2011 (ver. 1.1, March 2014): U.S. Geological Survey Scientific Investigations Report 2014–5030, 104 p. (http://pubs.usgs.gov/sir/2014/5030/) Gotvald, A.J.,2017, Methods for estimating selected low-flow frequency statistics and mean annual flow for ungaged locations on streams in North Georgia: U.S. Geological Survey Scientific Investigations Report 2017–5001, 25 p.

Oki, D.S., Rosa, S.N., and Yeung, C.W.,2010, Flood-frequency estimates for streams on Kaua'i, O'ahu, Moloka'i, Maui, and Hawai'i, State of Hawai'i: U.S. Geological Survey Scientific Investigations Report 2010-5035, 121 p. (http://pubs.usgs.gov/sir/2010/5035/) Gingerich, S.B.,2005, Median and low-flow characteristics for streams under natural and diverted conditions, northeast Maui, Hawaii: U.S. Geological Survey Scientific Investigations Report 2004-5262, 72 p. (http://pubs.usgs.gov/sir/2004/5262/pdf/sir2004-5262.pdf)

Fontaine, R.A., Wong, M.F., Matsuoka, Iwao,1992, Estimation of Median Streamflows at Perennial Stream Sites in Hawaii: U.S. Geological Survey Water-Resources Investigations Report 92-4099, 37 p. (http://pubs.er.usgs.gov/usgspubs/wri/wri924099)

Hortness, J.E.,2006, Estimating Low-Flow Frequency Statistics for Unregulated Streams in Idaho: U.S. Geological Survey Scientific Investigations Report 2006-5035, 31 p. (http://pubs.usgs.gov/sir/2006/5035/pdf/sir20065035.pdf)

Wood, M.S., Fosness, R.L., Skinner, K.D., and Veilleux, A.G.,2016, Estimating peak-flow frequency statistics for selected gaged and ungaged sites in naturally flowing streams and rivers in Idaho: U.S. Geological Survey Scientific Investigations Report 2016–5083, 56 p. (http://dx.doi.org/10.3133/sir20165083)

Hortness, J.E., and Berenbrock, Charles, 2001, Estimating Monthly and Annual Streamflow Statistics at Ungaged Sites in Idaho: U.S. Geological Survey Water-Resources Investigations Report 01–4093, 36 p. (http://idaho.usgs.gov/PDF/wri014093/index.html)

Over, T.M., Riley, J.D., Sharpe, J.B., and Arvin, Donald,2014, Estimation of regional flow-duration curves for Indiana and Illinois: U.S. Geological Survey Scientific Investigations Report 2014–5177, 24 p. and additional downloads, Tables 2–5, 8–13, and 18 (http://dx.doi.org/10.3133/sir20145177)

Soong, D.T., Ishii, A.L., Sharpe, J.B., and Avery, C.F.,2004, Estimating Flood-Peak Discharge Magnitudes and Frequencies for Rural Streams in Illinois, U.S. Geological Survey Scientific Investigations Report 2004-5103. 147 p. (http://il.water.usgs.gov/pubs/sir2004-5103.pdf) Over, T.M., Saito, R.J., Veilleux, A.G., Sharpe, J.B., Soong, D.T., and Ishii, A.L.,2016, Estimation of peak discharge quantiles for selected annual exceedance probabilities in northeastern Illinois: U.S. Geological Survey Scientific Investigations Report 2016-5050, 50 p. (http://dx.doi.org/10.3133/sir20165050)

Rao, A.R.,2005, Flood-Frequency Relationships for Indiana: Joint Transportation Research Program, Purdue University, FHWA/IN/JTRP-2005/18, 14 p.

(https://docs.lib.purdue.edu/cgi/viewcontent.cgi?article=1746&context=jtrp)

Robinson, B.A.,2013, Regional bankfull-channel dimensions of non-urban wadeable streams in Indiana: U.S. Geological Survey, Scientific Investigations Report 2013-5078, 33 p. (http://pubs.usgs.gov/sir/2013/5078/)

Martin, G.R., Fowler, K.K., and Arihood, L.D.,2016, Estimating selected low-flow frequency statistics and harmonic-mean flows for ungaged, unregulated streams in Indiana (ver 1.1, October 2016): U.S. Geological Survey Scientific Investigations Report 2016-5102, 45 p. (http://dx.doi.org/10.3133/sir20165102)

Arihood, L.D.; Glatfelter, D.R.,1991, Method for estimating low-flow characteristics of ungaged streams in Indiana: U.S. Geological Survey Water-Supply Paper 2372, 19 p. (http://onlinepubs.er.usgs.gov/djvu/WSP/wsp_2372.djvu)

Eash, D.A., and Barnes, K.K.,2012, Methods for estimating selected low-flow frequency statistics and harmonic mean flows for streams in Iowa: U.S. Geological Survey Scientific Investigations Report 2012-5171, 99 p. (http://pubs.usgs.gov/sir/2012/5171/)

Linhart, S.M., Nania, J.F., Sanders, C.L., Jr., and Archfield, S.A.,2012, Computing daily mean streamflow at ungaged locations in Iowa by using the Flow Anywhere and Flow Duration Curve Transfer statistical methods: U.S. Geological Survey Scientific Investigations Report 2012–5232, 50 p. (http://pubs.usgs.gov/sir/2012/5232/)

Eash, D.A., Barnes, K.K., and Veilleux, A.G.,2013, Methods for estimating annual exceedance-probability discharges for streams in Iowa, based on data through water year 2010: U.S. Geological Survey Scientific Investigations Report 2013-5086, 63 p. with a (http://pubs.usgs.gov/sir/2013/5086/)

Eash, D.A.,2015, Comparisons of estimates of annual exceedance-probability discharges for small drainage basins in Iowa, based on data through water year 2013: U.S. Geological Survey Scientific Investigations Report 2015–5055, 37 p.

(http://dx.doi.org/10.3133/sir20155055.)

Eash, D.A., Barnes, K.K., and O'Shea, P.S.,2016, Methods for estimating selected spring and fall low-flow frequency statistics for ungaged stream sites in lowa, based on data through June 2014: U.S. Geological Survey Scientific Investigations Report 2016–5111, 32 p. (http://dx.doi.org/10.3133/sir20165111)

Perry, C.A., Wolock, D.M., and Artman, J.C.,2004, Estimates of Flow Duration, Mean Flow, and Peak-Discharge Frequency Values for Kansas Stream Locations: U.S. Geological Survey Scientific Investigations Report 2004-5033, 651 p.

(http://water.usgs.gov/pubs/sir/2004/5033/pdf/sir2004.5033front.pdf)

Painter, C.C., Heimann, D.C., and Lanning-Rush, J.L., 2017, Methods for estimating annual exceedance-probability streamflows for streams in Kansas based on data through water

```
year 2015: U.S. Geological Survey Scientific Investigations Report 2017-5063, 20 p.
(https://doi.org/10.3133/sir20175063)
Hodgkins, G.A. and Martin, G.R., 2003, Estimating the Magnitude of Peak Flows for Streams
in Kentucky for Selected Recurrence Intervals: U.S. Geological Survey Water-Resources
Investigations Report 03-4180, 69 p. (http://water.usgs.gov/pubs/wri/wri034180/)
Martin, G.R., Ruhl, K.J., Moore, B.L., and Rose, M.F., 1997, Estimation of Peak-Discharge
Frequency of Urban Streams in Jefferson County, Kentucky: U.S. Geological Survey Water-
Resources Investigations Report 97-4219 (http://pubs.er.usgs.gov/publication/wri974219)
Martin, G.R., 2002, Estimating Mean Annual Streamflow of Rural Streams in Kentucky: U.S.
Geological Survey Water-Resources Investigations Report 02-4206, 35 p.
(http://pubs.er.usgs.gov/publication/wri024206)
Martin, G.R., and Arihood, L.D., 2010, Methods for estimating selected low-flow frequency
statistics for unregulated streams in Kentucky: U.S. Geological Survey Scientific
Investigations Report 2010-5217, 83 p. (http://pubs.usgs.gov/sir/2010/5217/)
Martin, G. R. and Ruhl, K. J., 1993, Regionalization of harmonic-mean streamflows in
Kentucky: U.S. Geological Survey Water-Resources Investigations Report 92-4173, 47 p., 1
pl. (http://pubs.er.usgs.gov/publication/wri924173StreamStats_KY_20140226.mdb)
Brockman, R. A., Agouridis, C. T., Workman, S. R., Ormsbee, L. E., Fogle, A. W., 2012,
Bankfull regional curves for the Inner and Outer Bluegrass Regions of Kentucky, Journal of
the American Water Resources Association, v. 48, no. 2, p. 391-406.
(http://onlinelibrary.wiley.com/doi/10.1111/j.1752-1688.2011.00621.x/full)
TR No.70, (2004) Regionalized Regression Equations for Estimating Low-Flow
Characteristics for selected Louisiana Streams
(http://la.water.usgs.gov/publications/pdfs/TR70.pdf)
TR No.60, (1998) Floods in Louisiana, Magnitude and Frequency, Fifth Edition (not
available)
Landers, M.N., 1985, Floodflow Frequency of Streams in the Alluvial Plain of the Lower
Mississippi River in Mississippi, Arkansas, and Louisiana: U.S. Geological Survey Water-
Resources Investigations Report 85-4150, 21 p.
(http://pubs.er.usgs.gov/publication/wri854150)
Lombard, P. J., Tasker, G. D., and Nielsen, M. G., 2003, August Median Streamflow on
Ungaged Streams in Eastern Aroostook County, Maine: U.S. Geological Survey Water-
Resources Investigations Report 03-4225, 20 p.
(http://water.usgs.gov/pubs/wri/wri034225/pdf/wrir03-4225.pdf)
Lombard, P. J., 2004, August Median Streamflow on Ungaged Streams in Eastern Coastal
Maine: U.S. Geological Survey Scientific Investigations Report 2004-5157, 15 p.
(http://water.usgs.gov/pubs/sir/2004/5157/)
Dudley, R.W., 2004, Estimating Monthly, Annual, and Low 7-Day, 10-Year Streamflows for
Ungaged Rivers in Maine: U.S. Geological Survey Scientific Investigations Report 2004-
5026, 22 p. (http://water.usgs.gov/pubs/sir/2004/5026/pdf/sir2004-5026.pdf)
Hodgkins, G. A., 1999, Estimating the Magnitude of Peak Flows for Streams in Maine for
Selected Recurrence Intervals: U.S. Geological Survey Water-Resources Investigations
Report 99-4008, 45 p. (https://pubs.er.usgs.gov/publication/wri994008)
Dudley, R.W., 2004, Hydraulic-Geometry Relations for Rivers in Coastal and Central Maine:
U.S. Geological Survey Scientific Investigations Report 2004-5042, 30 p
(http://pubs.usgs.gov/sir/2004/5042/pdf/sir2004-5042.pdf)
Lombard, P.J., 2010, June and August median streamflows estimated for ungaged streams
in southern Maine: U.S. Geological Survey Scientific Investigations Report 2010-5179, 16
```

p. (http://pubs.usgs.gov/sir/2010/5179/pdf/sir2010-5179.pdf)

Lombard, P.J., and Hodgkins, G.A.,2015, Peak flow regression equations for small, ungaged streams in Maine— Comparing map-based to field-based variables: U.S. Geological Survey Scientific Investigations Report 2015–5049, 12 p. (http://dx.doi.org/10.3133/sir20155049) Dudley, R.W.,2015, Regression equations for monthly and annual mean and selected percentile streamflows for ungaged rivers in Maine: U.S. Geological Survey Scientific Investigations Report 2015–5151, 35 p. (http://dx.doi.org/10.3133/sir20155151) Thomas, Jr., W.O. and Moglen, G.E.,2010, An Update of Regional Regression Equations for Maryland, Appendix 3 in Application of Hydrologic Methods in Maryland, Third Edition, September 2010: Maryland State Highway Administration and Maryland Department of the Environment, 38 p.

(http://gishydro.eng.umd.edu/HydroPanel/hydrology_panel_report_3rd_edition_final.pdf)
Chaplin, J.J.,2005, Development of regional curves relating bankfull-channel geometry and
discharge to drainage area for streams in Pennsylvania and selected areas of Maryland:
U.S. Geological Survey Scientific Investigations Report 2005-5147, 34 p.

(https://pubs.usgs.gov/sir/2005/5147/SIR2005-5147.pdf)

Ries, K.G., III,2000, Methods for estimating low-flow statistics for Massachusetts streams: U.S. Geological Survey Water Resources Investigations Report 00-4135, 81 p. (http://pubs.usgs.gov/wri/wri004135/)

Bent, G.C., and Steeves, P.A.,2006, A revised logistic regression equation and an automated procedure for mapping the probability of a stream flowing perennially in Massachusetts:

U.S. Geological Survey Scientific Investigations Report 2006-5031, 107 p.

(http://pubs.usgs.gov/sir/2006/5031/pdfs/SIR_2006-5031rev.pdf)

Bent, G.C., and Waite, A.M.,2013, Equations for estimating bankfull channel geometry and discharge for streams in Massachusetts: U.S. Geological Survey Scientific Investigations Report 2013–5155, 62 p., (http://pubs.usgs.gov/sir/2013/5155/)

Zarriello, P.J.,2017, Magnitude of flood flows at selected annual exceedance probabilities for streams in Massachusetts: U.S. Geological Survey Scientific Investigations Report 2016–5156, 99 p. (https://dx.doi.org/10.3133/sir20165156)

Holtschlag, D.J. and Croskey, H.M.,1984, Statistical Methods for Estimating Flow Characteristics of Michigan Streams: U.S. Geological Survey Water-Resources Investigations Report 84-4207, 80 p. (http://pubs.er.usgs.gov/usgspubs/wri/wri844207)

Lorenz, D.L., Sanocki, C.A., and Kocian, M.J.,2009, Techniques for Estimating the Magnitude and Frequency of Peak Flows on Small Streams in Minnesota Based on Data through Water Year 2005: U.S. Geological Survey Scientific Investigations Report 2009-5250, 54 p. (http://pubs.usgs.gov/sir/2009/5250/pdf/sir2009-5250.pdf)

Ziegeweid, J.R., Lorenz, D.L., Sanocki, C.A., and Czuba, C.R.,2015, Methods for estimating flow-duration curve and low-flow frequency statistics for ungaged locations on small streams in Minnesota: U.S. Geological Survey Scientific Investigations Report 2015-5170, 23 p. (http://dx.doi.org/10.3133/sir20155170)

Anderson, B.T.,2018, Flood frequency of rural streams in Mississippi, 2013: U.S. Geological Survey Scientific Investigations Report 2018–5148, 12 p.

(https://doi.org/10.3133/sir20185148)

Southard, R.E., and Veilleux, A.G.,2014, Methods for estimating annual exceedance-probability discharges and largest recorded floods for unregulated streams in rural Missouri: U.S. Geological Survey Scientific Investigations Report 2014–5165, 39 p. (http://pubs.usgs.gov/sir/2014/5165/)

Southard, R.E., 2013, Computed statistics at streamgages, and methods for estimating low-flow frequency statistics and development of regional regression equations for estimating

low-flow frequency statistics at ungaged locations in Missouri: U.S. Geological Survey Scientific Investigations Report 2013–5090, 28 p. (http://pubs.usgs.gov/sir/2013/5090/) Parrett, Charles and Hull, J.A.,1985, A method for estimating mean and low flows of streams in national forests of Montana: U.S. Geological Survey Water-Resources Investigations Report 85-4071, 13 p. (https://pubs.er.usgs.gov/publication/wri854071) Parrett, Charles and Cartier, K.D. ,1999, Methods for estimating monthly streamflow characteristics at ungaged sites in western Montana: U.S. Geological Survey Water-Supply Paper 2365, 30 p. (http://pubs.er.usgs.gov/publication/wsp2365)

Parrett, Charles and Johnson, D.R.,2004, Methods for Estimating Flood Frequency in Montana Based on Data through Water Year 1998: U.S. Geological Survey Water-Resources Investigations Report 03-4308, 102 p. (http://water.usgs.gov/pubs/wri/wri03-4308/) Sando, Roy, Sando, S.K., McCarthy, P.M., and Dutton, D.M.,2016, Methods for estimating peak-flow frequencies at ungaged sites in Montana based on data through water year 2011: U.S. Geological Survey Scientific Investigations Report 2015-5019-F, 30 p. (https://doi.org/10.3133/sir20155019)

McCarthy, P.M., Sando, Roy, Sando, S.K., and Dutton, D.M.,2016, Methods for estimating streamflow characteristics at ungaged sites in western Montana based on data through water year 2009: U.S. Geological Survey Scientific Investigations Report 2015–5019–G, 19 p. (https://doi.org/10.3133/sir20155019)

Soenksen, P.J., Miller, L.D., Sharpe, J.B. and Watton, J.R.,1999, Peak-Flow Frequency Relations and Evaluation of the Peak-Flow Gaging Network in Nebraska: U. S. Geological Survey Water-Resources Investigations Report 99-4032, 48 p,

(https://pubs.er.usgs.gov/publication/wri994032)

Flynn, R.H. and Tasker, G.D.,2002, Development of Regression Equations to Estimate Flow Durations and Low-Flow-Frequency Statistics in New Hampshire Streams: U.S.Geological Survey Scientific Investigations Report 02-4298, 66 p. (http://pubs.water.usgs.gov/wrir02-4298)

Olson, S.A.,2009, Estimation of flood discharges at selected recurrence intervals for streams in New Hampshire: U.S.Geological Survey Scientific Investigations Report 2008-5206, 57 p. (http://pubs.usgs.gov/sir/2008/5206/)

Flynn, R.H. and Tasker, G.D.,2004, Generalized Estimates from Streamflow Data of Annual and Seasonal Ground-Water-Recharge Rates for Drainage Basins in New Hampshire, U.S. Geological Survey Scientific Investigations Report 2004-5019, 67 p.

(http://pubs.usgs.gov/sir/2004/5019/http://pubs.usgs.gov/sir/2004/5019/)

Watson, K.M., and Schopp, R.D., 2009, Methodology for estimation of flood magnitude and frequency for New Jersey streams, U.S. Geological Survey Scientific Investigations Report 2009-5167, 51 p. (http://pubs.usgs.gov/sir/2009/5167/)

Watson, K.M., and McHugh, A.R.,2014, Regional regression equations for the estimation of selected monthly low-flow duration and frequency statistics at ungaged sites on streams in New Jersey: U.S. Geological Survey Scientific Investigations Report 2014-5004, 59 p. (baseline, period-or-record statistics)

(http://dx.doi.org/10.3133/sir20145004StreamStatsDB\2019_12_13_DataSource_table.xlsxDa Waltemeyer, S.D.,2002, Analysis of the magnitude and frequency of the 4-day, 3-year annual low flow on unregulated streams in New Mexico: U. S. Geological Survey Water-Resources Investigations Report 01-4271, 22 p.

(http://nm.water.usgs.gov/publications/abstracts/wrir01-4271.pdf)

Waltemeyer, S.D.,2008, Analysis of the Magnitude and Frequency of Peak Discharge and Maximum Observed Peak Discharge in New Mexico and Surrounding Areas: U.S. Geological

Survey Scientific Investigations Report 2008-5119, 105 p.

(http://pubs.usgs.gov/sir/2008/5119/)

Lumia, Richard, Freehafer, D.A., and Smith, M.J.,2006, Magnitude and Frequency of Floods in New York: U.S. Geological Survey Scientific Investigations Report 2006–5112, 152 p. (http://pubs.usgs.gov/sir/2006/5112/)

Stedfast, D.A.,1984, Evaluation of Six Methods for Estimating Magnitude and Frequency of Peak Discharges on Urban Streams in New York: U. S. Geological Survey Water-Resources Investigations Report 84-4350, 24 p.

(http://onlinepubs.er.usgs.gov/djvu/WRI/wrir_84_4350.djvu)

Mulvihill, C.I., Baldigo, B.P., Miller, S.J., and DeKoskie, Douglas,2009, Bankfull Discharge and Channel Characteristics of Streams in New York State: U.S. Geological Survey Scientific Investigations Report 2009-5144, 51 p. (http://pubs.usgs.gov/sir/2009/5144/) Barnes, C. R.,1986, Method for estimating low-flow statistics for ungaged streams in the lower Hudson River Basin, New York: U. S. Geological Survey Water-Resources Investigations Report 85-4070, 22 p.

(http://onlinepubs.er.usgs.gov/djvu/WRI/wrir_85_4070.djvu)

Randall, A.D., 2010, Low flow of streams in the Susquehanna River basin of New York: U.S. Geological Survey Scientific Investigations Report 2010-5063, 57 p.

(http://pubs.usgs.gov/sir/2010/5063/http://pubs.usgs.gov/sir/2010/5063/)

Gazoorian, C.L.,2015, Estimation of unaltered daily mean streamflow at ungaged streams of New York, excluding Long Island, water years 1961–2010: U.S. Geological Survey Scientific Investigations Report 2014–5220, 29 p. (https://pubs.usgs.gov/sir/2014/5220/) Giese, G. L. and Mason, R.R., Jr.,1993, Low-flow characteristics of streams in North Carolina: U.S. Geological Survey Water-Supply Paper 2403, 29 p.

(https://pubs.er.usgs.gov/publication/wsp2403)

Mason, Robert R., Jr.; Fuste, Luis A.; King, Jeffrey N.; Thomas, Wilbert O., Jr.,2002, The National Flood-Frequency Program -- Methods for Estimating Flood Magnitude and Frequency in Rural and Urban Areas in North Carolina, 2001: U.S. Geological Survey Fact Sheet 007-00, 4 p. (http://pubs.er.usgs.gov/publication/fs00700)

Weaver, J.C., Feaster, T.D., and Gotvald, A.J.,2009, Magnitude and frequency of rural floods in the Southeastern United States, through 2006—Volume 2, North Carolina: U.S. Geological Survey Scientific Investigations Report 2009–5158, 111 p.

(http://pubs.usgs.gov/sir/2009/5158/)

Williams-Sether, T.,2015, Regional regression equations to estimate peak-flow frequency at sites in North Dakota using data through 2009: U.S. Geological Survey Scientific Investigations Report 2015–5096, 12 p. (http://dx.doi.org/10.3133/sir20155096)

Koltun, G.F., Kula, S.P., and Puskas, B.M.,2006, A Streamflow Statistics (StreamStats) Web Application for Ohio: U.S. Geological Survey Scientific Investigations Report 2006-5312, 62 p. (http://pubs.usgs.gov/sir/2006/5312/)

Sherwood, J.M.,1994, Estimation of peak-frequency relations, flood hydrographs, and volume-duration-frequency relations of ungaged small urban streams in Ohio: U. S. Geological Survey Water-Supply Paper 2432, 42 p.

(http://pubs.er.usgs.gov/usgspubs/wsp/wsp2432)

Koltun, G. F., and Whitehead, M. T.,2002, Techniques for Estimating Selected Streamflow Characteristics of Rural, Unregulated Streams in Ohio: U. S. Geological Survey Water-Resources Investigations Report 02-4068, 50 p

(https://pubs.er.usgs.gov/publication/wri024068)

Koltun, G. F., and Schwartz, Ronald R.,1987, MULTIPLE-REGRESSION EQUATIONS FOR ESTIMATING LOW FLOWS AT UNGAGED STREAM SITES IN OHIO: U.S. Geological Survey

Water-Resources Investigations Report 86-4354, 39 p. (http://pubs.er.usgs.gov/usgspubs/wri/wri864354)

Koltun, G.F., and Kula, S.P.,2013, Methods for estimating selected low-flow statistics and development of annual flow-duration statistics for Ohio: U.S. Geological Survey Scientific Investigations Report 2012–5138, 195 p. (http://pubs.usgs.gov/sir/2012/5138/)

Koltun, G.F.,2019, Flood-frequency estimates for Ohio streamgages based on data through water year 2015 and techniques for estimating flood-frequency characteristics of rural, unregulated Ohio streams: U.S. Geological Survey Scientific Investigations Report 2019–5018, xx p. (https://dx.doi.org/10.3133/sir20195018)

Esralew, R.A., Smith, S.J.,2009, Methods for estimating flow-duration and annual mean-flow statistics for ungaged streams in Oklahoma: U.S. Geological Survey Scientific Investigations Report 2009-5267, 131 p. (http://pubs.usgs.gov/sir/2009/5267/) Smith, S.J., Lewis, J.M., and Graves, G.M.,2015, Methods for estimating the magnitude and frequency of peak streamflows at ungaged sites in and near the Oklahoma Panhandle: U.S. Geological Survey Scientific Investigations Report 2015-5134, 35 p.

(http://dx.doi.org/10.3133/sir20155134)

Lewis, J.M., Hunter, S.L., and Labriola, L.G.,2019, Methods for estimating the magnitude and frequency of peak streamflows for unregulated streams in Oklahoma developed by using streamflow data through 2017: U.S. Geological Survey Scientific Investigations Report 2019-5143, 39 p. (https://doi.org/10.3133/sir20195143)

Laenen, Antonius, 1980, Storm Runoff As Related to Urbanization in the Portland, Oregon - Vancouver, Washington Area: U.S. Geological Survey Open-File Report 80-689, 71 p. (http://pubs.er.usgs.gov/usgspubs/wri/wri834143)

Cooper, R.M.,2005, Estimation of Peak Discharges for Rural, Unregulated Streams in Western Oregon: U.S. Geological Survey Scientific Investigations Report 2005-5116, 76 p. (http://pubs.usgs.gov/sir/2005/5116/pdf/sir2005-5116.pdf)

Risley, John, Stonewall, Adam, and Haluska, Tana, 2008, Estimating flow-duration and low-flow frequency statistics for unregulated streams in Oregon: U.S. Geological Survey Scientific Investigations Report 2008-5126, 22 p. (http://pubs.usgs.gov/sir/2008/5126/) Cooper, Richard, 2006, Estimation of Peak Discharges for Rural, Unregulated Streams in Eastern Oregon, Oregon Water Resources Department OFR SW 06-001, Salem, OR. (https://digital.osl.state.or.us/islandora/object/osl%3A14736/datastream/OBJ/view) Stuckey, M.H., 2006, Low-flow, base-flow, and mean-flow regression equations for Pennsylvania streams: U.S. Geological Survey Scientific Investigations Report 2006-5130, 84 p. (http://pubs.usgs.gov/sir/2006/5130/)

Stuckey, M.H., Koerkle, E.H., and Ulrich, J.E.,2012, Estimation of baseline daily mean streamflows for ungaged locations on Pennsylvania streams, water years 1960–2008: U.S. Geological Survey Scientific Investigations Report 2012–5142, 61 p. (http://pubs.usgs.gov/sir/2012/5142/)

Clune, J.W., Chaplin, J.J., and White, K.E., 2018, Comparison of regression relations of bankfull discharge and channel geometry for the glaciated and nonglaciated settings of Pennsylvania and southern New York: U.S. Geological Survey Scientific Investigations Report 2018-5066, 20 p. (https://doi.org/10.3133/sir20185066)

Roland, M.A., and Stuckey, M.H.,2008, Regression equations for estimating flood flows at selected recurrence intervals for ungaged streams in Pennsylvania: U.S. Geological Survey Scientific Investigations Report 2008-5102, 57p. (http://pubs.usgs.gov/sir/2008/5102/) Zarriello, P.J., Ahearn, E.A., and Levin, S.B.,2012, Magnitude of flood flows for selected annual exceedance probabilities in Rhode Island through 2010: U.S. Geological Survey Scientific Investigations Report 2012-5109, 93 p. (http://pubs.usgs.gov/sir/2012/5109)

```
Bent, G.C., Steeves, P.A., and Waite, A.M.,2014, Equations for estimating selected streamflow statistics in Rhode Island: U.S. Geological Survey Scientific Investigations Report 2014-5010, 65 p. (http://dx.doi.org/10.3133/sir20145010)
```

Feaster, T.D., Gotvald, A.J., and Weaver, J.C.,2009, Magnitude and Frequency of Rural Floods in the Southeastern United States, 2006: Volume 3, South Carolina: U.S. Geological Survey Scientific Investigations Report 2009-5156, 226 p.

(http://pubs.usgs.gov/sir/2009/5156/)

Sando, Steven K.,1998, A Method for Estimating Magnitude and Frequency of Floods in South Dakota: U.S. Geological Survey Water-Resources Investigations Report 98-4055, 48 p. (http://pubs.water.usgs.gov/wri98-4055/)

Law, G.S., and Tasker G.D.,2003, Flood-Frequency Prediction Methods for Unregulated Streams of Tennessee, 2000: U.S. Geological Survey Water-Resources Investigations Report 03-4176, 79p. (http://pubs.usgs.gov/wri/wri034176/)

Neely, B.L., Jr.,1984, Flood Frequency and Storm Runoff of Urban Areas of Memphis and Shelby County, Tennessee: U.S. Geological Survey Water-Resources Investigations Report 84-4110, 51 p. (http://pubs.usgs.gov/wri/wrir_84-4110/)

Robbins, Clarence H.,1984, Synthesized Flood Frequency of Small Urban Streams in Tennessee: U.S. Geological Survey Water-Resources Investigations Report 84-4182, 24 p. (http://onlinepubs.er.usgs.gov/djvu/WRI/wrir_84_4182.djvu)

Law, G.S., Tasker, G.D., and Ladd, D.E., 2009, Streamflow-characteristic estimation methods for unregulated streams of Tennessee: U.S. Geological Survey Scientific Investigations Report 2009–5159, 212 p., 1 pl. (http://pubs.usgs.gov/sir/2009/5159/)

Asquith, W.H., Slade, R.M., Jr.,1999, Site-specific estimation of peak-stream flow frequency using generalized least squares regression for natural basins in Texas: U.S. Geological Survey Water-Resources Investigations Report 99-4172, 19 p. (http://pubs.water.usgs.gov/wri994172)

Asquith, William H.,1998, Peak-flow frequency for tributaries of the Colorado River downstream of Austin, Texas U.S. Geological Survey Water-Resources Investigations Report 98-4015, 26 p. (http://pubs.water.usgs.gov/wri98-4015/)

Raines, Timothy H.,1998, Peak-discharge frequency and potential extreme peak discharge for natural streams in the Brazos River basin, Texas: U.S. Geological Survey Water-Resources Investigations Report 98-4178, 47 p., 1 plate (http://pubs.water.usgs.gov/wri98-4178/)

Land, L.F., Schroeder, E.E. and Hampton, B.B.,1982, Techniques for Estimating the Magnitude and Frequency of Floods in the Dallas-Fort Worth Metropolitan Area, Texas: U.S. Geological Survey Water-Resources Investigations Report 82-18, 55 p. ()

Asquith, W.H., Slade, R. M., Lanning-Rush, Jennifer,1996, Peak-flow frequency and extreme flood potential for streams in the vicinity of the Highland Lakes, central Texas: U.S.

Geological Survey Water-Resources Investigations Report 96-4072

(http://onlinepubs.er.usgs.gov/djvu/WRI/wrir_96_4072_plt.djvu)

Liscum, Fred and Massey, B.C.,1980, Technique for Estimiating the Magnitude and Frequency of Floods in the Houston, Texas, Metropolitan Area: U.S. Geological Survey Water-Resources Investigations Report 80-17, 29 p. ()

Asquith, W.H., and Roussel, M.C.,2009, Regression equations for estimation of annual peak-streamflow frequency for undeveloped watersheds in Texas using an L-moment-based, PRESS-minimized, residual-adjusted approach: U.S. Geological Survey Scientific Investigations Report 2009–5087, 48 p. (http://pubs.usgs.gov/sir/2009/5087/)

Kenney, T.A., Wilkowske, C.D., and Wright, S.J.,2007, Methods for Estimating Magnitude and Frequency of Peak Flows for Natural Streams in Utah: U.S. Geological Survey Scientific

```
Investigations Report 2007-5158, 28 p. (http://pubs.usgs.gov/sir/2007/5158/)
Wilkowske, C.D., Kenney, T.A., and Wright, S.J., 2009, Methods for Estimating Monthly and
Annual Streamflow Statistics at Ungaged Sites in Utah: U.S. Geological Survey Scientific
Investigations Report 2008-5230, 62 p. (http://pubs.usgs.gov/sir/2008/5230/)
Olson, S.A., 2002, Flow-frequency characteristics of Vermont streams: U.S. Geological
Survey Water-Resources Investigations Report 02-4238, 47 p.
(http://pubs.usgs.gov/wri/wrir02-4238/)
Olson, S.A., 2014, Estimation of flood discharges at selected annual exceedance
probabilities for unregulated, rural streams in Vermont, with a section on Vermont regional
skew regression, by Veilleux, A.G.: U.S. Geological Survey Scientific Investigations Report
2014-5078, 27 p. plus appendixes. (http://pubs.usgs.gov/sir/2014/5078/)
Olson, S.A., and Brouillette, M.C., 2006, A logistic regression equation for estimating the
probability of a stream in Vermont having intermittent flow: U.S. Geological Survey
Scientific Investigations Report 2006-5217, 15 p. (https://pubs.usgs.gov/sir/2006/5217/)
Austin, S.H., Krstolic, J.L., and Wiegand, Ute, 2011, Low-flow characteristics of Virginia
streams: U.S. Geological Survey Scientific Investigations Report 2011-5143, 122 p. + 9
tables on CD. (http://pubs.usgs.gov/sir/2011/5143/)
Austin, S.H., Krstolic, J.L., and Wiegand, Ute, 2011, Peak-flow characteristics of Virginia
streams: U.S. Geological Survey Scientific Investigations Report 2011-5144, 106 p. + 3
tables and 2 appendixes on CD. (http://pubs.usgs.gov/sir/2011/5144/)
Austin, S.H., 2014, Methods and equations for estimating peak streamflow per square mile
in Virginia's urban basins: U.S. Geological Survey Scientific Investigations Report 2014-
5090, 25 p. (http://pubs.usgs.gov/sir/2014/5090/http://pubs.usgs.gov/sir/2014/5090/)
Curran, C.A. and Olsen, T.D., 2009, Estimating Low-Flow Frequency Statistics and
Hydrologic Analysis of Selected Streamflow-Gaging Stations, Nooksack River Basin,
Northwestern Washington and Canada: U.S. Geological Survey Scientific Investigations
Report 2009-5170, 44 p. (http://pubs.usgs.gov/sir/2009/5170/)
Curran, C.A., Eng, Ken, and Konrad, C.P., 2012, Analysis of low flows and selected methods
for estimating low-flow characteristics at partial-record and ungaged stream sites in
western Washington: U.S. Geological Survey Scientific Investigations Report 2012-5078, 46
p. (http://pubs.usgs.gov/sir/2012/5078/)
Mastin, M.C., Konrad, C.P., Veilleux, A.G., and Tecca, A.E., 2016, Magnitude, frequency, and
trends of floods at gaged and ungaged sites in Washington, based on data through water
year 2014 (ver 1.1, October 2016): U.S. Geological Survey Scientific Investigations Report
2016-5118, 70 p. (http://dx.doi.org/10.3133/sir20165118)
Wiley, Jeffrey B., 2008, Estimating Selected Streamflow Statistics Representative of 1930-
2002 in West Virginia: U.S. Geological Survey Scientific Investigations Report 2008-5105,
24 p. (http://pubs.usgs.gov/sir/2008/5105/)
Wiley, Jeffrey B., 1987, Techniques for estimating flood depth frequency relations for
streams in West Virginia: U.S. Geological Survey Water-Resources Investigations Report
87-4111, 17 p. (http://pubs.er.usgs.gov/usgspubs/wri/wri874111)
Wiley, J.B., and Atkins, J.T., Jr., 2010, Estimation of flood-frequency discharges for rural,
unregulated streams in West Virginia: U.S. Geological Survey Scientific Investigations
Report 2010-5033, 78 p. (http://pubs.usgs.gov/sir/2010/5033/)
Wiley, J.B., and Atkins, J.T., Jr., 2010, Estimation of selected seasonal streamflow statistics
representative of 1930-2002 in West Virginia: U.S. Geological Survey Scientific
Investigations Report 2010-5185, 20 p. (http://pubs.usgs.gov/sir/2010/5185/)
Conger, Duane H., 1986, Estimating Magnitude and Frequency of Floods for Wisconsin
```

Urban Streams: U.S. Geological Survey Water-Resources Investigations Report 86-4005, 18

p. (http://pubs.er.usgs.gov/publication/wri864005)

Walker, J.F., Peppler, M.C., Danz, M.E., and Hubbard, L.E., 2017, Flood-frequency characteristics of Wisconsin streams (ver. 2.1, December 2017): Reston, Virginia, U.S. Geological Survey Scientific Investigations Report 2016–5140, 33 p., 1 plate, 2 appendixes (https://doi.org/10.3133/sir20165140)

Miller, Kirk A.,2003, Peak-flow Characteristics of Wyoming Streams: U.S. Geological Survey Water-Resources Investigations Report 03-4107, 79 p.

(http://pubs.usgs.gov/wri/wri034107/)

Ramos-Ginés, Orlando,1999, Estimation of Magnitude and Frequency of Floods for Streams in Puerto Rico: New Empirical Models: U. S. Geological Survey Water-Resources Investigations Report 99-4142, 41 p. (http://pubs.usgs.gov/wri/wri994142/)

Moody, J.A., 2012, An analytical method for predicting postwildfire peak discharges: U.S. Geological Survey Scientific Investigations Report 2011-5236, 36 p.

(https://pubs.usgs.gov/sir/2011/5236/)

Flow-Duration Statistics Parameters [Statewide Low Flow WRIR00 4135]

Parameter Code	Parameter Name	Value	Units	Min Limit	Max Limit
DRNAREA	Drainage Area	0.0906	square miles	1.61	149
DRFTPERSTR	Stratified Drift per Stream Length	-100000	square mile per mile	0	1.29
MAREGION	Massachusetts Region	1	dimensionless	0	1
BSLDEM250	Mean Basin Slope from 250K DEM	4.012	percent	0.32	24.6

Flow-Duration Statistics Flow Report[Statewide Low Flow WRIR00 4135]

Statistic Value Unit

Flow-Duration Statistics Citations

Sauer, Vernon B.; Thomas, W. O., Jr.; Stricker, V. A.; Wilson, K. V.,1983, Flood characteristics of urban watersheds in the United States: U.S. Geological Survey Water-Supply Paper 2207, 63 p. (http://pubs.er.usgs.gov/publication/wsp2207)
()

Anderson, B.T.,2020, Magnitude and frequency of floods in Alabama, 2015: U.S. Geological Survey Scientific Investigations Report 2020-5032, 148 p.

(https://doi.org/10.3133/sir20205032)

Hedgecock, T.S.,2004, Magnitude and Frequency of Floods on Small Rural Streams in Alabama: U. S. Geological Survey Scientific Investigations Report 2004-5135, 10 p. (http://pubs.usgs.gov/sir/2004/5135/)

Hedgecock, T.S., 2010, Magnitude and Frequency of Floods for Urban Streams in Alabama, 2007: U.S Geological Survey Scientific Investigations Report 2010-5012, 17p.

(https://pubs.usgs.gov/sir/2010/5012/)

Wiley, J.B., and Curran, J.H.,2003, Estimating annual high-flow statistics and monthly and seasonal low-flow statistics for ungaged sites on streams in Alaska and conterminous basins in Canada: U.S. Geological Survey Water-Resources Investigations Report 03-4114, 61 p. (http://water.usgs.gov/pubs/wri/wri034114/pdf/wri034114_v1.10.pdf)
Brabets, Timothy P.,1996, Evaluation of the streamflow-gaging network of Alaska in providing regional streamflow information: U.S. Geological Survey Water-Resources Investigations Report 96-4001, 98 p. (http://pubs.er.usgs.gov/usgspubs/wri/wri964001)
Curran, J.H., Barth, N.A., Veilleux, A.G., and Ourso, R.T.,2016, Estimating Flood Magnitude and Frequency at Gaged and Ungaged Sites on Streams in Alaska and Conterminous Basins in Canada, Based on Data through Water Year 2012: U.S. Geological Survey Scientific Investigations Report 2016-5024, 47 p.

(http://dx.doi.org/10.3133/sir20165024http://dx.doi.org/10.3133/sir20165024)
Southard, R.E.,2010, Estimation of the Magnituude and Frequency of Floods in Urban
Basins in Missouri: U.S. Geological Survey Scientific Investigations Report 2010-5073, 27
p. (http://pubs.usgs.gov/sir/2010/5073/)

Waltemeyer, S.D., Analysis of the Magnitude and Frequency of Peak Discharges for the Navajo Nation in Arizona, Utah, Colorado, and New Mexico: U. S. Geological Survey Scientific Investigations Report2006-5306, 42 p. (http://pubs.usgs.gov/sir/2006/5306/) Paretti, N.V., Kennedy, J.R., Turney, L.A., and Veilleux, A.G.,2014, Methods for estimating magnitude and frequency of floods in Arizona, developed with unregulated and rural peakflow data through water year 2010: U.S. Geological Survey Scientific Investigations Report 2014-5211, 61 p., http://dx.doi.org/10.3133/sir20145211.

(http://pubs.usgs.gov/sir/2014/5211/)

Kennedy, J.R., Paretti, N.V., and Veilleux, A.G.,2014, Methods for estimating magnitude and frequency of 1-, 3-, 7-, 15-, and 30-day flood-duration flows in Arizona: U.S. Geological Survey Scientific Investigations Report 2014–5109, 35 p.

(http://pubs.usgs.gov/sir/2014/5109/)

Funkhouser, J.E., Eng, Ken, and Moix, M.W.,2008, Low-Flow Characteristics and Regionalization of Low Flow Characteristics for Selected Streams in Arkansas: U. S. Geological Survey Scientific Investigations Report 2008-5065, 161 p. (http://pubs.usgs.gov/sir/2008/5065/pdf/SIR2008-5065.pdf)

Breaker, B.K.,2015, Dry season mean monthly flow and harmonic mean flow regression equations for selected ungaged basins in Arkansas: U.S. Geological Survey Scientific Investigations Report 2015–5031, 25 p. (http://pubs.usgs.gov/sir/2015/5031/)

Wagner, D.M., Krieger, J.D., and Veilleux, A.G.,2016, Methods for estimating annual exceedance probability discharges for streams in Arkansas, based on data through water year 2013: U.S. Geological Survey Scientific Investigations Report 2016-5081, 136 p. (http://dx.doi.org/10.3133/sir20165081)

Thomas, B.E, Hjalmarson, H.W., and Waltemeyer, S.D.,1997, Methods for Estimating Magnitude and Frequency of Floods in the Southwestern United States: U.S. Water-Supply Paper 2433, 196 p. (http://pubs.er.usgs.gov/publication/wsp2433)

Gotvald, A.J., Barth, N.A., Veilleux, A.G., and Parrett, Charles, 2012, Methods for determining magnitude and frequency of floods in California, based on data through water year 2006: U.S. Geological Survey Scientific Investigations Report 2012–5113, 38 p., 1 pl. (http://pubs.usgs.gov/sir/2012/5113/)

Sanocki, C.A., Williams-Sether, T., Steeves, P.A., and Christensen, V.G.,2019, Techniques for Estimating the Magnitude and Frequency of Peak Flows on Small Streams in the Binational U.S. and Canadian Lake of the Woods-Rainy River Basin Upstream from Kenora,

Ontario, Canada, Based on Data through Water Year 2013 : U.S. Geological Survey Scientific Investigations Report 2019-5012, 17 p. (https://doi.org/10.3133/sir20195012)

Capesius, J.P., and Stephens, V. C.,2009, Regional Regression Equations for Estimation of Natural Streamflow Statistics in Colorado: U. S. Geological Survey Scientific Investigations Report 2009-5136, 32 p.

(http://pubs.usgs.gov/sir/2009/5136/http://pubs.usgs.gov/sir/2009/5136/)

Kohn, M.S., Stevens, M.R., Harden, T.M., Godaire, J.E., Klinger, R.E., and Mommandi, A.,2016, Paleoflood investigations to improve peak-streamflow regional-regression equations for natural streamflow in eastern Colorado, 2015: U.S. Geological Survey Scientific Investigations Report 2016–5099, 58 p. (http://dx.doi.org/10.3133/sir20165099) Ahearn, E.A.,2004, Regression Equations for Estimating Flood Flows for the 2-, 10-, 25-, 50-, 100-, and 500-Year Recurrence Intervals in Connecticut: U.S. Geological Survey SRI 2004-5160, 62 p. (http://water.usgs.gov/pubs/sir/2004/5160/)

Ahearn, E.A.,2010, Regional regression equations to estimate flow-duration statistics in Connecticut: U. S. Geological Survey Scientific Investigations Report 2010-5052, 45 p. (http://pubs.usgs.gov/sir/2010/5052/)

Ries, K.G., III, and Dillow, J.J.A.,2006, Magnitude and frequency of floods in Delaware: Scientific Investigations Report 2006-5146, 59 p. (http://pubs.usgs.gov/sir/2006/5146/) Carpenter, D.H., and Hayes, D.C.,1996, Low-flow characteristics of streams in Maryland and Delaware: U.S. Geological Survey Water-Resources Investigations Report 94-4020, 113 p., 10 plates (http://pubs.er.usgs.gov/usgspubs/wri/wri944020)

Franklin, M.A. and Losey, G.T.,1984, Magnitude and Frequency of Floods from Urban Streams in Leon County, Florida: U.S. Geological Survey Water-Resources Investigations Report 84-4004, 37 p. (http://pubs.er.usgs.gov/publication/wri844004)

Lopez, M.A. and Woodham, W. M.,1983, Magnitude and frequency of flooding on small urban watersheds in the Tampa Bay area, west-central Florida: U.S. Geological Survey Water-Resources Investigations Report 82-42, 52 p.

(https://pubs.er.usgs.gov/publication/wri8242)

Rumenik, R. P.; Grubbs, J. W.,1996, Methods for estimating low-flow characteristics of ungaged streams in selected areas, northern Florida: U.S. Geological Survey Water-Resources Investigations Report 96-4124, 28 p.

(https://doi.org/10.3133/wri964124https://doi.org/10.3133/wri964124)

Verdi, R.J., and Dixon, J.F.,2011, Magnitude and Frequency of Floods for Rural Streams in Florida, 2006: U.S. Geological Survey Scientific Investigations Report 2011-5034, 69 p., 1 pl. (http://pubs.usgs.gov/sir/2011/5034/)

Inman, E.J.,2000, Lagtime relations for urban streams in Georgia: U.S. Geological Survey Water-Resources Investigations Report 00-4049, 12 p.

(http://ga.water.usgs.gov/pubs/wrir/wrir004049/pdf/wrir00-4049.pdf)

Gotvald, A.J., Feaster, T.D., and Weaver, J.C.,2009, Magnitude and Frequency of Rural Floods in the Southeastern United States, 2006: Volume 1, Georgia: U.S. Geological Survey Scientific Investigations Report 2009-5043, 120 p. (http://pubs.usgs.gov/sir/2009/5043/) Feaster, T.D., Gotvald, A.J., and Weaver, J.C.,2014, Methods for estimating the magnitude and frequency of floods for urban and small, rural streams in Georgia, South Carolina, and North Carolina, 2011 (ver. 1.1, March 2014): U.S. Geological Survey Scientific Investigations Report 2014–5030, 104 p. (http://pubs.usgs.gov/sir/2014/5030/) Gotvald, A.J.,2017, Methods for estimating selected low-flow frequency statistics and mean annual flow for ungaged locations on streams in North Georgia: U.S. Geological

(https://doi.org/10.3133/sir20175001)

Survey Scientific Investigations Report 2017-5001, 25 p.

Oki, D.S., Rosa, S.N., and Yeung, C.W.,2010, Flood-frequency estimates for streams on Kaua'i, O'ahu, Moloka'i, Maui, and Hawai'i, State of Hawai'i: U.S. Geological Survey Scientific Investigations Report 2010-5035, 121 p. (http://pubs.usgs.gov/sir/2010/5035/) Gingerich, S.B.,2005, Median and low-flow characteristics for streams under natural and diverted conditions, northeast Maui, Hawaii: U.S. Geological Survey Scientific Investigations Report 2004-5262, 72 p. (http://pubs.usgs.gov/sir/2004/5262/pdf/sir2004-5262.pdf)

Fontaine, R.A., Wong, M.F., Matsuoka, Iwao,1992, Estimation of Median Streamflows at Perennial Stream Sites in Hawaii: U.S. Geological Survey Water-Resources Investigations Report 92-4099, 37 p. (http://pubs.er.usgs.gov/usgspubs/wri/wri924099)

Hortness, J.E.,2006, Estimating Low-Flow Frequency Statistics for Unregulated Streams in Idaho: U.S. Geological Survey Scientific Investigations Report 2006-5035, 31 p. (http://pubs.usgs.gov/sir/2006/5035/pdf/sir20065035.pdf)

Wood, M.S., Fosness, R.L., Skinner, K.D., and Veilleux, A.G.,2016, Estimating peak-flow frequency statistics for selected gaged and ungaged sites in naturally flowing streams and rivers in Idaho: U.S. Geological Survey Scientific Investigations Report 2016–5083, 56 p. (http://dx.doi.org/10.3133/sir20165083)

Hortness, J.E., and Berenbrock, Charles, 2001, Estimating Monthly and Annual Streamflow Statistics at Ungaged Sites in Idaho: U.S. Geological Survey Water-Resources Investigations Report 01–4093, 36 p. (http://idaho.usgs.gov/PDF/wri014093/index.html) Over, T.M., Riley, J.D., Sharpe, J.B., and Arvin, Donald, 2014, Estimation of regional flow-duration curves for Indiana and Illinois: U.S. Geological Survey Scientific Investigations Report 2014–5177, 24 p. and additional downloads, Tables 2–5, 8–13, and 18 (http://dx.doi.org/10.3133/sir20145177)

Soong, D.T., Ishii, A.L., Sharpe, J.B., and Avery, C.F.,2004, Estimating Flood-Peak Discharge Magnitudes and Frequencies for Rural Streams in Illinois, U.S. Geological Survey Scientific Investigations Report 2004-5103. 147 p. (http://il.water.usgs.gov/pubs/sir2004-5103.pdf) Over, T.M., Saito, R.J., Veilleux, A.G., Sharpe, J.B., Soong, D.T., and Ishii, A.L.,2016, Estimation of peak discharge quantiles for selected annual exceedance probabilities in northeastern Illinois: U.S. Geological Survey Scientific Investigations Report 2016-5050, 50 p. (http://dx.doi.org/10.3133/sir20165050)

Rao, A.R.,2005, Flood-Frequency Relationships for Indiana: Joint Transportation Research Program, Purdue University, FHWA/IN/JTRP-2005/18, 14 p.

(https://docs.lib.purdue.edu/cgi/viewcontent.cgi?article=1746&context=jtrp)

Robinson, B.A.,2013, Regional bankfull-channel dimensions of non-urban wadeable streams in Indiana: U.S. Geological Survey, Scientific Investigations Report 2013-5078, 33 p. (http://pubs.usgs.gov/sir/2013/5078/)

Martin, G.R., Fowler, K.K., and Arihood, L.D.,2016, Estimating selected low-flow frequency statistics and harmonic-mean flows for ungaged, unregulated streams in Indiana (ver 1.1, October 2016): U.S. Geological Survey Scientific Investigations Report 2016-5102, 45 p. (http://dx.doi.org/10.3133/sir20165102)

Arihood, L.D.; Glatfelter, D.R.,1991, Method for estimating low-flow characteristics of ungaged streams in Indiana: U.S. Geological Survey Water-Supply Paper 2372, 19 p. (http://onlinepubs.er.usgs.gov/djvu/WSP/wsp_2372.djvu)

Eash, D.A., and Barnes, K.K.,2012, Methods for estimating selected low-flow frequency statistics and harmonic mean flows for streams in Iowa: U.S. Geological Survey Scientific Investigations Report 2012-5171, 99 p. (http://pubs.usgs.gov/sir/2012/5171/)

Linhart, S.M., Nania, J.F., Sanders, C.L., Jr., and Archfield, S.A., 2012, Computing daily mean streamflow at ungaged locations in lowa by using the Flow Anywhere and Flow

Duration Curve Transfer statistical methods: U.S. Geological Survey Scientific Investigations Report 2012–5232, 50 p. (http://pubs.usgs.gov/sir/2012/5232/) Eash, D.A., Barnes, K.K., and Veilleux, A.G.,2013, Methods for estimating annual exceedance-probability discharges for streams in Iowa, based on data through water year 2010: U.S. Geological Survey Scientific Investigations Report 2013-5086, 63 p. with a (http://pubs.usgs.gov/sir/2013/5086/)

Eash, D.A.,2015, Comparisons of estimates of annual exceedance-probability discharges for small drainage basins in Iowa, based on data through water year 2013: U.S. Geological Survey Scientific Investigations Report 2015–5055, 37 p.

(http://dx.doi.org/10.3133/sir20155055.)

Eash, D.A., Barnes, K.K., and O'Shea, P.S.,2016, Methods for estimating selected spring and fall low-flow frequency statistics for ungaged stream sites in lowa, based on data through June 2014: U.S. Geological Survey Scientific Investigations Report 2016–5111, 32 p. (http://dx.doi.org/10.3133/sir20165111)

Perry, C.A., Wolock, D.M., and Artman, J.C.,2004, Estimates of Flow Duration, Mean Flow, and Peak-Discharge Frequency Values for Kansas Stream Locations: U.S. Geological Survey Scientific Investigations Report 2004-5033, 651 p.

(http://water.usgs.gov/pubs/sir/2004/5033/pdf/sir2004.5033front.pdf)

Painter, C.C., Heimann, D.C., and Lanning-Rush, J.L.,2017, Methods for estimating annual exceedance-probability streamflows for streams in Kansas based on data through water year 2015: U.S. Geological Survey Scientific Investigations Report 2017–5063, 20 p. (https://doi.org/10.3133/sir20175063)

Hodgkins, G.A. and Martin, G.R.,2003, Estimating the Magnitude of Peak Flows for Streams in Kentucky for Selected Recurrence Intervals: U.S. Geological Survey Water-Resources Investigations Report 03-4180, 69 p. (http://water.usgs.gov/pubs/wri/wri034180/)
Martin, G.R., Ruhl, K.J., Moore, B.L., and Rose, M.F.,1997, Estimation of Peak-Discharge Frequency of Urban Streams in Jefferson County, Kentucky: U.S. Geological Survey Water-Resources Investigations Report 97-4219 (http://pubs.er.usgs.gov/publication/wri974219)
Martin, G.R.,2002, Estimating Mean Annual Streamflow of Rural Streams in Kentucky: U.S. Geological Survey Water-Resources Investigations Report 02-4206, 35 p. (http://pubs.er.usgs.gov/publication/wri024206)

Martin, G.R., and Arihood, L.D.,2010, Methods for estimating selected low-flow frequency statistics for unregulated streams in Kentucky: U.S. Geological Survey Scientific Investigations Report 2010-5217, 83 p. (http://pubs.usgs.gov/sir/2010/5217/)

Martin, G. R. and Ruhl, K. J.,1993, Regionalization of harmonic-mean streamflows in Kentucky: U.S. Geological Survey Water-Resources Investigations Report 92-4173, 47 p., 1 pl. (http://pubs.er.usgs.gov/publication/wri924173StreamStats_KY_20140226.mdb) Brockman, R. A., Agouridis, C. T., Workman, S. R., Ormsbee, L. E., Fogle, A. W.,2012, Bankfull regional curves for the Inner and Outer Bluegrass Regions of Kentucky, Journal of the American Water Resources Association, v. 48, no. 2, p. 391-406.

(http://onlinelibrary.wiley.com/doi/10.1111/j.1752-1688.2011.00621.x/full)

TR No.70, (2004) Regionalized Regression Equations for Estimating Low-Flow Characteristics for selected Louisiana Streams

(http://la.water.usgs.gov/publications/pdfs/TR70.pdf)

TR No.60, (1998) Floods in Louisiana, Magnitude and Frequency, Fifth Edition (not available)

Landers, M.N.,1985, Floodflow Frequency of Streams in the Alluvial Plain of the Lower Mississippi River in Mississippi, Arkansas, and Louisiana: U.S. Geological Survey Water-

```
Resources Investigations Report 85-4150, 21 p.
```

(http://pubs.er.usgs.gov/publication/wri854150)

Lombard, P. J., Tasker, G. D., and Nielsen, M. G.,2003, August Median Streamflow on Ungaged Streams in Eastern Aroostook County, Maine: U.S. Geological Survey Water-Resources Investigations Report 03-4225, 20 p.

(http://water.usgs.gov/pubs/wri/wri034225/pdf/wrir03-4225.pdf)

Lombard, P. J.,2004, August Median Streamflow on Ungaged Streams in Eastern Coastal Maine: U.S. Geological Survey Scientific Investigations Report 2004-5157, 15 p. (http://water.usgs.gov/pubs/sir/2004/5157/)

Dudley, R.W.,2004, Estimating Monthly, Annual, and Low 7-Day, 10-Year Streamflows for Ungaged Rivers in Maine: U.S. Geological Survey Scientific Investigations Report 2004-5026, 22 p. (http://water.usgs.gov/pubs/sir/2004/5026/pdf/sir2004-5026.pdf)

Hodgkins, G. A.,1999, Estimating the Magnitude of Peak Flows for Streams in Maine for Selected Recurrence Intervals: U.S. Geological Survey Water-Resources Investigations Report 99-4008, 45 p. (https://pubs.er.usgs.gov/publication/wri994008)

Dudley, R.W.,2004, Hydraulic-Geometry Relations for Rivers in Coastal and Central Maine: U.S. Geological Survey Scientific Investigations Report 2004-5042, 30 p (http://pubs.usgs.gov/sir/2004/5042/pdf/sir2004-5042.pdf)

Lombard, P.J.,2010, June and August median streamflows estimated for ungaged streams in southern Maine: U.S. Geological Survey Scientific Investigations Report 2010-5179, 16 p. (http://pubs.usgs.gov/sir/2010/5179/pdf/sir2010-5179.pdf)

Lombard, P.J., and Hodgkins, G.A.,2015, Peak flow regression equations for small, ungaged streams in Maine— Comparing map-based to field-based variables: U.S. Geological Survey Scientific Investigations Report 2015–5049, 12 p. (http://dx.doi.org/10.3133/sir20155049) Dudley, R.W.,2015, Regression equations for monthly and annual mean and selected percentile streamflows for ungaged rivers in Maine: U.S. Geological Survey Scientific Investigations Report 2015–5151, 35 p. (http://dx.doi.org/10.3133/sir20155151) Thomas, Jr., W.O. and Moglen, G.E.,2010, An Update of Regional Regression Equations for Maryland, Appendix 3 in Application of Hydrologic Methods in Maryland, Third Edition, September 2010: Maryland State Highway Administration and Maryland Department of the

(http://gishydro.eng.umd.edu/HydroPanel/hydrology_panel_report_3rd_edition_final.pdf)
Chaplin, J.J.,2005, Development of regional curves relating bankfull-channel geometry and
discharge to drainage area for streams in Pennsylvania and selected areas of Maryland:
U.S. Geological Survey Scientific Investigations Report 2005-5147, 34 p.

(https://pubs.usgs.gov/sir/2005/5147/SIR2005-5147.pdf)

Environment, 38 p.

Ries, K.G., III,2000, Methods for estimating low-flow statistics for Massachusetts streams: U.S. Geological Survey Water Resources Investigations Report 00-4135, 81 p. (http://pubs.usgs.gov/wri/wri004135/)

Bent, G.C., and Steeves, P.A.,2006, A revised logistic regression equation and an automated procedure for mapping the probability of a stream flowing perennially in Massachusetts: U.S. Geological Survey Scientific Investigations Report 2006-5031, 107 p.

(http://pubs.usgs.gov/sir/2006/5031/pdfs/SIR_2006-5031rev.pdf)

Bent, G.C., and Waite, A.M.,2013, Equations for estimating bankfull channel geometry and discharge for streams in Massachusetts: U.S. Geological Survey Scientific Investigations Report 2013–5155, 62 p., (http://pubs.usgs.gov/sir/2013/5155/)

Zarriello, P.J.,2017, Magnitude of flood flows at selected annual exceedance probabilities for streams in Massachusetts: U.S. Geological Survey Scientific Investigations Report 2016-5156, 99 p. (https://dx.doi.org/10.3133/sir20165156)

Holtschlag, D.J. and Croskey, H.M.,1984, Statistical Methods for Estimating Flow Characteristics of Michigan Streams: U.S. Geological Survey Water-Resources Investigations Report 84-4207, 80 p. (http://pubs.er.usgs.gov/usgspubs/wri/wri844207) Lorenz, D.L., Sanocki, C.A., and Kocian, M.J.,2009, Techniques for Estimating the Magnitude and Frequency of Peak Flows on Small Streams in Minnesota Based on Data through Water Year 2005: U.S. Geological Survey Scientific Investigations Report 2009-5250, 54 p. (http://pubs.usgs.gov/sir/2009/5250/pdf/sir2009-5250.pdf) Ziegeweid, J.R., Lorenz, D.L., Sanocki, C.A., and Czuba, C.R.,2015, Methods for estimating

Ziegeweid, J.R., Lorenz, D.L., Sanocki, C.A., and Czuba, C.R.,2015, Methods for estimating flow-duration curve and low-flow frequency statistics for ungaged locations on small streams in Minnesota: U.S. Geological Survey Scientific Investigations Report 2015–5170, 23 p. (http://dx.doi.org/10.3133/sir20155170)

Anderson, B.T.,2018, Flood frequency of rural streams in Mississippi, 2013: U.S. Geological Survey Scientific Investigations Report 2018–5148, 12 p.

(https://doi.org/10.3133/sir20185148)

Southard, R.E., and Veilleux, A.G.,2014, Methods for estimating annual exceedance-probability discharges and largest recorded floods for unregulated streams in rural Missouri: U.S. Geological Survey Scientific Investigations Report 2014–5165, 39 p. (http://pubs.usgs.gov/sir/2014/5165/)

Southard, R.E.,2013, Computed statistics at streamgages, and methods for estimating low-flow frequency statistics and development of regional regression equations for estimating low-flow frequency statistics at ungaged locations in Missouri: U.S. Geological Survey Scientific Investigations Report 2013–5090, 28 p. (http://pubs.usgs.gov/sir/2013/5090/) Parrett, Charles and Hull, J.A.,1985, A method for estimating mean and low flows of streams in national forests of Montana: U.S. Geological Survey Water-Resources Investigations Report 85-4071, 13 p. (https://pubs.er.usgs.gov/publication/wri854071) Parrett, Charles and Cartier, K.D. ,1999, Methods for estimating monthly streamflow characteristics at ungaged sites in western Montana: U. S. Geological Survey Water-Supply Paper 2365, 30 p. (http://pubs.er.usgs.gov/publication/wsp2365)

Parrett, Charles and Johnson, D.R.,2004, Methods for Estimating Flood Frequency in Montana Based on Data through Water Year 1998: U.S. Geological Survey Water-Resources Investigations Report 03-4308, 102 p. (http://water.usgs.gov/pubs/wri/wri03-4308/) Sando, Roy, Sando, S.K., McCarthy, P.M., and Dutton, D.M.,2016, Methods for estimating peak-flow frequencies at ungaged sites in Montana based on data through water year 2011: U.S. Geological Survey Scientific Investigations Report 2015-5019-F, 30 p. (https://doi.org/10.3133/sir20155019)

McCarthy, P.M., Sando, Roy, Sando, S.K., and Dutton, D.M.,2016, Methods for estimating streamflow characteristics at ungaged sites in western Montana based on data through water year 2009: U.S. Geological Survey Scientific Investigations Report 2015–5019–G, 19 p. (https://doi.org/10.3133/sir20155019)

Soenksen, P.J., Miller, L.D., Sharpe, J.B. and Watton, J.R.,1999, Peak-Flow Frequency Relations and Evaluation of the Peak-Flow Gaging Network in Nebraska: U. S. Geological Survey Water-Resources Investigations Report 99-4032, 48 p,

(https://pubs.er.usgs.gov/publication/wri994032)

Flynn, R.H. and Tasker, G.D.,2002, Development of Regression Equations to Estimate Flow Durations and Low-Flow-Frequency Statistics in New Hampshire Streams: U.S.Geological Survey Scientific Investigations Report 02-4298, 66 p. (http://pubs.water.usgs.gov/wrir02-4298)

Olson, S.A., 2009, Estimation of flood discharges at selected recurrence intervals for streams in New Hampshire: U.S.Geological Survey Scientific Investigations Report 2008-

5206, 57 p. (http://pubs.usgs.gov/sir/2008/5206/)

Flynn, R.H. and Tasker, G.D.,2004, Generalized Estimates from Streamflow Data of Annual and Seasonal Ground-Water-Recharge Rates for Drainage Basins in New Hampshire, U.S. Geological Survey Scientific Investigations Report 2004-5019, 67 p.

(http://pubs.usgs.gov/sir/2004/5019/http://pubs.usgs.gov/sir/2004/5019/)

Watson, K.M., and Schopp, R.D., 2009, Methodology for estimation of flood magnitude and frequency for New Jersey streams, U.S. Geological Survey Scientific Investigations Report 2009-5167, 51 p. (http://pubs.usgs.gov/sir/2009/5167/)

Watson, K.M., and McHugh, A.R.,2014, Regional regression equations for the estimation of selected monthly low-flow duration and frequency statistics at ungaged sites on streams in New Jersey: U.S. Geological Survey Scientific Investigations Report 2014-5004, 59 p. (baseline, period-or-record statistics)

(http://dx.doi.org/10.3133/sir20145004StreamStatsDB\2019_12_13_DataSource_table.xlsxDa Waltemeyer, S.D.,2002, Analysis of the magnitude and frequency of the 4-day, 3-year annual low flow on unregulated streams in New Mexico: U. S. Geological Survey Water-Resources Investigations Report 01-4271, 22 p.

(http://nm.water.usgs.gov/publications/abstracts/wrir01-4271.pdf)

Waltemeyer, S.D., 2008, Analysis of the Magnitude and Frequency of Peak Discharge and Maximum Observed Peak Discharge in New Mexico and Surrounding Areas: U.S. Geological Survey Scientific Investigations Report 2008-5119, 105 p.

(http://pubs.usgs.gov/sir/2008/5119/)

Lumia, Richard, Freehafer, D.A., and Smith, M.J.,2006, Magnitude and Frequency of Floods in New York: U.S. Geological Survey Scientific Investigations Report 2006–5112, 152 p. (http://pubs.usgs.gov/sir/2006/5112/)

Stedfast, D.A.,1984, Evaluation of Six Methods for Estimating Magnitude and Frequency of Peak Discharges on Urban Streams in New York: U. S. Geological Survey Water-Resources Investigations Report 84-4350, 24 p.

(http://onlinepubs.er.usgs.gov/djvu/WRI/wrir_84_4350.djvu)

Mulvihill, C.I., Baldigo, B.P., Miller, S.J., and DeKoskie, Douglas,2009, Bankfull Discharge and Channel Characteristics of Streams in New York State: U.S. Geological Survey Scientific Investigations Report 2009-5144, 51 p. (http://pubs.usgs.gov/sir/2009/5144/) Barnes, C. R.,1986, Method for estimating low-flow statistics for ungaged streams in the lower Hudson River Basin, New York: U. S. Geological Survey Water-Resources Investigations Report 85-4070, 22 p.

(http://onlinepubs.er.usgs.gov/djvu/WRI/wrir_85_4070.djvu)

Randall, A.D.,2010, Low flow of streams in the Susquehanna River basin of New York: U.S. Geological Survey Scientific Investigations Report 2010-5063, 57 p.

(http://pubs.usgs.gov/sir/2010/5063/http://pubs.usgs.gov/sir/2010/5063/)

Gazoorian, C.L.,2015, Estimation of unaltered daily mean streamflow at ungaged streams of New York, excluding Long Island, water years 1961–2010: U.S. Geological Survey Scientific Investigations Report 2014–5220, 29 p. (https://pubs.usgs.gov/sir/2014/5220/) Giese, G. L. and Mason, R.R., Jr.,1993, Low-flow characteristics of streams in North Carolina: U.S. Geological Survey Water-Supply Paper 2403, 29 p.

(https://pubs.er.usgs.gov/publication/wsp2403)

Mason, Robert R., Jr.; Fuste, Luis A.; King, Jeffrey N.; Thomas, Wilbert O., Jr.,2002, The National Flood-Frequency Program -- Methods for Estimating Flood Magnitude and Frequency in Rural and Urban Areas in North Carolina, 2001: U.S. Geological Survey Fact Sheet 007-00, 4 p. (http://pubs.er.usgs.gov/publication/fs00700)

Weaver, J.C., Feaster, T.D., and Gotvald, A.J.,2009, Magnitude and frequency of rural floods in the Southeastern United States, through 2006—Volume 2, North Carolina: U.S. Geological Survey Scientific Investigations Report 2009–5158, 111 p.

(http://pubs.usgs.gov/sir/2009/5158/)

Williams-Sether, T.,2015, Regional regression equations to estimate peak-flow frequency at sites in North Dakota using data through 2009: U.S. Geological Survey Scientific Investigations Report 2015-5096, 12 p. (http://dx.doi.org/10.3133/sir20155096)

Koltun, G.F., Kula, S.P., and Puskas, B.M., 2006, A Streamflow Statistics (StreamStats) Web Application for Ohio: U.S. Geological Survey Scientific Investigations Report 2006-5312, 62 p. (http://pubs.usgs.gov/sir/2006/5312/)

Sherwood, J.M.,1994, Estimation of peak-frequency relations, flood hydrographs, and volume-duration-frequency relations of ungaged small urban streams in Ohio: U. S. Geological Survey Water-Supply Paper 2432, 42 p.

(http://pubs.er.usgs.gov/usgspubs/wsp/wsp2432)

Koltun, G. F., and Whitehead, M. T.,2002, Techniques for Estimating Selected Streamflow Characteristics of Rural, Unregulated Streams in Ohio: U. S. Geological Survey Water-Resources Investigations Report 02-4068, 50 p

(https://pubs.er.usgs.gov/publication/wri024068)

Koltun, G. F., and Schwartz, Ronald R.,1987, MULTIPLE-REGRESSION EQUATIONS FOR ESTIMATING LOW FLOWS AT UNGAGED STREAM SITES IN OHIO: U.S. Geological Survey Water-Resources Investigations Report 86-4354, 39 p.

(http://pubs.er.usgs.gov/usgspubs/wri/wri864354)

Koltun, G.F., and Kula, S.P.,2013, Methods for estimating selected low-flow statistics and development of annual flow-duration statistics for Ohio: U.S. Geological Survey Scientific Investigations Report 2012–5138, 195 p. (http://pubs.usgs.gov/sir/2012/5138/)

Koltun, G.F.,2019, Flood-frequency estimates for Ohio streamgages based on data through water year 2015 and techniques for estimating flood-frequency characteristics of rural, unregulated Ohio streams: U.S. Geological Survey Scientific Investigations Report 2019–5018, xx p. (https://dx.doi.org/10.3133/sir20195018)

Esralew, R.A., Smith, S.J.,2009, Methods for estimating flow-duration and annual meanflow statistics for ungaged streams in Oklahoma: U.S. Geological Survey Scientific Investigations Report 2009-5267, 131 p. (http://pubs.usgs.gov/sir/2009/5267/)

Smith, S.J., Lewis, J.M., and Graves, G.M.,2015, Methods for estimating the magnitude and frequency of peak streamflows at ungaged sites in and near the Oklahoma Panhandle: U.S. Geological Survey Scientific Investigations Report 2015-5134, 35 p.

(http://dx.doi.org/10.3133/sir20155134)

Lewis, J.M., Hunter, S.L., and Labriola, L.G.,2019, Methods for estimating the magnitude and frequency of peak streamflows for unregulated streams in Oklahoma developed by using streamflow data through 2017: U.S. Geological Survey Scientific Investigations Report 2019-5143, 39 p. (https://doi.org/10.3133/sir20195143)

Laenen, Antonius,1980, Storm Runoff As Related to Urbanization in the Portland, Oregon - Vancouver, Washington Area: U.S. Geological Survey Open-File Report 80-689, 71 p. (http://pubs.er.usgs.gov/usgspubs/wri/wri834143)

Cooper, R.M.,2005, Estimation of Peak Discharges for Rural, Unregulated Streams in Western Oregon: U.S. Geological Survey Scientific Investigations Report 2005-5116, 76 p. (http://pubs.usgs.gov/sir/2005/5116/pdf/sir2005-5116.pdf)

Risley, John, Stonewall, Adam, and Haluska, Tana, 2008, Estimating flow-duration and low-flow frequency statistics for unregulated streams in Oregon: U.S. Geological Survey Scientific Investigations Report 2008-5126, 22 p. (http://pubs.usgs.gov/sir/2008/5126/)

Cooper, Richard, 2006, Estimation of Peak Discharges for Rural, Unregulated Streams in Eastern Oregon, Oregon Water Resources Department OFR SW 06-001, Salem, OR. (https://digital.osl.state.or.us/islandora/object/osl%3A14736/datastream/OBJ/view) Stuckey, M.H., 2006, Low-flow, base-flow, and mean-flow regression equations for Pennsylvania streams: U.S. Geological Survey Scientific Investigations Report 2006-5130, 84 p. (http://pubs.usgs.gov/sir/2006/5130/)

Stuckey, M.H., Koerkle, E.H., and Ulrich, J.E.,2012, Estimation of baseline daily mean streamflows for ungaged locations on Pennsylvania streams, water years 1960–2008: U.S. Geological Survey Scientific Investigations Report 2012–5142, 61 p.

(http://pubs.usgs.gov/sir/2012/5142/)

Clune, J.W., Chaplin, J.J., and White, K.E.,2018, Comparison of regression relations of bankfull discharge and channel geometry for the glaciated and nonglaciated settings of Pennsylvania and southern New York: U.S. Geological Survey Scientific Investigations Report 2018-5066, 20 p. (https://doi.org/10.3133/sir20185066)

Roland, M.A., and Stuckey, M.H.,2008, Regression equations for estimating flood flows at selected recurrence intervals for ungaged streams in Pennsylvania: U.S. Geological Survey Scientific Investigations Report 2008-5102, 57p. (http://pubs.usgs.gov/sir/2008/5102/) Zarriello, P.J., Ahearn, E.A., and Levin, S.B.,2012, Magnitude of flood flows for selected annual exceedance probabilities in Rhode Island through 2010: U.S. Geological Survey Scientific Investigations Report 2012-5109, 93 p. (http://pubs.usgs.gov/sir/2012/5109) Bent, G.C., Steeves, P.A., and Waite, A.M.,2014, Equations for estimating selected streamflow statistics in Rhode Island: U.S. Geological Survey Scientific Investigations Report 2014-5010, 65 p. (http://dx.doi.org/10.3133/sir20145010)

Feaster, T.D., Gotvald, A.J., and Weaver, J.C.,2009, Magnitude and Frequency of Rural Floods in the Southeastern United States, 2006: Volume 3, South Carolina: U.S. Geological Survey Scientific Investigations Report 2009-5156, 226 p.

(http://pubs.usgs.gov/sir/2009/5156/)

Sando, Steven K.,1998, A Method for Estimating Magnitude and Frequency of Floods in South Dakota: U.S. Geological Survey Water-Resources Investigations Report 98-4055, 48 p. (http://pubs.water.usgs.gov/wri98-4055/)

Law, G.S., and Tasker G.D.,2003, Flood-Frequency Prediction Methods for Unregulated Streams of Tennessee, 2000: U.S. Geological Survey Water-Resources Investigations Report 03-4176, 79p. (http://pubs.usgs.gov/wri/wri034176/)

Neely, B.L., Jr.,1984, Flood Frequency and Storm Runoff of Urban Areas of Memphis and Shelby County, Tennessee: U.S. Geological Survey Water-Resources Investigations Report 84-4110, 51 p. (http://pubs.usgs.gov/wri/wrir_84-4110/)

Robbins, Clarence H.,1984, Synthesized Flood Frequency of Small Urban Streams in Tennessee: U.S. Geological Survey Water-Resources Investigations Report 84-4182, 24 p. (http://onlinepubs.er.usgs.gov/djvu/WRI/wrir_84_4182.djvu)

Law, G.S., Tasker, G.D., and Ladd, D.E.,2009, Streamflow-characteristic estimation methods for unregulated streams of Tennessee: U.S. Geological Survey Scientific Investigations Report 2009–5159, 212 p., 1 pl. (http://pubs.usgs.gov/sir/2009/5159/)

Asquith, W.H., Slade, R.M., Jr.,1999, Site-specific estimation of peak-stream flow frequency using generalized least squares regression for natural basins in Texas: U.S. Geological Survey Water-Resources Investigations Report 99-4172, 19 p.

(http://pubs.water.usgs.gov/wri994172)

Asquith, William H.,1998, Peak-flow frequency for tributaries of the Colorado River downstream of Austin, Texas U.S. Geological Survey Water-Resources Investigations Report 98-4015, 26 p. (http://pubs.water.usgs.gov/wri98-4015/)

```
Raines, Timothy H., 1998, Peak-discharge frequency and potential extreme peak discharge
for natural streams in the Brazos River basin, Texas: U.S. Geological Survey Water-
Resources Investigations Report 98-4178, 47 p., 1 plate (http://pubs.water.usgs.gov/wri98-
4178/)
Land, L.F., Schroeder, E.E. and Hampton, B.B., 1982, Techniques for Estimating the
Magnitude and Frequency of Floods in the Dallas-Fort Worth Metropolitan Area, Texas: U.S.
Geological Survey Water-Resources Investigations Report 82-18, 55 p. ()
Asquith, W.H., Slade, R. M., Lanning-Rush, Jennifer, 1996, Peak-flow frequency and extreme
flood potential for streams in the vicinity of the Highland Lakes, central Texas: U.S.
Geological Survey Water-Resources Investigations Report 96-4072
(http://onlinepubs.er.usgs.gov/djvu/WRI/wrir_96_4072_plt.djvu)
Liscum, Fred and Massey, B.C., 1980, Technique for Estimiating the Magnitude and
Frequency of Floods in the Houston, Texas, Metropolitan Area: U.S. Geological Survey
Water-Resources Investigations Report 80-17, 29 p. ()
Asquith, W.H., and Roussel, M.C., 2009, Regression equations for estimation of annual
peak-streamflow frequency for undeveloped watersheds in Texas using an L-moment-
based, PRESS-minimized, residual-adjusted approach: U.S. Geological Survey Scientific
Investigations Report 2009-5087, 48 p. (http://pubs.usgs.gov/sir/2009/5087/)
Kenney, T.A., Wilkowske, C.D., and Wright, S.J., 2007, Methods for Estimating Magnitude
and Frequency of Peak Flows for Natural Streams in Utah: U.S. Geological Survey Scientific
Investigations Report 2007-5158, 28 p. (http://pubs.usgs.gov/sir/2007/5158/)
Wilkowske, C.D., Kenney, T.A., and Wright, S.J., 2009, Methods for Estimating Monthly and
Annual Streamflow Statistics at Ungaged Sites in Utah: U.S. Geological Survey Scientific
Investigations Report 2008-5230, 62 p. (http://pubs.usgs.gov/sir/2008/5230/)
Olson, S.A., 2002, Flow-frequency characteristics of Vermont streams: U.S. Geological
Survey Water-Resources Investigations Report 02-4238, 47 p.
(http://pubs.usgs.gov/wri/wrir02-4238/)
Olson, S.A., 2014, Estimation of flood discharges at selected annual exceedance
probabilities for unregulated, rural streams in Vermont, with a section on Vermont regional
skew regression, by Veilleux, A.G.: U.S. Geological Survey Scientific Investigations Report
2014-5078, 27 p. plus appendixes. (http://pubs.usgs.gov/sir/2014/5078/)
Olson, S.A., and Brouillette, M.C., 2006, A logistic regression equation for estimating the
probability of a stream in Vermont having intermittent flow: U.S. Geological Survey
Scientific Investigations Report 2006-5217, 15 p. (https://pubs.usgs.gov/sir/2006/5217/)
Austin, S.H., Krstolic, J.L., and Wiegand, Ute, 2011, Low-flow characteristics of Virginia
streams: U.S. Geological Survey Scientific Investigations Report 2011-5143, 122 p. + 9
tables on CD. (http://pubs.usgs.gov/sir/2011/5143/)
Austin, S.H., Krstolic, J.L., and Wiegand, Ute, 2011, Peak-flow characteristics of Virginia
streams: U.S. Geological Survey Scientific Investigations Report 2011-5144, 106 p. + 3
tables and 2 appendixes on CD. (http://pubs.usgs.gov/sir/2011/5144/)
Austin, S.H., 2014, Methods and equations for estimating peak streamflow per square mile
in Virginia's urban basins: U.S. Geological Survey Scientific Investigations Report 2014-
5090, 25 p. (http://pubs.usgs.gov/sir/2014/5090/http://pubs.usgs.gov/sir/2014/5090/)
Curran, C.A. and Olsen, T.D., 2009, Estimating Low-Flow Frequency Statistics and
Hydrologic Analysis of Selected Streamflow-Gaging Stations, Nooksack River Basin,
Northwestern Washington and Canada: U.S. Geological Survey Scientific Investigations
Report 2009-5170, 44 p. (http://pubs.usgs.gov/sir/2009/5170/)
Curran, C.A., Eng, Ken, and Konrad, C.P., 2012, Analysis of low flows and selected methods
```

for estimating low-flow characteristics at partial-record and ungaged stream sites in

western Washington: U.S. Geological Survey Scientific Investigations Report 2012-5078, 46 p. (http://pubs.usgs.gov/sir/2012/5078/)

Mastin, M.C., Konrad, C.P., Veilleux, A.G., and Tecca, A.E., 2016, Magnitude, frequency, and trends of floods at gaged and ungaged sites in Washington, based on data through water year 2014 (ver 1.1, October 2016): U.S. Geological Survey Scientific Investigations Report 2016–5118, 70 p. (http://dx.doi.org/10.3133/sir20165118)

Wiley, Jeffrey B.,2008, Estimating Selected Streamflow Statistics Representative of 1930–2002 in West Virginia: U.S. Geological Survey Scientific Investigations Report 2008-5105, 24 p. (http://pubs.usgs.gov/sir/2008/5105/)

Wiley, Jeffrey B.,1987, Techniques for estimating flood depth frequency relations for streams in West Virginia: U.S. Geological Survey Water-Resources Investigations Report 87-4111, 17 p. (http://pubs.er.usgs.gov/usgspubs/wri/wri874111)

Wiley, J.B., and Atkins, J.T., Jr.,2010, Estimation of flood-frequency discharges for rural, unregulated streams in West Virginia: U.S. Geological Survey Scientific Investigations Report 2010-5033, 78 p. (http://pubs.usgs.gov/sir/2010/5033/)

Wiley, J.B., and Atkins, J.T., Jr.,2010, Estimation of selected seasonal streamflow statistics representative of 1930-2002 in West Virginia: U.S. Geological Survey Scientific Investigations Report 2010-5185, 20 p. (http://pubs.usgs.gov/sir/2010/5185/)

Conger, Duane H.,1986, Estimating Magnitude and Frequency of Floods for Wisconsin Urban Streams: U.S. Geological Survey Water-Resources Investigations Report 86-4005, 18 p. (http://pubs.er.usgs.gov/publication/wri864005)

Walker, J.F., Peppler, M.C., Danz, M.E., and Hubbard, L.E., 2017, Flood-frequency characteristics of Wisconsin streams (ver. 2.1, December 2017): Reston, Virginia, U.S. Geological Survey Scientific Investigations Report 2016–5140, 33 p., 1 plate, 2 appendixes (https://doi.org/10.3133/sir20165140)

Miller, Kirk A., 2003, Peak-flow Characteristics of Wyoming Streams: U.S. Geological Survey Water-Resources Investigations Report 03-4107, 79 p.

(http://pubs.usgs.gov/wri/wri034107/)

Ramos-Ginés, Orlando,1999, Estimation of Magnitude and Frequency of Floods for Streams in Puerto Rico: New Empirical Models: U. S. Geological Survey Water-Resources Investigations Report 99-4142, 41 p. (http://pubs.usgs.gov/wri/wri994142/)

Moody, J.A.,2012, An analytical method for predicting postwildfire peak discharges: U.S. Geological Survey Scientific Investigations Report 2011-5236, 36 p.

(https://pubs.usgs.gov/sir/2011/5236/)

August Flow-Duration Statistics Parameters[Statewide Low Flow WRIR00 4135]					
Parameter Code	Parameter Name	Value	Units	Min Limit	Max Limit
DRNAREA	Drainage Area	0.0906	square miles	1.61	149
BSLDEM250	Mean Basin Slope from 250K DEM	4.012	percent	0.32	24.6
DRFTPERSTR	Stratified Drift per Stream Length	-100000	square mile per mile	0	1.29
MAREGION	Massachusetts Region	1	dimensionless	0	1

Statistic Value Unit

August Flow-Duration Statistics Citations

Sauer, Vernon B.; Thomas, W. O., Jr.; Stricker, V. A.; Wilson, K. V.,1983, Flood characteristics of urban watersheds in the United States: U.S. Geological Survey Water-Supply Paper 2207, 63 p. (http://pubs.er.usgs.gov/publication/wsp2207)
()

Anderson, B.T.,2020, Magnitude and frequency of floods in Alabama, 2015: U.S. Geological Survey Scientific Investigations Report 2020–5032, 148 p.

(https://doi.org/10.3133/sir20205032)

Hedgecock, T.S.,2004, Magnitude and Frequency of Floods on Small Rural Streams in Alabama: U. S. Geological Survey Scientific Investigations Report 2004-5135, 10 p. (http://pubs.usgs.gov/sir/2004/5135/)

Hedgecock, T.S.,2010, Magnitude and Frequency of Floods for Urban Streams in Alabama, 2007: U.S Geological Survey Scientific Investigations Report 2010-5012, 17p. (https://pubs.usgs.gov/sir/2010/5012/)

Wiley, J.B., and Curran, J.H.,2003, Estimating annual high-flow statistics and monthly and seasonal low-flow statistics for ungaged sites on streams in Alaska and conterminous basins in Canada: U.S. Geological Survey Water-Resources Investigations Report 03-4114, 61 p. (http://water.usgs.gov/pubs/wri/wri034114/pdf/wri034114_v1.10.pdf)
Brabets, Timothy P.,1996, Evaluation of the streamflow-gaging network of Alaska in providing regional streamflow information: U.S. Geological Survey Water-Resources Investigations Report 96-4001, 98 p. (http://pubs.er.usgs.gov/usgspubs/wri/wri964001)
Curran, J.H., Barth, N.A., Veilleux, A.G., and Ourso, R.T.,2016, Estimating Flood Magnitude

and Frequency at Gaged and Ungaged Sites on Streams in Alaska and Conterminous Basins in Canada, Based on Data through Water Year 2012: U.S. Geological Survey Scientific Investigations Report 2016-5024, 47 p.

(http://dx.doi.org/10.3133/sir20165024http://dx.doi.org/10.3133/sir20165024)
Southard, R.E.,2010, Estimation of the Magnituude and Frequency of Floods in Urban
Basins in Missouri: U.S. Geological Survey Scientific Investigations Report 2010-5073, 27
p. (http://pubs.usgs.gov/sir/2010/5073/)

Waltemeyer, S.D., Analysis of the Magnitude and Frequency of Peak Discharges for the Navajo Nation in Arizona, Utah, Colorado, and New Mexico: U. S. Geological Survey Scientific Investigations Report2006-5306, 42 p. (http://pubs.usgs.gov/sir/2006/5306/) Paretti, N.V., Kennedy, J.R., Turney, L.A., and Veilleux, A.G.,2014, Methods for estimating magnitude and frequency of floods in Arizona, developed with unregulated and rural peakflow data through water year 2010: U.S. Geological Survey Scientific Investigations Report 2014-5211, 61 p., http://dx.doi.org/10.3133/sir20145211.

(http://pubs.usgs.gov/sir/2014/5211/)

Kennedy, J.R., Paretti, N.V., and Veilleux, A.G.,2014, Methods for estimating magnitude and frequency of 1-, 3-, 7-, 15-, and 30-day flood-duration flows in Arizona: U.S. Geological Survey Scientific Investigations Report 2014–5109, 35 p.

(http://pubs.usgs.gov/sir/2014/5109/)

Funkhouser, J.E., Eng, Ken, and Moix, M.W.,2008, Low-Flow Characteristics and Regionalization of Low Flow Characteristics for Selected Streams in Arkansas: U. S.

Geological Survey Scientific Investigations Report 2008-5065, 161 p. (http://pubs.usgs.gov/sir/2008/5065/pdf/SIR2008-5065.pdf)

Breaker, B.K.,2015, Dry season mean monthly flow and harmonic mean flow regression equations for selected ungaged basins in Arkansas: U.S. Geological Survey Scientific Investigations Report 2015–5031, 25 p. (http://pubs.usgs.gov/sir/2015/5031/) Wagner, D.M., Krieger, J.D., and Veilleux, A.G.,2016, Methods for estimating annual exceedance probability discharges for streams in Arkansas, based on data through water year 2013: U.S. Geological Survey Scientific Investigations Report 2016–5081, 136 p. (http://dx.doi.org/10.3133/sir20165081)

Thomas, B.E, Hjalmarson, H.W., and Waltemeyer, S.D.,1997, Methods for Estimating Magnitude and Frequency of Floods in the Southwestern United States: U.S. Water-Supply Paper 2433, 196 p. (http://pubs.er.usgs.gov/publication/wsp2433)

Gotvald, A.J., Barth, N.A., Veilleux, A.G., and Parrett, Charles, 2012, Methods for determining magnitude and frequency of floods in California, based on data through water year 2006: U.S. Geological Survey Scientific Investigations Report 2012–5113, 38 p., 1 pl. (http://pubs.usgs.gov/sir/2012/5113/)

Sanocki, C.A., Williams-Sether, T., Steeves, P.A., and Christensen, V.G.,2019, Techniques for Estimating the Magnitude and Frequency of Peak Flows on Small Streams in the Binational U.S. and Canadian Lake of the Woods-Rainy River Basin Upstream from Kenora, Ontario, Canada, Based on Data through Water Year 2013: U.S. Geological Survey Scientific Investigations Report 2019–5012, 17 p. (https://doi.org/10.3133/sir20195012)
Capesius, J.P., and Stephens, V. C.,2009, Regional Regression Equations for Estimation of Natural Streamflow Statistics in Colorado: U. S. Geological Survey Scientific Investigations Report 2009-5136, 32 p.

(http://pubs.usgs.gov/sir/2009/5136/http://pubs.usgs.gov/sir/2009/5136/)

Kohn, M.S., Stevens, M.R., Harden, T.M., Godaire, J.E., Klinger, R.E., and Mommandi, A.,2016, Paleoflood investigations to improve peak-streamflow regional-regression equations for natural streamflow in eastern Colorado, 2015: U.S. Geological Survey Scientific Investigations Report 2016–5099, 58 p. (http://dx.doi.org/10.3133/sir20165099) Ahearn, E.A.,2004, Regression Equations for Estimating Flood Flows for the 2-, 10-, 25-, 50-, 100-, and 500-Year Recurrence Intervals in Connecticut: U.S. Geological Survey SRI 2004-5160, 62 p. (http://water.usgs.gov/pubs/sir/2004/5160/)

Ahearn, E.A.,2010, Regional regression equations to estimate flow-duration statistics in Connecticut: U. S. Geological Survey Scientific Investigations Report 2010-5052, 45 p. (http://pubs.usgs.gov/sir/2010/5052/)

Ries, K.G., III, and Dillow, J.J.A.,2006, Magnitude and frequency of floods in Delaware: Scientific Investigations Report 2006-5146, 59 p. (http://pubs.usgs.gov/sir/2006/5146/) Carpenter, D.H., and Hayes, D.C.,1996, Low-flow characteristics of streams in Maryland and Delaware: U.S. Geological Survey Water-Resources Investigations Report 94-4020, 113 p., 10 plates (http://pubs.er.usgs.gov/usgspubs/wri/wri944020)

Franklin, M.A. and Losey, G.T.,1984, Magnitude and Frequency of Floods from Urban Streams in Leon County, Florida: U.S. Geological Survey Water-Resources Investigations Report 84-4004, 37 p. (http://pubs.er.usgs.gov/publication/wri844004)

Lopez, M.A. and Woodham, W. M.,1983, Magnitude and frequency of flooding on small urban watersheds in the Tampa Bay area, west-central Florida: U.S. Geological Survey Water-Resources Investigations Report 82-42, 52 p.

(https://pubs.er.usgs.gov/publication/wri8242)

Rumenik, R. P.; Grubbs, J. W.,1996, Methods for estimating low-flow characteristics of ungaged streams in selected areas, northern Florida: U.S. Geological Survey Water-

Resources Investigations Report 96-4124, 28 p.

(https://doi.org/10.3133/wri964124https://doi.org/10.3133/wri964124)

Verdi, R.J., and Dixon, J.F.,2011, Magnitude and Frequency of Floods for Rural Streams in Florida, 2006: U.S. Geological Survey Scientific Investigations Report 2011–5034, 69 p., 1 pl. (http://pubs.usgs.gov/sir/2011/5034/)

Inman, E.J.,2000, Lagtime relations for urban streams in Georgia: U.S. Geological Survey Water-Resources Investigations Report 00-4049, 12 p.

(http://ga.water.usgs.gov/pubs/wrir/wrir004049/pdf/wrir00-4049.pdf)

Gotvald, A.J., Feaster, T.D., and Weaver, J.C.,2009, Magnitude and Frequency of Rural Floods in the Southeastern United States, 2006: Volume 1, Georgia: U.S. Geological Survey Scientific Investigations Report 2009-5043, 120 p. (http://pubs.usgs.gov/sir/2009/5043/) Feaster, T.D., Gotvald, A.J., and Weaver, J.C.,2014, Methods for estimating the magnitude and frequency of floods for urban and small, rural streams in Georgia, South Carolina, and North Carolina, 2011 (ver. 1.1, March 2014): U.S. Geological Survey Scientific Investigations Report 2014–5030, 104 p. (http://pubs.usgs.gov/sir/2014/5030/) Gotvald, A.J.,2017, Methods for estimating selected low-flow frequency statistics and mean annual flow for ungaged locations on streams in North Georgia: U.S. Geological Survey Scientific Investigations Report 2017–5001, 25 p.

(https://doi.org/10.3133/sir20175001)

Oki, D.S., Rosa, S.N., and Yeung, C.W.,2010, Flood-frequency estimates for streams on Kaua'i, O'ahu, Moloka'i, Maui, and Hawai'i, State of Hawai'i: U.S. Geological Survey Scientific Investigations Report 2010-5035, 121 p. (http://pubs.usgs.gov/sir/2010/5035/) Gingerich, S.B.,2005, Median and low-flow characteristics for streams under natural and diverted conditions, northeast Maui, Hawaii: U.S. Geological Survey Scientific Investigations Report 2004-5262, 72 p. (http://pubs.usgs.gov/sir/2004/5262/pdf/sir2004-5262.pdf)

Fontaine, R.A., Wong, M.F., Matsuoka, Iwao,1992, Estimation of Median Streamflows at Perennial Stream Sites in Hawaii: U.S. Geological Survey Water-Resources Investigations Report 92-4099, 37 p. (http://pubs.er.usgs.gov/usgspubs/wri/wri924099)

Hortness, J.E.,2006, Estimating Low-Flow Frequency Statistics for Unregulated Streams in Idaho: U.S. Geological Survey Scientific Investigations Report 2006-5035, 31 p. (http://pubs.usgs.gov/sir/2006/5035/pdf/sir20065035.pdf)

Wood, M.S., Fosness, R.L., Skinner, K.D., and Veilleux, A.G.,2016, Estimating peak-flow frequency statistics for selected gaged and ungaged sites in naturally flowing streams and rivers in Idaho: U.S. Geological Survey Scientific Investigations Report 2016–5083, 56 p. (http://dx.doi.org/10.3133/sir20165083)

Hortness, J.E., and Berenbrock, Charles, 2001, Estimating Monthly and Annual Streamflow Statistics at Ungaged Sites in Idaho: U.S. Geological Survey Water-Resources Investigations Report 01–4093, 36 p. (http://idaho.usgs.gov/PDF/wri014093/index.html) Over, T.M., Riley, J.D., Sharpe, J.B., and Arvin, Donald, 2014, Estimation of regional flow-duration curves for Indiana and Illinois: U.S. Geological Survey Scientific Investigations Report 2014–5177, 24 p. and additional downloads, Tables 2–5, 8–13, and 18 (http://dx.doi.org/10.3133/sir20145177)

Soong, D.T., Ishii, A.L., Sharpe, J.B., and Avery, C.F.,2004, Estimating Flood-Peak Discharge Magnitudes and Frequencies for Rural Streams in Illinois, U.S. Geological Survey Scientific Investigations Report 2004-5103. 147 p. (http://il.water.usgs.gov/pubs/sir2004-5103.pdf) Over, T.M., Saito, R.J., Veilleux, A.G., Sharpe, J.B., Soong, D.T., and Ishii, A.L.,2016, Estimation of peak discharge quantiles for selected annual exceedance probabilities in

northeastern Illinois: U.S. Geological Survey Scientific Investigations Report 2016-5050, 50 p. (http://dx.doi.org/10.3133/sir20165050)

Rao, A.R.,2005, Flood-Frequency Relationships for Indiana: Joint Transportation Research Program, Purdue University, FHWA/IN/JTRP-2005/18, 14 p.

(https://docs.lib.purdue.edu/cgi/viewcontent.cgi?article=1746&context=jtrp)

Robinson, B.A.,2013, Regional bankfull-channel dimensions of non-urban wadeable streams in Indiana: U.S. Geological Survey, Scientific Investigations Report 2013-5078, 33 p. (http://pubs.usgs.gov/sir/2013/5078/)

Martin, G.R., Fowler, K.K., and Arihood, L.D.,2016, Estimating selected low-flow frequency statistics and harmonic-mean flows for ungaged, unregulated streams in Indiana (ver 1.1, October 2016): U.S. Geological Survey Scientific Investigations Report 2016-5102, 45 p. (http://dx.doi.org/10.3133/sir20165102)

Arihood, L.D.; Glatfelter, D.R.,1991, Method for estimating low-flow characteristics of ungaged streams in Indiana: U.S. Geological Survey Water-Supply Paper 2372, 19 p. (http://onlinepubs.er.usgs.gov/djvu/WSP/wsp_2372.djvu)

Eash, D.A., and Barnes, K.K.,2012, Methods for estimating selected low-flow frequency statistics and harmonic mean flows for streams in Iowa: U.S. Geological Survey Scientific Investigations Report 2012-5171, 99 p. (http://pubs.usgs.gov/sir/2012/5171/)

Linhart, S.M., Nania, J.F., Sanders, C.L., Jr., and Archfield, S.A.,2012, Computing daily mean streamflow at ungaged locations in Iowa by using the Flow Anywhere and Flow Duration Curve Transfer statistical methods: U.S. Geological Survey Scientific Investigations Report 2012–5232, 50 p. (http://pubs.usgs.gov/sir/2012/5232/)

Eash, D.A., Barnes, K.K., and Veilleux, A.G.,2013, Methods for estimating annual exceedance-probability discharges for streams in Iowa, based on data through water year 2010: U.S. Geological Survey Scientific Investigations Report 2013-5086, 63 p. with a (http://pubs.usgs.gov/sir/2013/5086/)

Eash, D.A.,2015, Comparisons of estimates of annual exceedance-probability discharges for small drainage basins in Iowa, based on data through water year 2013: U.S. Geological Survey Scientific Investigations Report 2015-5055, 37 p.

(http://dx.doi.org/10.3133/sir20155055.)

Eash, D.A., Barnes, K.K., and O'Shea, P.S.,2016, Methods for estimating selected spring and fall low-flow frequency statistics for ungaged stream sites in lowa, based on data through June 2014: U.S. Geological Survey Scientific Investigations Report 2016–5111, 32 p. (http://dx.doi.org/10.3133/sir20165111)

Perry, C.A., Wolock, D.M., and Artman, J.C.,2004, Estimates of Flow Duration, Mean Flow, and Peak-Discharge Frequency Values for Kansas Stream Locations: U.S. Geological Survey Scientific Investigations Report 2004-5033, 651 p.

(http://water.usgs.gov/pubs/sir/2004/5033/pdf/sir2004.5033front.pdf)

Painter, C.C., Heimann, D.C., and Lanning-Rush, J.L.,2017, Methods for estimating annual exceedance-probability streamflows for streams in Kansas based on data through water year 2015: U.S. Geological Survey Scientific Investigations Report 2017–5063, 20 p. (https://doi.org/10.3133/sir20175063)

Hodgkins, G.A. and Martin, G.R.,2003, Estimating the Magnitude of Peak Flows for Streams in Kentucky for Selected Recurrence Intervals: U.S. Geological Survey Water-Resources Investigations Report 03-4180, 69 p. (http://water.usgs.gov/pubs/wri/wri034180/) Martin, G.R., Ruhl, K.J., Moore, B.L., and Rose, M.F.,1997, Estimation of Peak-Discharge Frequency of Urban Streams in Jefferson County, Kentucky: U.S. Geological Survey Water-

Resources Investigations Report 97-4219 (http://pubs.er.usgs.gov/publication/wri974219)

```
Martin, G.R., 2002, Estimating Mean Annual Streamflow of Rural Streams in Kentucky: U.S.
Geological Survey Water-Resources Investigations Report 02-4206, 35 p.
(http://pubs.er.usgs.gov/publication/wri024206)
Martin, G.R., and Arihood, L.D., 2010, Methods for estimating selected low-flow frequency
statistics for unregulated streams in Kentucky: U.S. Geological Survey Scientific
Investigations Report 2010-5217, 83 p. (http://pubs.usgs.gov/sir/2010/5217/)
Martin, G. R. and Ruhl, K. J., 1993, Regionalization of harmonic-mean streamflows in
Kentucky: U.S. Geological Survey Water-Resources Investigations Report 92-4173, 47 p., 1
pl. (http://pubs.er.usgs.gov/publication/wri924173StreamStats_KY_20140226.mdb)
Brockman, R. A., Agouridis, C. T., Workman, S. R., Ormsbee, L. E., Fogle, A. W., 2012,
Bankfull regional curves for the Inner and Outer Bluegrass Regions of Kentucky, Journal of
the American Water Resources Association, v. 48, no. 2, p. 391-406.
(http://onlinelibrary.wiley.com/doi/10.1111/j.1752-1688.2011.00621.x/full)
TR No.70, (2004) Regionalized Regression Equations for Estimating Low-Flow
Characteristics for selected Louisiana Streams
(http://la.water.usgs.gov/publications/pdfs/TR70.pdf)
TR No.60, (1998) Floods in Louisiana, Magnitude and Frequency, Fifth Edition (not
available)
Landers, M.N., 1985, Floodflow Frequency of Streams in the Alluvial Plain of the Lower
Mississippi River in Mississippi, Arkansas, and Louisiana: U.S. Geological Survey Water-
Resources Investigations Report 85-4150, 21 p.
(http://pubs.er.usgs.gov/publication/wri854150)
Lombard, P. J., Tasker, G. D., and Nielsen, M. G., 2003, August Median Streamflow on
Ungaged Streams in Eastern Aroostook County, Maine: U.S. Geological Survey Water-
Resources Investigations Report 03-4225, 20 p.
(http://water.usgs.gov/pubs/wri/wri034225/pdf/wrir03-4225.pdf)
Lombard, P. J., 2004, August Median Streamflow on Ungaged Streams in Eastern Coastal
Maine: U.S. Geological Survey Scientific Investigations Report 2004-5157, 15 p.
(http://water.usgs.gov/pubs/sir/2004/5157/)
Dudley, R.W., 2004, Estimating Monthly, Annual, and Low 7-Day, 10-Year Streamflows for
Ungaged Rivers in Maine: U.S. Geological Survey Scientific Investigations Report 2004-
5026, 22 p. (http://water.usgs.gov/pubs/sir/2004/5026/pdf/sir2004-5026.pdf)
Hodgkins, G. A., 1999, Estimating the Magnitude of Peak Flows for Streams in Maine for
Selected Recurrence Intervals: U.S. Geological Survey Water-Resources Investigations
Report 99-4008, 45 p. (https://pubs.er.usgs.gov/publication/wri994008)
Dudley, R.W., 2004, Hydraulic-Geometry Relations for Rivers in Coastal and Central Maine:
U.S. Geological Survey Scientific Investigations Report 2004-5042, 30 p
(http://pubs.usgs.gov/sir/2004/5042/pdf/sir2004-5042.pdf)
Lombard, P.J., 2010, June and August median streamflows estimated for ungaged streams
in southern Maine: U.S. Geological Survey Scientific Investigations Report 2010-5179, 16
p. (http://pubs.usgs.gov/sir/2010/5179/pdf/sir2010-5179.pdf)
Lombard, P.J., and Hodgkins, G.A., 2015, Peak flow regression equations for small, ungaged
streams in Maine— Comparing map-based to field-based variables: U.S. Geological Survey
Scientific Investigations Report 2015-5049, 12 p. (http://dx.doi.org/10.3133/sir20155049)
Dudley, R.W., 2015, Regression equations for monthly and annual mean and selected
percentile streamflows for ungaged rivers in Maine: U.S. Geological Survey Scientific
Investigations Report 2015-5151, 35 p. (http://dx.doi.org/10.3133/sir20155151)
Thomas, Jr., W.O. and Moglen, G.E., 2010, An Update of Regional Regression Equations for
```

Maryland, Appendix 3 in Application of Hydrologic Methods in Maryland, Third Edition,

September 2010: Maryland State Highway Administration and Maryland Department of the Environment, 38 p.

(http://gishydro.eng.umd.edu/HydroPanel/hydrology_panel_report_3rd_edition_final.pdf)
Chaplin, J.J.,2005, Development of regional curves relating bankfull-channel geometry and
discharge to drainage area for streams in Pennsylvania and selected areas of Maryland:
U.S. Geological Survey Scientific Investigations Report 2005-5147, 34 p.

(https://pubs.usgs.gov/sir/2005/5147/SIR2005-5147.pdf)

Ries, K.G., III,2000, Methods for estimating low-flow statistics for Massachusetts streams: U.S. Geological Survey Water Resources Investigations Report 00-4135, 81 p. (http://pubs.usgs.gov/wri/wri004135/)

Bent, G.C., and Steeves, P.A.,2006, A revised logistic regression equation and an automated procedure for mapping the probability of a stream flowing perennially in Massachusetts: U.S. Geological Survey Scientific Investigations Report 2006–5031, 107 p. (http://pubs.usgs.gov/sir/2006/5031/pdfs/SIR_2006-5031rev.pdf)

Bent, G.C., and Waite, A.M.,2013, Equations for estimating bankfull channel geometry and discharge for streams in Massachusetts: U.S. Geological Survey Scientific Investigations Report 2013-5155, 62 p., (http://pubs.usgs.gov/sir/2013/5155/)

Zarriello, P.J.,2017, Magnitude of flood flows at selected annual exceedance probabilities for streams in Massachusetts: U.S. Geological Survey Scientific Investigations Report 2016-5156, 99 p. (https://dx.doi.org/10.3133/sir20165156)

Holtschlag, D.J. and Croskey, H.M.,1984, Statistical Methods for Estimating Flow Characteristics of Michigan Streams: U.S. Geological Survey Water-Resources Investigations Report 84-4207, 80 p. (http://pubs.er.usgs.gov/usgspubs/wri/wri844207) Lorenz, D.L., Sanocki, C.A., and Kocian, M.J.,2009, Techniques for Estimating the Magnitude and Frequency of Peak Flows on Small Streams in Minnesota Based on Data through Water Year 2005: U.S. Geological Survey Scientific Investigations Report 2009-5250, 54 p. (http://pubs.usgs.gov/sir/2009/5250/pdf/sir2009-5250.pdf)

Ziegeweid, J.R., Lorenz, D.L., Sanocki, C.A., and Czuba, C.R.,2015, Methods for estimating flow-duration curve and low-flow frequency statistics for ungaged locations on small streams in Minnesota: U.S. Geological Survey Scientific Investigations Report 2015–5170, 23 p. (http://dx.doi.org/10.3133/sir20155170)

Anderson, B.T.,2018, Flood frequency of rural streams in Mississippi, 2013: U.S. Geological Survey Scientific Investigations Report 2018–5148, 12 p.

(https://doi.org/10.3133/sir20185148)

Southard, R.E., and Veilleux, A.G.,2014, Methods for estimating annual exceedance-probability discharges and largest recorded floods for unregulated streams in rural Missouri: U.S. Geological Survey Scientific Investigations Report 2014–5165, 39 p. (http://pubs.usgs.gov/sir/2014/5165/)

Southard, R.E.,2013, Computed statistics at streamgages, and methods for estimating low-flow frequency statistics and development of regional regression equations for estimating low-flow frequency statistics at ungaged locations in Missouri: U.S. Geological Survey Scientific Investigations Report 2013–5090, 28 p. (http://pubs.usgs.gov/sir/2013/5090/) Parrett, Charles and Hull, J.A.,1985, A method for estimating mean and low flows of streams in national forests of Montana: U.S. Geological Survey Water-Resources Investigations Report 85-4071, 13 p. (https://pubs.er.usgs.gov/publication/wri854071) Parrett, Charles and Cartier, K.D. ,1999, Methods for estimating monthly streamflow characteristics at ungaged sites in western Montana: U.S. Geological Survey Water-Supply Paper 2365, 30 p. (http://pubs.er.usgs.gov/publication/wsp2365)

Parrett, Charles and Johnson, D.R.,2004, Methods for Estimating Flood Frequency in Montana Based on Data through Water Year 1998: U.S. Geological Survey Water-Resources Investigations Report 03-4308, 102 p. (http://water.usgs.gov/pubs/wri/wri03-4308/) Sando, Roy, Sando, S.K., McCarthy, P.M., and Dutton, D.M.,2016, Methods for estimating peak-flow frequencies at ungaged sites in Montana based on data through water year 2011: U.S. Geological Survey Scientific Investigations Report 2015–5019–F, 30 p. (https://doi.org/10.3133/sir20155019)

McCarthy, P.M., Sando, Roy, Sando, S.K., and Dutton, D.M.,2016, Methods for estimating streamflow characteristics at ungaged sites in western Montana based on data through water year 2009: U.S. Geological Survey Scientific Investigations Report 2015-5019-G, 19 p. (https://doi.org/10.3133/sir20155019)

Soenksen, P.J., Miller, L.D., Sharpe, J.B. and Watton, J.R.,1999, Peak-Flow Frequency Relations and Evaluation of the Peak-Flow Gaging Network in Nebraska: U. S. Geological Survey Water-Resources Investigations Report 99-4032, 48 p,

(https://pubs.er.usgs.gov/publication/wri994032)

Flynn, R.H. and Tasker, G.D.,2002, Development of Regression Equations to Estimate Flow Durations and Low-Flow-Frequency Statistics in New Hampshire Streams: U.S.Geological Survey Scientific Investigations Report 02-4298, 66 p. (http://pubs.water.usgs.gov/wrir02-4298)

Olson, S.A.,2009, Estimation of flood discharges at selected recurrence intervals for streams in New Hampshire: U.S.Geological Survey Scientific Investigations Report 2008-5206, 57 p. (http://pubs.usgs.gov/sir/2008/5206/)

Flynn, R.H. and Tasker, G.D.,2004, Generalized Estimates from Streamflow Data of Annual and Seasonal Ground-Water-Recharge Rates for Drainage Basins in New Hampshire, U.S. Geological Survey Scientific Investigations Report 2004-5019, 67 p.

(http://pubs.usgs.gov/sir/2004/5019/http://pubs.usgs.gov/sir/2004/5019/)

Watson, K.M., and Schopp, R.D., 2009, Methodology for estimation of flood magnitude and frequency for New Jersey streams, U.S. Geological Survey Scientific Investigations Report 2009-5167, 51 p. (http://pubs.usgs.gov/sir/2009/5167/)

Watson, K.M., and McHugh, A.R.,2014, Regional regression equations for the estimation of selected monthly low-flow duration and frequency statistics at ungaged sites on streams in New Jersey: U.S. Geological Survey Scientific Investigations Report 2014-5004, 59 p. (baseline, period-or-record statistics)

(http://dx.doi.org/10.3133/sir20145004StreamStatsDB\2019_12_13_DataSource_table.xlsxDa Waltemeyer, S.D.,2002, Analysis of the magnitude and frequency of the 4-day, 3-year annual low flow on unregulated streams in New Mexico: U. S. Geological Survey Water-Resources Investigations Report 01-4271, 22 p.

(http://nm.water.usgs.gov/publications/abstracts/wrir01-4271.pdf)

Waltemeyer, S.D., 2008, Analysis of the Magnitude and Frequency of Peak Discharge and Maximum Observed Peak Discharge in New Mexico and Surrounding Areas: U.S. Geological Survey Scientific Investigations Report 2008-5119, 105 p.

(http://pubs.usgs.gov/sir/2008/5119/)

Lumia, Richard, Freehafer, D.A., and Smith, M.J.,2006, Magnitude and Frequency of Floods in New York: U.S. Geological Survey Scientific Investigations Report 2006–5112, 152 p. (http://pubs.usgs.gov/sir/2006/5112/)

Stedfast, D.A.,1984, Evaluation of Six Methods for Estimating Magnitude and Frequency of Peak Discharges on Urban Streams in New York: U. S. Geological Survey Water-Resources Investigations Report 84-4350, 24 p.

(http://onlinepubs.er.usgs.gov/djvu/WRI/wrir_84_4350.djvu)

Mulvihill, C.I., Baldigo, B.P., Miller, S.J., and DeKoskie, Douglas,2009, Bankfull Discharge and Channel Characteristics of Streams in New York State: U.S. Geological Survey Scientific Investigations Report 2009-5144, 51 p. (http://pubs.usgs.gov/sir/2009/5144/) Barnes, C. R.,1986, Method for estimating low-flow statistics for ungaged streams in the lower Hudson River Basin, New York: U. S. Geological Survey Water-Resources Investigations Report 85-4070, 22 p.

(http://onlinepubs.er.usgs.gov/djvu/WRI/wrir_85_4070.djvu)

Randall, A.D., 2010, Low flow of streams in the Susquehanna River basin of New York: U.S. Geological Survey Scientific Investigations Report 2010-5063, 57 p.

(http://pubs.usgs.gov/sir/2010/5063/http://pubs.usgs.gov/sir/2010/5063/)

Gazoorian, C.L.,2015, Estimation of unaltered daily mean streamflow at ungaged streams of New York, excluding Long Island, water years 1961–2010: U.S. Geological Survey Scientific Investigations Report 2014–5220, 29 p. (https://pubs.usgs.gov/sir/2014/5220/) Giese, G. L. and Mason, R.R., Jr.,1993, Low-flow characteristics of streams in North Carolina: U.S. Geological Survey Water-Supply Paper 2403, 29 p.

(https://pubs.er.usgs.gov/publication/wsp2403)

Mason, Robert R., Jr.; Fuste, Luis A.; King, Jeffrey N.; Thomas, Wilbert O., Jr.,2002, The National Flood-Frequency Program -- Methods for Estimating Flood Magnitude and Frequency in Rural and Urban Areas in North Carolina, 2001: U.S. Geological Survey Fact Sheet 007-00, 4 p. (http://pubs.er.usgs.gov/publication/fs00700)

Weaver, J.C., Feaster, T.D., and Gotvald, A.J.,2009, Magnitude and frequency of rural floods in the Southeastern United States, through 2006—Volume 2, North Carolina: U.S. Geological Survey Scientific Investigations Report 2009–5158, 111 p.

(http://pubs.usgs.gov/sir/2009/5158/)

Williams-Sether, T.,2015, Regional regression equations to estimate peak-flow frequency at sites in North Dakota using data through 2009: U.S. Geological Survey Scientific Investigations Report 2015–5096, 12 p. (http://dx.doi.org/10.3133/sir20155096) Koltun, G.F., Kula, S.P., and Puskas, B.M.,2006, A Streamflow Statistics (StreamStats) Web

Application for Ohio: U.S. Geological Survey Scientific Investigations Report 2006-5312, 62 p. (http://pubs.usgs.gov/sir/2006/5312/)

Sherwood, J.M.,1994, Estimation of peak-frequency relations, flood hydrographs, and volume-duration-frequency relations of ungaged small urban streams in Ohio: U. S. Geological Survey Water-Supply Paper 2432, 42 p.

(http://pubs.er.usgs.gov/usgspubs/wsp/wsp2432)

Koltun, G. F., and Whitehead, M. T.,2002, Techniques for Estimating Selected Streamflow Characteristics of Rural, Unregulated Streams in Ohio: U. S. Geological Survey Water-Resources Investigations Report 02-4068, 50 p

(https://pubs.er.usgs.gov/publication/wri024068)

Koltun, G. F., and Schwartz, Ronald R.,1987, MULTIPLE-REGRESSION EQUATIONS FOR ESTIMATING LOW FLOWS AT UNGAGED STREAM SITES IN OHIO: U.S. Geological Survey Water-Resources Investigations Report 86-4354, 39 p.

(http://pubs.er.usgs.gov/usgspubs/wri/wri864354)

Koltun, G.F., and Kula, S.P.,2013, Methods for estimating selected low-flow statistics and development of annual flow-duration statistics for Ohio: U.S. Geological Survey Scientific Investigations Report 2012–5138, 195 p. (http://pubs.usgs.gov/sir/2012/5138/)

Koltun, G.F.,2019, Flood-frequency estimates for Ohio streamgages based on data through water year 2015 and techniques for estimating flood-frequency characteristics of rural, unregulated Ohio streams: U.S. Geological Survey Scientific Investigations Report 2019–5018, xx p. (https://dx.doi.org/10.3133/sir20195018)

Esralew, R.A., Smith, S.J.,2009, Methods for estimating flow-duration and annual mean-flow statistics for ungaged streams in Oklahoma: U.S. Geological Survey Scientific Investigations Report 2009-5267, 131 p. (http://pubs.usgs.gov/sir/2009/5267/) Smith, S.J., Lewis, J.M., and Graves, G.M.,2015, Methods for estimating the magnitude and frequency of peak streamflows at ungaged sites in and near the Oklahoma Panhandle: U.S. Geological Survey Scientific Investigations Report 2015-5134, 35 p. (http://dx.doi.org/10.3133/sir20155134)

Lewis, J.M., Hunter, S.L., and Labriola, L.G.,2019, Methods for estimating the magnitude and frequency of peak streamflows for unregulated streams in Oklahoma developed by using streamflow data through 2017: U.S. Geological Survey Scientific Investigations Report 2019–5143, 39 p. (https://doi.org/10.3133/sir20195143)

Laenen, Antonius,1980, Storm Runoff As Related to Urbanization in the Portland, Oregon - Vancouver, Washington Area: U.S. Geological Survey Open-File Report 80-689, 71 p. (http://pubs.er.usgs.gov/usgspubs/wri/wri834143)

Cooper, R.M.,2005, Estimation of Peak Discharges for Rural, Unregulated Streams in Western Oregon: U.S. Geological Survey Scientific Investigations Report 2005-5116, 76 p. (http://pubs.usgs.gov/sir/2005/5116/pdf/sir2005-5116.pdf)

Risley, John, Stonewall, Adam, and Haluska, Tana, 2008, Estimating flow-duration and low-flow frequency statistics for unregulated streams in Oregon: U.S. Geological Survey Scientific Investigations Report 2008-5126, 22 p. (http://pubs.usgs.gov/sir/2008/5126/) Cooper, Richard, 2006, Estimation of Peak Discharges for Rural, Unregulated Streams in Eastern Oregon, Oregon Water Resources Department OFR SW 06-001, Salem, OR. (https://digital.osl.state.or.us/islandora/object/osl%3A14736/datastream/OBJ/view) Stuckey, M.H., 2006, Low-flow, base-flow, and mean-flow regression equations for Pennsylvania streams: U.S. Geological Survey Scientific Investigations Report 2006-5130, 84 p. (http://pubs.usgs.gov/sir/2006/5130/)

Stuckey, M.H., Koerkle, E.H., and Ulrich, J.E.,2012, Estimation of baseline daily mean streamflows for ungaged locations on Pennsylvania streams, water years 1960–2008: U.S. Geological Survey Scientific Investigations Report 2012–5142, 61 p. (http://pubs.usgs.gov/sir/2012/5142/)

Clune, J.W., Chaplin, J.J., and White, K.E., 2018, Comparison of regression relations of bankfull discharge and channel geometry for the glaciated and nonglaciated settings of Pennsylvania and southern New York: U.S. Geological Survey Scientific Investigations Report 2018-5066, 20 p. (https://doi.org/10.3133/sir20185066)

Roland, M.A., and Stuckey, M.H.,2008, Regression equations for estimating flood flows at selected recurrence intervals for ungaged streams in Pennsylvania: U.S. Geological Survey Scientific Investigations Report 2008-5102, 57p. (http://pubs.usgs.gov/sir/2008/5102/) Zarriello, P.J., Ahearn, E.A., and Levin, S.B.,2012, Magnitude of flood flows for selected annual exceedance probabilities in Rhode Island through 2010: U.S. Geological Survey Scientific Investigations Report 2012-5109, 93 p. (http://pubs.usgs.gov/sir/2012/5109) Bent, G.C., Steeves, P.A., and Waite, A.M.,2014, Equations for estimating selected streamflow statistics in Rhode Island: U.S. Geological Survey Scientific Investigations Report 2014-5010, 65 p. (http://dx.doi.org/10.3133/sir20145010)

Feaster, T.D., Gotvald, A.J., and Weaver, J.C.,2009, Magnitude and Frequency of Rural Floods in the Southeastern United States, 2006: Volume 3, South Carolina: U.S. Geological Survey Scientific Investigations Report 2009-5156, 226 p.

(http://pubs.usgs.gov/sir/2009/5156/)

Sando, Steven K.,1998, A Method for Estimating Magnitude and Frequency of Floods in South Dakota: U.S. Geological Survey Water-Resources Investigations Report 98-4055, 48

```
p. (http://pubs.water.usgs.gov/wri98-4055/)
Law, G.S., and Tasker G.D., 2003, Flood-Frequency Prediction Methods for Unregulated
Streams of Tennessee, 2000: U.S. Geological Survey Water-Resources Investigations
Report 03-4176, 79p. (http://pubs.usgs.gov/wri/wri034176/)
Neely, B.L., Jr., 1984, Flood Frequency and Storm Runoff of Urban Areas of Memphis and
Shelby County, Tennessee: U.S. Geological Survey Water-Resources Investigations Report
84-4110, 51 p. (http://pubs.usgs.gov/wri/wrir_84-4110/)
Robbins, Clarence H., 1984, Synthesized Flood Frequency of Small Urban Streams in
Tennessee: U.S. Geological Survey Water-Resources Investigations Report 84-4182, 24 p.
(http://onlinepubs.er.usgs.gov/djvu/WRI/wrir_84_4182.djvu)
Law, G.S., Tasker, G.D., and Ladd, D.E., 2009, Streamflow-characteristic estimation methods
for unregulated streams of Tennessee: U.S. Geological Survey Scientific Investigations
Report 2009-5159, 212 p., 1 pl. (http://pubs.usgs.gov/sir/2009/5159/)
Asquith, W.H., Slade, R.M., Jr., 1999, Site-specific estimation of peak-stream flow
frequency using generalized least squares regression for natural basins in Texas: U.S.
Geological Survey Water-Resources Investigations Report 99-4172, 19 p.
(http://pubs.water.usgs.gov/wri994172)
Asquith, William H.,1998, Peak-flow frequency for tributaries of the Colorado River
downstream of Austin, Texas U.S. Geological Survey Water-Resources Investigations
Report 98-4015, 26 p. (http://pubs.water.usgs.gov/wri98-4015/)
Raines, Timothy H., 1998, Peak-discharge frequency and potential extreme peak discharge
for natural streams in the Brazos River basin, Texas: U.S. Geological Survey Water-
Resources Investigations Report 98-4178, 47 p., 1 plate (http://pubs.water.usgs.gov/wri98-
4178/)
Land, L.F., Schroeder, E.E. and Hampton, B.B., 1982, Techniques for Estimating the
Magnitude and Frequency of Floods in the Dallas-Fort Worth Metropolitan Area, Texas: U.S.
Geological Survey Water-Resources Investigations Report 82-18, 55 p. ()
Asquith, W.H., Slade, R. M., Lanning-Rush, Jennifer, 1996, Peak-flow frequency and extreme
flood potential for streams in the vicinity of the Highland Lakes, central Texas: U.S.
Geological Survey Water-Resources Investigations Report 96-4072
(http://onlinepubs.er.usgs.gov/djvu/WRI/wrir_96_4072_plt.djvu)
Liscum, Fred and Massey, B.C., 1980, Technique for Estimiating the Magnitude and
Frequency of Floods in the Houston, Texas, Metropolitan Area: U.S. Geological Survey
Water-Resources Investigations Report 80-17, 29 p. ()
Asquith, W.H., and Roussel, M.C., 2009, Regression equations for estimation of annual
peak-streamflow frequency for undeveloped watersheds in Texas using an L-moment-
based, PRESS-minimized, residual-adjusted approach: U.S. Geological Survey Scientific
Investigations Report 2009-5087, 48 p. (http://pubs.usgs.gov/sir/2009/5087/)
Kenney, T.A., Wilkowske, C.D., and Wright, S.J., 2007, Methods for Estimating Magnitude
and Frequency of Peak Flows for Natural Streams in Utah: U.S. Geological Survey Scientific
Investigations Report 2007-5158, 28 p. (http://pubs.usgs.gov/sir/2007/5158/)
Wilkowske, C.D., Kenney, T.A., and Wright, S.J., 2009, Methods for Estimating Monthly and
Annual Streamflow Statistics at Ungaged Sites in Utah: U.S. Geological Survey Scientific
Investigations Report 2008-5230, 62 p. (http://pubs.usgs.gov/sir/2008/5230/)
Olson, S.A., 2002, Flow-frequency characteristics of Vermont streams: U.S. Geological
Survey Water-Resources Investigations Report 02-4238, 47 p.
(http://pubs.usgs.gov/wri/wrir02-4238/)
Olson, S.A., 2014, Estimation of flood discharges at selected annual exceedance
```

probabilities for unregulated, rural streams in Vermont, with a section on Vermont regional

```
skew regression, by Veilleux, A.G.: U.S. Geological Survey Scientific Investigations Report
2014-5078, 27 p. plus appendixes. (http://pubs.usgs.gov/sir/2014/5078/)
Olson, S.A., and Brouillette, M.C., 2006, A logistic regression equation for estimating the
probability of a stream in Vermont having intermittent flow: U.S. Geological Survey
Scientific Investigations Report 2006-5217, 15 p. (https://pubs.usgs.gov/sir/2006/5217/)
Austin, S.H., Krstolic, J.L., and Wiegand, Ute, 2011, Low-flow characteristics of Virginia
streams: U.S. Geological Survey Scientific Investigations Report 2011-5143, 122 p. + 9
tables on CD. (http://pubs.usgs.gov/sir/2011/5143/)
Austin, S.H., Krstolic, J.L., and Wiegand, Ute, 2011, Peak-flow characteristics of Virginia
streams: U.S. Geological Survey Scientific Investigations Report 2011-5144, 106 p. + 3
tables and 2 appendixes on CD. (http://pubs.usgs.gov/sir/2011/5144/)
Austin, S.H., 2014, Methods and equations for estimating peak streamflow per square mile
in Virginia's urban basins: U.S. Geological Survey Scientific Investigations Report 2014-
5090, 25 p. (http://pubs.usgs.gov/sir/2014/5090/http://pubs.usgs.gov/sir/2014/5090/)
Curran, C.A. and Olsen, T.D., 2009, Estimating Low-Flow Frequency Statistics and
Hydrologic Analysis of Selected Streamflow-Gaging Stations, Nooksack River Basin,
Northwestern Washington and Canada: U.S. Geological Survey Scientific Investigations
Report 2009-5170, 44 p. (http://pubs.usgs.gov/sir/2009/5170/)
Curran, C.A., Eng, Ken, and Konrad, C.P., 2012, Analysis of low flows and selected methods
for estimating low-flow characteristics at partial-record and ungaged stream sites in
western Washington: U.S. Geological Survey Scientific Investigations Report 2012-5078, 46
p. (http://pubs.usgs.gov/sir/2012/5078/)
Mastin, M.C., Konrad, C.P., Veilleux, A.G., and Tecca, A.E., 2016, Magnitude, frequency, and
trends of floods at gaged and ungaged sites in Washington, based on data through water
year 2014 (ver 1.1, October 2016): U.S. Geological Survey Scientific Investigations Report
2016-5118, 70 p. (http://dx.doi.org/10.3133/sir20165118)
Wiley, Jeffrey B., 2008, Estimating Selected Streamflow Statistics Representative of 1930-
2002 in West Virginia: U.S. Geological Survey Scientific Investigations Report 2008-5105,
24 p. (http://pubs.usgs.gov/sir/2008/5105/)
Wiley, Jeffrey B., 1987, Techniques for estimating flood depth frequency relations for
streams in West Virginia: U.S. Geological Survey Water-Resources Investigations Report
87-4111, 17 p. (http://pubs.er.usgs.gov/usgspubs/wri/wri874111)
Wiley, J.B., and Atkins, J.T., Jr., 2010, Estimation of flood-frequency discharges for rural,
unregulated streams in West Virginia: U.S. Geological Survey Scientific Investigations
Report 2010-5033, 78 p. (http://pubs.usgs.gov/sir/2010/5033/)
Wiley, J.B., and Atkins, J.T., Jr., 2010, Estimation of selected seasonal streamflow statistics
representative of 1930-2002 in West Virginia: U.S. Geological Survey Scientific
Investigations Report 2010-5185, 20 p. (http://pubs.usgs.gov/sir/2010/5185/)
Conger, Duane H., 1986, Estimating Magnitude and Frequency of Floods for Wisconsin
Urban Streams: U.S. Geological Survey Water-Resources Investigations Report 86-4005, 18
p. (http://pubs.er.usgs.gov/publication/wri864005)
Walker, J.F., Peppler, M.C., Danz, M.E., and Hubbard, L.E., 2017, Flood-frequency
characteristics of Wisconsin streams (ver. 2.1, December 2017): Reston, Virginia, U.S.
Geological Survey Scientific Investigations Report 2016-5140, 33 p., 1 plate, 2 appendixes
```

Miller, Kirk A.,2003, Peak-flow Characteristics of Wyoming Streams: U.S. Geological Survey Water-Resources Investigations Report 03-4107, 79 p. (http://pubs.usgs.gov/wri/wri034107/)

(https://doi.org/10.3133/sir20165140)

Ramos-Ginés, Orlando,1999, Estimation of Magnitude and Frequency of Floods for Streams in Puerto Rico: New Empirical Models: U. S. Geological Survey Water-Resources Investigations Report 99-4142, 41 p. (http://pubs.usgs.gov/wri/wri994142/) Moody, J.A.,2012, An analytical method for predicting postwildfire peak discharges: U.S. Geological Survey Scientific Investigations Report 2011-5236, 36 p. (https://pubs.usgs.gov/sir/2011/5236/)

Bankfull Statistics Parameters [Bankfull Statewide SIR2013 5155]

Parameter Code	Parameter Name	Value	Units	Min Limit	Max Limit
DRNAREA	Drainage Area	0.0906	square miles	0.6	329
BSLDEM10M	Mean Basin Slope from 10m DEM	11.842	percent	2.2	23.9

Bankfull Statistics Disclaimers[Bankfull Statewide SIR2013 5155]

One or more of the parameters is outside the suggested range. Estimates were extrapolated with unknown errors

Bankfull Statistics Flow Report[Bankfull Statewide SIR2013 5155]

Statistic	Value	Unit
Bankfull Width	6.39	ft
Bankfull Depth	0.51	ft
Bankfull Area	3.2	ft^2
Bankfull Streamflow	8.95	ft^3/s

Bankfull Statistics Citations

Bent, G.C., and Waite, A.M.,2013, Equations for estimating bankfull channel geometry and discharge for streams in Massachusetts: U.S. Geological Survey Scientific Investigations Report 2013–5155, 62 p., (http://pubs.usgs.gov/sir/2013/5155/)

Probability Statistics Parameters[Perennial Flow Probability]

Parameter Code	Parameter Name	Value	Units	Min Limit	Max Limit
DRNAREA	Drainage Area	0.0906	square miles	0.01	1.99

Parameter Code	Parameter Name	Value	Units	Min Limit	Max Limit
PCTSNDGRV	Percent Underlain By Sand And Gravel	0	percent	0	100
FOREST	Percent Forest	99.23	percent	0	100
MAREGION	Massachusetts Region	1	dimensionless	0	1

Probability Statistics Flow Report[Perennial Flow Probability]

PII: Prediction Interval-Lower, PIu: Prediction Interval-Upper, SEp: Standard Error of Prediction, SE: Standard Error (other -- see report)

Statistic	Value	Unit	PC
Probability Stream Flowing Perennially	0.245	dim	71

Probability Statistics Citations

Bent, G.C., and Steeves, P.A.,2006, A revised logistic regression equation and an automated procedure for mapping the probability of a stream flowing perennially in Massachusetts: U.S. Geological Survey Scientific Investigations Report 2006–5031, 107 p. (http://pubs.usgs.gov/sir/2006/5031/pdfs/SIR_2006-5031rev.pdf)

USGS Data Disclaimer: Unless otherwise stated, all data, metadata and related materials are considered to satisfy the quality standards relative to the purpose for which the data were collected. Although these data and associated metadata have been reviewed for accuracy and completeness and approved for release by the U.S. Geological Survey (USGS), no warranty expressed or implied is made regarding the display or utility of the data for other purposes, nor on all computer systems, nor shall the act of distribution constitute any such warranty.

USGS Software Disclaimer: This software has been approved for release by the U.S. Geological Survey (USGS). Although the software has been subjected to rigorous review, the USGS reserves the right to update the software as needed pursuant to further analysis and review. No warranty, expressed or implied, is made by the USGS or the U.S. Government as to the functionality of the software and related material nor shall the fact of release constitute any such warranty. Furthermore, the software is released on condition that neither the USGS nor the U.S. Government shall be held liable for any damages resulting from its authorized or unauthorized use.

USGS Product Names Disclaimer: Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

Application Version: 4.4.0


S-MJR-2 StreamStats Report

Region ID: MA

Workspace ID: MA20200828024143728000

Clicked Point (Latitude, Longitude): 42.41516, -72.46782

Time: 2020-08-27 22:41:59 -0400



Basin Characteristics				
Parameter Code	Parameter Description	Value	Unit	
DRNAREA	Area that drains to a point on a stream	0.21	square miles	
ELEV	Mean Basin Elevation	758	feet	
LC06STOR	Percentage of water bodies and wetlands determined from the NLCD 2006	0	percent	
BSLDEM250	Mean basin slope computed from 1:250K DEM	8.113	percent	
DRFTPERSTR	Area of stratified drift per unit of stream length	0.13	square mile per mile	

Parameter Code	Parameter Description	Value	Unit
MAREGION	Region of Massachusetts 0 for Eastern 1 for Western	1	dimensionless
BSLDEM10M	Mean basin slope computed from 10 m DEM	10.885	percent
PCTSNDGRV	Percentage of land surface underlain by sand and gravel deposits	22.71	percent
FOREST	Percentage of area covered by forest	91.39	percent
ACRSDFT	Area underlain by stratified drift	0.0471	square miles
CENTROIDX	Basin centroid horizontal (x) location in state plane coordinates	120682.7	meters
CENTROIDY	Basin centroid vertical (y) location in state plane units	907103.8	meters
CRSDFT	Percentage of area of coarse-grained stratified drift	22.71	percent
LAKEAREA	Percentage of Lakes and Ponds	0	percent
LC11DEV	Percentage of developed (urban) land from NLCD 2011 classes 21-24	3.49	percent
LC11IMP	Average percentage of impervious area determined from NLCD 2011 impervious dataset	0.0544	percent
MAXTEMPC	Mean annual maximum air temperature over basin area, in degrees Centigrade	13.7	feet per mi
OUTLETX	Basin outlet horizontal (x) location in state plane coordinates	120345	feet
OUTLETY	Basin outlet vertical (y) location in state plane coordinates	907635	feet
PRECPRIS00	Basin average mean annual precipitation for 1971 to 2000 from PRISM	48	inches
STRMTOT	total length of all mapped streams (1:24,000-scale) in the basin	0.36	miles
WETLAND	Percentage of Wetlands	3.87	percent

Peak-Flow Statistics Parameters [Peak Statewide 2016 5156]				
Parameter Code	Parameter Name	Value Units	Min Limit	Max Limit

Parameter Code	Parameter Name	Value U	Jnits	Min Limit	Max Limit
DRNAREA	Drainage Area		square niles	0.16	512
ELEV	Mean Basin Elevation	758 f	eet	80.6	1948
LC06STOR	Percent Storage from NLCD2006	0 р	percent	0	32.3

Peak-Flow Statistics Flow Report[Peak Statewide 2016 5156]

PII: Prediction Interval-Lower, PIu: Prediction Interval-Upper, SEp: Standard Error of Prediction, SE: Standard Error (other -- see report)

Statistic	Value	Unit	PII	Plu	SEp
2 Year Peak Flood	16.8	ft^3/s	8.36	33.7	42.3
5 Year Peak Flood	29.1	ft^3/s	14.2	59.4	43.4
10 Year Peak Flood	39.5	ft^3/s	18.8	82.8	44.7
25 Year Peak Flood	55.1	ft^3/s	25.3	120	47.1
50 Year Peak Flood	68.4	ft^3/s	30.3	154	49.4
100 Year Peak Flood	82.9	ft^3/s	35.5	193	51.8
200 Year Peak Flood	99	ft^3/s	41.1	238	54.1
500 Year Peak Flood	123	ft^3/s	48.6	312	57.6

Peak-Flow Statistics Citations

Zarriello, P.J.,2017, Magnitude of flood flows at selected annual exceedance probabilities for streams in Massachusetts: U.S. Geological Survey Scientific Investigations Report 2016-5156, 99 p. (https://dx.doi.org/10.3133/sir20165156)

Low-Flow Statistic					
Parameter Code	Parameter Name	Value	Units	Min Limit	Max Limit
DRNAREA	Drainage Area	0.21	square miles	1.61	149
BSLDEM250	Mean Basin Slope from 250K DEM	8.113	percent	0.32	24.6
DRFTPERSTR	Stratified Drift per Stream Length	0.13	square mile per mile	0	1.29

Parameter Code	Parameter Name	Value Units	Min Limit	Max Limit
MAREGION	Massachusetts Region	1 dimensionles	s 0	1

Low-Flow Statistics Disclaimers [Statewide Low Flow WRIR00 4135]

One or more of the parameters is outside the suggested range. Estimates were extrapolated with unknown errors

Low-Flow Statistics Flow Report[Statewide Low Flow WRIR00 4135]

Statistic	Value	Unit
7 Day 2 Year Low Flow	0.0235	ft^3/s
7 Day 10 Year Low Flow	0.0121	ft^3/s

Low-Flow Statistics Citations

Ries, K.G., III,2000, Methods for estimating low-flow statistics for Massachusetts streams: U.S. Geological Survey Water Resources Investigations Report 00-4135, 81 p. (http://pubs.usgs.gov/wri/wri004135/)

Flow-Duration Statistics Parameters [Statewide Low Flow WRIR00 4135]

Parameter Code	Parameter Name	Value	Units	Min Limit	Max Limit
DRNAREA	Drainage Area	0.21	square miles	1.61	149
DRFTPERSTR	Stratified Drift per Stream Length	0.13	square mile per mile	0	1.29
MAREGION	Massachusetts Region	1	dimensionless	0	1
BSLDEM250	Mean Basin Slope from 250K DEM	8.113	percent	0.32	24.6

Flow-Duration Statistics Disclaimers [Statewide Low Flow WRIR00 4135]

One or more of the parameters is outside the suggested range. Estimates were extrapolated with unknown errors

Flow-Duration Statistics Flow Report[Statewide Low Flow WRIR00 4135]

Statistic Value Unit

Statistic	Value	Unit
50 Percent Duration	0.194	ft^3/s
60 Percent Duration	0.124	ft^3/s
70 Percent Duration	0.0894	ft^3/s
75 Percent Duration	0.0719	ft^3/s
80 Percent Duration	0.0729	ft^3/s
85 Percent Duration	0.0556	ft^3/s
90 Percent Duration	0.0475	ft^3/s
95 Percent Duration	0.0277	ft^3/s
98 Percent Duration	0.0183	ft^3/s
99 Percent Duration	0.0126	ft^3/s

Flow-Duration Statistics Citations

Ries, K.G., III,2000, Methods for estimating low-flow statistics for Massachusetts streams: U.S. Geological Survey Water Resources Investigations Report 00-4135, 81 p. (http://pubs.usgs.gov/wri/wri004135/)

August Flow-Duration Statistics Parameters [Statewide Low Flow WRIR00 4135]

Parameter Code	Parameter Name	Value	Units	Min Limit	Max Limit
DRNAREA	Drainage Area	0.21	square miles	1.61	149
BSLDEM250	Mean Basin Slope from 250K DEM	8.113	percent	0.32	24.6
DRFTPERSTR	Stratified Drift per Stream Length	0.13	square mile per mile	0	1.29
MAREGION	Massachusetts Region	1	dimensionless	0	1

August Flow-Duration Statistics Disclaimers[Statewide Low Flow WRIR00 4135]

One or more of the parameters is outside the suggested range. Estimates were extrapolated with unknown errors

August Flow-Duration Statistics Flow Report[Statewide Low Flow WRIR00 4135]

Statistic Value Unit

Statistic	Value	Unit
August 50 Percent Duration	0.0582	ft^3/s

August Flow-Duration Statistics Citations

Ries, K.G., III,2000, Methods for estimating low-flow statistics for Massachusetts streams: U.S. Geological Survey Water Resources Investigations Report 00-4135, 81 p. (http://pubs.usgs.gov/wri/wri004135/)

Bankfull Statistics Parameters [Bankfull Statewide SIR2013 5155]

Parameter Code	Parameter Name	Value	Units	Min Limit	Max Limit
DRNAREA	Drainage Area	0.21	square miles	0.6	329
BSLDEM10M	Mean Basin Slope from 10m DEM	10.885	percent	2.2	23.9

Bankfull Statistics Disclaimers[Bankfull Statewide SIR2013 5155]

One or more of the parameters is outside the suggested range. Estimates were extrapolated with unknown errors

Bankfull Statistics Flow Report[Bankfull Statewide SIR2013 5155]

Statistic	Value	Unit
Bankfull Width	8.77	ft
Bankfull Depth	0.642	ft
Bankfull Area	5.53	ft^2
Bankfull Streamflow	15.8	ft^3/s

Bankfull Statistics Citations

Bent, G.C., and Waite, A.M.,2013, Equations for estimating bankfull channel geometry and discharge for streams in Massachusetts: U.S. Geological Survey Scientific Investigations Report 2013–5155, 62 p., (http://pubs.usgs.gov/sir/2013/5155/)

Probability Statistics Parameters[Perennial Flow Probability]

Parameter Code	Parameter Name	Value U	Jnits	Min Limit	Max Limit
DRNAREA	Drainage Area	0.21 s	square miles	0.01	1.99
PCTSNDGRV	Percent Underlain By Sand And Gravel	22.71 p	percent	0	100
FOREST	Percent Forest	91.39 p	percent	0	100
MAREGION	Massachusetts Region	1 d	dimensionless	0	1

Probability Statistics Flow Report[Perennial Flow Probability]

PII: Prediction Interval-Lower, Plu: Prediction Interval-Upper, SEp: Standard Error of Prediction, SE: Standard Error (other -- see report)

Statistic	Value	Unit	PC
Probability Stream Flowing Perennially	0.535	dim	71

Probability Statistics Citations

Bent, G.C., and Steeves, P.A.,2006, A revised logistic regression equation and an automated procedure for mapping the probability of a stream flowing perennially in Massachusetts: U.S. Geological Survey Scientific Investigations Report 2006–5031, 107 p. (http://pubs.usgs.gov/sir/2006/5031/pdfs/SIR_2006-5031rev.pdf)

USGS Data Disclaimer: Unless otherwise stated, all data, metadata and related materials are considered to satisfy the quality standards relative to the purpose for which the data were collected. Although these data and associated metadata have been reviewed for accuracy and completeness and approved for release by the U.S. Geological Survey (USGS), no warranty expressed or implied is made regarding the display or utility of the data for other purposes, nor on all computer systems, nor shall the act of distribution constitute any such warranty.

USGS Software Disclaimer: This software has been approved for release by the U.S. Geological Survey (USGS). Although the software has been subjected to rigorous review, the USGS reserves the right to update the software as needed pursuant to further analysis and review. No warranty, expressed or implied, is made by the USGS or the U.S. Government as to the functionality of the software and related material nor shall the fact of release constitute any such warranty. Furthermore, the software is released on condition that neither the USGS nor the U.S. Government shall be held liable for any damages resulting from its authorized or unauthorized use.

USGS Product Names Disclaimer: Any use of trade, firm, or product names is for descriptive purposes only and does not imply endorsement by the U.S. Government.

Application Version: 4.4.0

ATTACHMENT C
Abutter Information
(Certified Abutter List, Abutter Notification
& Affidavit of Service)




TOWN OF SHUTESBURY CERTIFIED 100' ABUTTERS LIST FOR PARCEL ZU-2 OFF PRATT CORNER RD

MAP	LOT	OWNER	CO-OWNER	MAILING ADDRESS	TOWN	ST	ZIP	LOCATION
ZU		2 W D COWLS INC		P O BOX 9677	NORTH AMHERS	AMF	01059	PRATT CORNER RD
U		3 TOWN OF AMHERST		4 BOLTWOOD AVENUE	AMHERST	MA	01002	PRATT CORNER RD
U		6 TOWN OF AMHERST		4 BOLTWOOD AVENUE	AMHERST	MA	1002	2 PRATT CORNER RD
U		57 TRAPANI JOSEF G	TRAPANI ANNA E	692 PRATT CORNER RD	AMHERST	MA	01002	692 PRATT CORNER RD
ZV		1 POVERTY MOUNTAIN FARM, LLC	C/O WEIR, K. & BANFIELD-WEIR, C	760 PRATT CORNER RD	AMHERST	MA	01002	PRATT CORNER RD

FOR: TRC

650 Suffolk ST, Lowell, MA 01854
Molly Lennon, Environmental Scientist
Mlennon@trccompanies.com

Respectfully, Leslie Bracebridge, Assessors Clerk for Kevin Rudden, Administratve Assessor 9/8/2020





Town of Shutesbury, Massachusetts

Selected Parcel: PRATT CORNER RD ID: ZU-2

Printed 9/8/2020 from http://www.mainstreetmaps.com/ma/shutesbury/public.asp







This map is for informational purposes only. It is not for appraisal of, description of, or conveyance of land. The Town of Shutesbury, Massachusetts and MainStreetGIS, LLC assume no legal responsibility for the information contained herein

SHUTESBURY CONSERVATION COMMISSION NOTIFICATION TO ABUTTERS

In accordance with the second paragraph of the Massachusetts Wetlands Protection Act (G.L. Ch. 131 §40), and §10.05(4)(a) of 310 CMR 10.00, and the Shutesbury Wetlands Protection Bylaw and regulations, you are hereby notified of a public hearing on the matter described below.

A.	An ANRAD has been filed with the Shutesbury Conservation Commission.
В.	The name of the applicant is: <u>W.D. Cowls, Inc.</u>
C.	The address/lot number of the land where the activity is proposed: <u>Pratt Corner Road</u> , <u>Shutesbury</u> , <u>MA (Parcel ID: ZU-2)</u>
D.	The proposed activity is: Review of delineated wetland resources.
	,
E.	A Public Hearing regarding this ANRAD will be held on: November 12, 2020
F.	Public Participation will be via Virtual Means Only: Governor Baker issued an Emergency Order on March 12, 2020 allowing public bodies greater flexibility in utilizing technology in the conduct of meetings under the Open Meeting Law. The Shutesbury Conservation Commission greatly values the participation of its citizens in the public meeting process, but given the current circumstances and recommendations to limit or avoid public gatherings, including Governor Baker's State of Emergency, together with the present closure of Shutesbury Town Hall, the Town has decided to implement the "remote participation" procedures allowed under Governor Baker's Emergency Order for all boards, committees, and commissions. Remote access information will be published on the Shutesbury meeting calendar: www.shutesbury.org/node/2. Click on the agenda for the meeting you wish to attend.
G	The ANRAD may be examined on the Shutesbury Conservation Commission website:

Notice of the public hearing, including date, time, and place will be published at least five business days in advance in **Greenfield Recorder** or the **Hampshire Daily Gazette**.

from the applicant or the applicant's representative.

For more information about this application or the Wetlands Protection Act, contact the Shutesbury Conservation Commission (concom@shutesbury.org or 413.259.3792) or the Department of Environmental Protection (DEP) Western Region Office at (413.784.1100). For information about the Shutesbury Wetlands Protection Bylaw, contact the Shutesbury Conservation Commission.

shutesbury.org/concom. A paper copy may be obtained, for a fee, from the Shutesbury Town Clerk: townclerk@shutesbury.org or 413.259.1204. Copies may also be obtained

AFFIDAVIT OF SERVICE

I, <u>Jeff Brandt</u>, hereby certify under the pains and penalties of perjury that on <u>October 26, 2020</u>
I gave notification to abutters in compliance with the Shutesbury Wetlands Protection Bylaw and regulations as well as the second paragraph of the Massachusetts General Laws, Chapter 131,
Section 40 and the DEP Guide to Abutter Notification in connection with the following matter:


An Abbreviated Notice of Resource Area Delineation application was filed under the Massachusetts Wetlands Protection Act by <u>W.D. Cowls, Inc.</u> with the Shutesbury Conservation Commission on <u>October 26, 2020</u> for the property located <u>off Pratt Corner Road, Shutesbury, Massachusetts (Assessor's ID ZU-2)</u>.


The form of the notification, and a list of the abutters to whom it was given and their addresses, are attached to this Affidavit of Service.

Jeff Brandt	
871	_10/26/2020
Signature	Date


ATTACHMENT D Figure 1: Delineated Resources Map (September 2020)










Coordinate System: NAD 1983 StatePlane New Hampshire FIPS 2800 Feet; Map Rotat

















Coordinate System: NAD 1983 StatePlane New Hampshire FIPS 2800 Feet; Map Rotation: 0
























Coordinate System: NAD 1983 StatePlane New Hampshire FIPS 2800 Feet; Map Rotation: 0









Coordinate System: NAD 1983 StatePlane New Hampshire FIPS 2800 Fee

